



Mini Review

The redox interplay between nitrite and nitric oxide: From the gut to the brain

Cassilda Pereira¹, Nuno R. Ferreira¹, Bárbara S. Rocha¹, Rui M. Barbosa, João Laranjinha^{*}

Department of Pharmacy and Center for Neurosciences and Cell Biology, University of Coimbra, Health Sciences Campus, Azinhaga de Santa Comba, 3000-548 Coimbra, Portugal

ARTICLE INFO

Article history:

Received 10 February 2013

Accepted 9 April 2013

Keywords:

Dietary nitrate

Nitrite

Nitric oxide

Gut

Brain

Glutamate

ABSTRACT

The reversible redox conversion of nitrite and nitric oxide (*NO) in a physiological setting is now widely accepted. Nitrite has long been identified as a stable intermediate of *NO oxidation but several lines of evidence support the reduction of nitrite to nitric oxide *in vivo*. In the gut, this notion implies that nitrate from dietary sources fuels the longstanding production of nitrite in the oral cavity followed by univalent reduction to *NO in the stomach. Once formed, *NO boosts a network of reactions, including the production of higher nitrogen oxides that may have a physiological impact via the post-translational modification of proteins and lipids. Dietary compounds, such as polyphenols, and different prandial states (secreting specific gastric mediators) modulate the outcome of these reactions. The gut has unusual characteristics that modulate nitrite and *NO redox interplay: (1) wide range of pH (neutral vs acidic) and oxygen tension (*c.a.* 70 Torr in the stomach and nearly anoxic in the colon), (2) variable lumen content and (3) highly developed enteric nervous system (sensitive to *NO and dietary compounds, such as glutamate). The redox interplay of nitrite and *NO might also participate in the regulation of brain homeostasis upon neuronal glutamatergic stimulation in a process facilitated by ascorbate and a localized and transient decrease of oxygen tension. In a way reminiscent of that occurring in the stomach, a nitrite/ *NO /ascorbate redox interplay in the brain at glutamatergic synapses, contributing to local *NO increase, may impact on *NO -mediated process.

We here discuss the implications of the redox conversion of nitrite to *NO in the gut, how nitrite-derived *NO may signal from the digestive to the central nervous system, influencing brain function, as well as a putative ascorbate-driven nitrite/ *NO pathway occurring in the brain.

© 2013 The Authors. Published by Elsevier B.V. Open access under [CC BY-NC-ND license](http://creativecommons.org/licenses/by-nc-nd/3.0/).

Contents

From nitric oxide to nitrite and back	277
Nitrite and nitric oxide in the gut: luminal warfare or welfare?	277
Reduction by dietary polyphenols	278
Protein nitrosation	278
Protein and lipid nitration	278
Redox signaling by nitrite and nitric oxide from gut mucosa and beyond	279
Mitochondrial nitric oxide–nitrite cycle	279
Hypoxic signaling	279
Hemoglobin and myoglobin	279
Xanthine oxidase (XO)	280
The enteric nervous system	280
Nitrite and nitric oxide in the central nervous system	280

^{*} Corresponding author. Tel.: +351 239488466.

E-mail address: laranjin@ci.uc.pt (J. Laranjinha).

¹ These authors contributed equally to this paper.

Basic tenets of nitric oxide in the brain: production, actions, diffusion and inactivation	280
Where nitric oxide and nitrite come together	280
Nitric oxide/ nitrite/ ascorbate cycle in the brain	281
Acknowledgments	281
References	281

From nitric oxide to nitrite and back

Nitric oxide ($\cdot\text{NO}$), an ubiquitous free radical produced via a highly regulated enzymatic process *in vivo*, modulates a wide range of signaling pathways in virtually all body organs and systems [1]. Physiological mechanisms, such as the regulation of vascular tone, host response as well as learning and memory formation are tightly regulated by this diffusible free radical [2]. Upon its synthesis *via* NO synthases (NOS), $\cdot\text{NO}$ may embark in reactions with biomolecular targets, being ultimately neutralized into the supposedly inert oxidation products nitrite and nitrate [3]. Both anions have been disregarded as inert products devoid of physiological relevance but recently it was demonstrated that nitrate and nitrite might be stepwisely reduced back to $\cdot\text{NO}$ *in vivo*.

The so-called nitrate-nitrite-NO pathway, by opposition to the classical L-arginine-NOS pathway, implies that nitrate and nitrite fuel the production of $\cdot\text{NO}$, a process particularly evident in the gastrointestinal compartment for nitrate from dietary sources originates nitrite in the oral cavity and $\cdot\text{NO}$ in the stomach [4]. The gut is therefore a primary site for the production of significantly high $\cdot\text{NO}$ concentrations [5,6]. The nitrite reduction to $\cdot\text{NO}$ has also been described in other compartments/tissues under hypoxic conditions, a condition in which several hemoproteins (e.g., myoglobin, hemoglobin, xanthine oxidase, among others) acquire a nitrite-reductase activity, ensuring $\cdot\text{NO}$ bioavailability under oxygen deprivation [7]. Taken together, these observations suggest a reciprocal relationship between nitrite and $\cdot\text{NO}$ during redox signaling mechanisms.

Nitrite and nitric oxide in the gut: luminal warfare or welfare?

The production of $\cdot\text{NO}$ from nitrate obligates the intermediary formation of nitrite [4]. Nitrate, obtained from the diet or endogenously produced upon $\cdot\text{NO}$ oxidation, is mixed with saliva in the oral cavity and travels along the gut being absorbed in the small

intestine. The salivary glands then recover *c.a.* 20% of circulating nitrate and secrete it into the oral cavity where it is reduced to nitrite by the local microflora and mixed with saliva, thus establishing the enterosalivary circulation of nitrate [8,9].

Under the acidic conditions of the gastric juice, nitrite is protonated to nitrous acid (HNO_2), which then decomposes into different nitrogen oxides, depending on the redox microenvironment and gastric content [10,11]. Under normal fasting conditions, HNO_2 yields $\cdot\text{NO}$ and nitrogen dioxide radical ($\cdot\text{NO}_2$) which may trigger signaling cascades by direct interaction with hemoproteins, such as soluble guanylate cyclase in the case of $\cdot\text{NO}$, or by oxidizing and nitrating proteins and lipids in the case of $\cdot\text{NO}_2$ [1,12]. Nitric oxide and $\cdot\text{NO}_2$ may also combine to produce dinitrogen trioxide (N_2O_3) [13], a nitrosating agent. Dinitrogen trioxide reacts with gastric chloride, phosphate and bicarbonate anions to form nitrosyl compounds that nitrosate secondary amines, a process that has been claimed to support a deleterious effect of dietary nitrite [14]. However, N_2O_3 hydrolysis to nitrite and HNO_2 is kinetically favored rather than N-nitrosation [14]. Of note, nitrite has also been shown to induce S-nitrosation within the gastric compartment, suggesting that this posttranslational modification may also be considered as part as nitrite redox signaling [15]. Finally, two $\cdot\text{NO}_2$ molecules may also combine to generate dinitrogen tetroxide (N_2O_4) which under aqueous solutions yields both nitrate and nitrite [16].

However, this scenario is expected to change after a meal. During a postprandial period, ascorbic acid is secreted together with gastric juice into the lumen and, upon reaction with nitrite, is oxidized to ascorbyl radical whereas nitrite is univalently reduced to $\cdot\text{NO}$ (Fig. 1A) [6,17]. Consequently, in the presence of endogenous reducing agents, the network of reactions is shifted towards $\cdot\text{NO}$ production [6,18,19]. This raised the question as to whether dietary products could modulate the intricate dynamics of nitrite and $\cdot\text{NO}$ in the acidic gastric lumen. Here, we highlight three mechanisms by which nitrite signals *in vivo* through redox chemistry.




Fig. 1. EPR analysis of the reaction between caffeic acid or ascorbic acid with nitrite. (A) EPR signal of ascorbate radical (line 1) and caffeic acid semiquinone radical (line 3) obtained under flow conditions upon mixing 2 mM of the compounds with 4 mM of nitrite at pH 2.0. Line 2 is the mixture of caffeic acid with nitrite alkalized immediately before being pumped to the EPR cavity. (B) *In vivo* $\cdot\text{NO}$ production in the stomach of healthy volunteers following consumption of lettuce and the dietary products and beverages indicated. Reproduced from Gago B, Lundberg JO, Barbosa RM, Laranjinha J. Red wine-dependent reduction of nitrite to nitric oxide in the stomach. *Free Radic Biol Med* 43(9):1233–42; 2007 and Rocha BS, Gago B, Barbosa RM, Laranjinha J. Dietary polyphenols generate nitric oxide from nitrite in the stomach and induce smooth muscle relaxation. *Toxicology* 265(1–2):41–8; 2009 with permission from Elsevier.

Reduction by dietary polyphenols

The paradigm defining polyphenols as global antioxidants has nowadays been largely discredited for many reasons reviewed elsewhere [20]. Yet, more than 8000 different polyphenols are provided by the human diet and their health benefits are well established from the epidemiologic viewpoint, suggesting that other mechanisms operate *in vivo* [21]. One of them relies precisely on the nitrite-NO interplay in the gastric lumen. Foods (and beverages, including the red wine) rich in polyphenols, in a way reminiscent of ascorbate, have been shown to boost $\cdot\text{NO}$ production from nitrite at acidic pH in the human stomach [6,18,19]. Upon the consumption of lettuce (as a rich source of nitrite) and of red wine, apples or black tea (as rich sources of polyphenols) the intragastric $\cdot\text{NO}$ levels in humans rise by about three orders of magnitude (Fig. 1B) [19]. Mechanistically, we have shown that dietary nitrite is univalently reduced to $\cdot\text{NO}$ while the polyphenol is oxidized to the corresponding o-semiquinone radical (Fig. 1A) [6]. As a result, two radicals are produced: $\cdot\text{NO}$ and the phenoxyl radical. They can both combine to produce nitroso derivatives or, due to the high oxygen tension in the stomach (≈ 70 Torr [22]), $\cdot\text{NO}$ may be oxidized to $\cdot\text{NO}_2$ which, in turn, can combine with the phenoxyl radicals yielding nitrated phenols [23]. Of note, the trimolecular reaction of $\cdot\text{NO}$ with oxygen, considered to be too slow to be pertinent *in vivo* ($k=8 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$) [24], acquires relevance in the gastric compartment as the steady state concentrations of both reagents are unusually high.

Although thermodynamically feasible [25], the mechanistic details of nitrite reduction to $\cdot\text{NO}$ at the expenses of polyphenols in the gastric environment still need further exploration. Overall, it is now important to recognize that the interaction of nitrite with polyphenols in the gastric lumen produces $\cdot\text{NO}$, which diffuses toward deep regions of the gastric mucosa [26,27], participating in distinct physiological mechanisms such as smooth muscle relaxation [19], regulation of mucosal blood flow and mucus thickness [28,29] and also in the eradication of pathogens such as *Helicobacter pylori* [30].

Protein nitrosation

Wine has been identified as one of the polyphenol-containing dietary sources that promote $\cdot\text{NO}$ production from nitrite in the stomach (See Fig. 1B). Along with polyphenols, ethanol is a major component of wine that can interact with nitrite under gastric conditions [31]. The reaction of aliphatic alcohols, with nitrite-derived reactive nitrogen species, such as HNO_2 and N_2O_3 may lead to the formation of alkyl nitrites of the corresponding alcohols via O-nitrosation [32]. This family of organic nitrites are known for their potent vasodilator activity [31]. Along these lines we have demonstrated that, under acidic gastric conditions, ethyl nitrite is formed in micromolar concentrations from the reaction of red wine or distilled alcoholic drinks with nitrite at concentrations that can be easily found physiologically [31]. Of note, ethyl nitrite is as a potent nitrosating agent [33] and may indirectly mediate $\cdot\text{NO}$ effects, as supported by relaxation experiments with blood vessels and gastric strips [31].

Several chemical groups other than alcohols can be nitrosated as a result of the nitrite chemistry in the stomach, namely aromatic compounds, amines and thiols, yielding C-, N-, S- nitroso species [34,35] along with heme moieties that produce heme-nitrosyls (heme-nitrosylation) [36,37]. A further level in the complex chemistry that may take place in the gastric compartment is added by the observation that different physiological/pathophysiological conditions may redirect the gastric chemistry such as, for instance, an achlorhydric stomach, where the neutral pH facilitates the formation of N-nitroso compounds (reviewed in

[38]). The biological significance of many of these modifications remains obscure but S-nitrosation has received particular attention. The posttranslational modification of a critical cysteine residue in a protein by S-nitrosation can be relevant on the regulation of protein function [39] and has been implicated in controlling oxygen delivery to tissues, modulating the function or activity of transcription factors, enzymes, membrane receptors and ion channels [37]. Mechanistically, S-nitrosation may occur between $\cdot\text{NO}$ and a thiol group, if a thyl radical is formed in the cysteine residue [40] or, more importantly, by the action nitrogen oxides (formally addition of a nitrosonium equivalent, NO^+), such as N_2O_3 formed, for instance, by the reaction of $\cdot\text{NO}$ with oxygen [41]. It may be also considered that two different thiols can undergo a fast transnitrosation reaction, which may explain in part the high liability of S-nitrosothiols [42,43].

The stomach is a proper environment for protein S-nitrosation for diet, saliva, gastric secretions (containing glutathione) and mucus glycoproteins (cysteine-rich mucins) constitute potential targets for S-nitrosation [28]. Considering the perfect timing (nitrite consumption) and place for mucins nitrosation, we have tried to elucidate the effects of the nitrite rich chemistry on the mucus layer from a physiological and pathophysiological point a view. A nitrosation pattern in the mucus proteins was found and, moreover, can be modulated with diet components and endogenous reductants (unpublished observations). It is tempting to speculate that several $\cdot\text{NO}$ -like effects of nitrite described in the stomach, such as stimulation of the mucosal blood flow and mucus generation, might proceed via intermediate formation of S-nitrosothiols which can act as stable carriers of $\cdot\text{NO}$. Finally, it must be added that many of the compounds formed in the gastric lumen from acidified nitrite are fairly stable and readily absorbed which suggests systemic effects [4].

Protein and lipid nitration

In the gastric lumen, due to the low pH (*c.a.* 2 under fasting) and unusually high nitrite concentrations (1–2 mM upon a nitrate load [44]), nitrite triggers a complex network of chemical reactions yielding oxidizing and nitrating agents. Different nitrite-dependent nitrating pathways have been pointed out but $\cdot\text{NO}_2$ seems to be the main intermediate in most of them [45,46]. In the stomach, two important sources of $\cdot\text{NO}_2$ may be forwarded: $\cdot\text{NO}$ autoxidation and peroxyntitrous acid (ONOOH) formation. As aforementioned, the high concentrations of $\cdot\text{NO}$ and oxygen achieved in the gastric lumen afford physiological significance to an apparently irrelevant reaction *in vivo* ($\cdot\text{NO}$ autoxidation). In addition, $\cdot\text{NO}$ and superoxide radical ($\text{O}_2^{\cdot-}$) combine at near limit diffusion rate ($k=6.7 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$) to produce ONOOH [47]. At physiological pH, ONOOH exists in equilibrium with peroxyntitrite anion (ONOO^-) ($\text{pKa}=6.8$) but in the acidic gastric juice ONOOH is the predominant form [48]. Peroxyntitrous acid undergoes homolytic cleavage to produce $\cdot\text{NO}_2$ and hydroxyl radical ($\cdot\text{OH}$) [49]. To this pathway aids the high carbon dioxide (CO_2) tension in the headspace and bicarbonate in the gastric juice that via the production of nitrosoperoxy carbonate (ONOOCOO) drives the formation of not only $\cdot\text{NO}_2$ but also carbonate radical ($\text{CO}_3^{\cdot-}$), which is an oxidant stronger than $\cdot\text{NO}_2$ [50,51]. This facilitates the univalent oxidation or tyrosine to tyrosil radical (tyr-O^{\cdot}) (the first step of protein tyrosine nitration) and further radical combination of tyr-O^{\cdot} and $\cdot\text{NO}_2$ ($k=3.0 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$) [51].

Given this complex web of reactions, it is likely that dietary nitrite activates oxidative signaling pathways in the stomach through protein posttranslational modifications. Accordingly, we have observed nitrite-dependent protein (tyrosine) nitration *in vivo* in the stomach [46]. We have observed that pepsin, a gastric protease involved in the breakdown of dietary and mucosal

proteins, is nitrated by both inorganic nitrite and human saliva upon a nitrate load [52]. Pepsin nitration is inhibited by urate, pointing to $\cdot\text{NO}_2$ as the nitrating agent arising from nitrite. Moreover, this posttranslational modification is associated with a decrease of the proteolytic activity of the enzyme. These results strongly support the hypothesis of nitrite, from dietary sources, to signal through nitration reactions with functional and physiological implications.

In addition to endogenous protein nitration, dietary compounds may also be targeted for nitrite-dependent nitration, notoriously dietary lipids, which have already been described to go through structural modifications upon exposure to acidified nitrite *in vitro* [53]. Therefore, lipid nitration afforded by gastric nitrite is a hypothesis that requires further investigation [46].

Taken together, these considerations support the concept that at acidic pH, protein and lipid nitration may constitute a yet poorly characterized pathway accounting for nitrite signaling *in vivo* [46].

Redox signaling by nitrite and nitric oxide from gut mucosa and beyond

Once formed in the acidic gastric lumen, and in addition to reactions with radicals, nitrogen oxides or macromolecules, $\cdot\text{NO}$ diffuses towards the gastric mucosa [26,27], hence triggering physiological responses. It increases mucus thickness (diets supplemented with nitrate have been shown to increase MUC6 expression, [54]), enhances mucosal blood flow (likely by binding to endothelial guanylate cyclase in vascular smooth muscle cells) and induces smooth muscle relaxation, influencing gastric emptying rates (reviewed in [10]). There is a vast collection of biochemical reactions in which $\cdot\text{NO}$ participates [11] that go beyond the scope of this review but we would like to emphasize that $\cdot\text{NO}$ can be also oxidized to nitrite and nitrate, contributing to a pool of these anions *in vivo* [55]. However, the documented increase of plasma and tissue nitrate and nitrite upon a nitrate load [56] supports a dietary contribution to the *in vivo* pool of those anions via, for instance, the enterosalivary recirculation.

In tissues, nitrite may trigger distinct signaling pathways, either via $\cdot\text{NO}$ production or in a $\cdot\text{NO}$ -independent manner. An interesting example of the later is myeloperoxidase (MPO), that can use nitrite as a substrate to produce oxidizing (compound I and II) and nitrating ($\cdot\text{NO}_2$) agents [57]. This reaction is of particular interest throughout the gut mucosa due to the continuous recruitment of polymorphonuclear cells that express this peroxidase as part of their antimicrobial arsenal [58]. Therefore, nitrite may be expected to trigger protein or lipid nitration reactions (at pH 7.4) through MPO activation.

Below, we will briefly address some of nitrite-dependent signaling pathways.

Mitochondrial nitric oxide-nitrite cycle

Nitric oxide is known to regulate mitochondrial function through mechanisms that include mitochondrial biogenesis [59], reactive oxygen species production [60], cytochrome c release [61] and inhibition of mitochondrial Complex I (NADH-ubiquinone oxidoreductase) and Complex IV (cytochrome c oxidase, CcOX) [62]. CcOX, in turn, is endowed with a nitrite-reductase activity [63,64] as well with a nitric oxide oxidase activity, particularly under conditions of low electron flux and in the presence of oxygen (reviewed in [65]), indicating that the enzyme contributes to nitrite physiological levels. In connection with these observations it is interesting to note that recently, nitrite was shown to regulate mitochondrial biogenesis in a $\cdot\text{NO}$ - and soluble guanylate cyclase-independent manner [66].

Complex I can be inhibited by $\cdot\text{NO}$ via S-nitrosation after a long exposure to high $\cdot\text{NO}$ concentrations [67] and crucially depends on the structural conformation of the enzyme which, in turn, depends on the availability of oxygen and NADH. Under hypoxic conditions, Complex I undergoes a transition between the active form and the de-activated form, exposing a critical cysteine residue that when S-nitrosated fully inhibits the enzyme [68,69]. Apparently, during re-oxygenation, the inhibited S-nitrosated Complex I has a fundamental contribution for cell protection, preventing the accumulation of reducing equivalents and ROS (reactive oxygen species) production [70]. Regarding the inhibition of complex IV (CcOX) by $\cdot\text{NO}$ and related species, two mechanisms have generated consensus [71,72]. A first reaction pathway involves the formation of a NO-bound (nitrosyl) enzyme in the ferrous heme a_3 center and a second one produces nitrite-bound CcOX on the ferric heme a_3 center. Both pathways lead to the reversible inhibition of the enzyme, and CcOX can restore its activity via dissociation of both, $\cdot\text{NO}$, the process being slow and light-sensitive, and nitrite, the process being more rapid and light-sensitive. Sarti and co-workers, showed that both mechanisms can occur. Under conditions of low electron flux and high oxygen concentration, the 'nitrite' uncompetitive inhibition pathway prevails, whereas the electron flux is increased, and the oxygen concentration reduced, the oxygen-competitive 'nitrosyl' pathway tends to take over [71,72]. Recent evidences in favor of a nitrite reductase activity of CcOX [73] under acidic conditions and low oxygen tension, producing $\cdot\text{NO}$, support that such a nitrite-reductase activity plays a role in hypoxic cell signaling.

Taken together, the reaction of mitochondrial complexes I and IV with $\cdot\text{NO}$ may trigger physiological or pathological events [67,74] and it is of note that $\cdot\text{NO}$ bioavailability is intimately related to the oxygen tension for conditions in which oxygen decreases in tissues below normoxic conditions, the activity of NOSs to produce $\cdot\text{NO}$ is progressively slowed. However, the fact that the anoxic environment promotes tissue acidification, favouring the reduction of nitrite to $\cdot\text{NO}$, highlights the importance of the tissue/cell distribution of nitrite concerning the $\cdot\text{NO}$ chemistry [4,75].

Hypoxic signaling

The nitrite-dependent redox signaling under low oxygen tension is largely dependent on its reduction to $\cdot\text{NO}$ by several enzymes that acquire nitrite-reductase activity under these circumstances (for a recent review see [7]).

Hemoglobin and myoglobin

Several enzymes ensure a nitrite-reductase activity along the oxygen gradient but deoxygenated hemoglobin (plasma) and myoglobin (tissues) are of foremost importance. Although at different rates, both proteins may participate in the production $\cdot\text{NO}$ by the same mechanism [76]. Nitrite interacts with the heme group of deoxyhemoglobin [deoxyHb(Fe^{2+})], producing $\cdot\text{NO}$ and methemoglobin [methHb(Fe^{3+})] [77,78] at rates that increase when the protein conformation is changing from the R-to-T state (from oxy to deoxy), being maximized when hemoglobin is 50% saturated with oxygen [79]. It has been described that $\cdot\text{NO}$ hence formed in the erythrocyte may originate N_2O_3 via its combination with $\cdot\text{NO}_2$ which is present as a ferrous nitrogen dioxide [$\cdot\text{NO}_2(\text{Fe}^{2+})$] in equilibrium with [$\text{NO}_2^-(\text{Fe}^{3+})$] [80]. The functional significance of this reaction is that N_2O_3 is much more stable than $\cdot\text{NO}$ in the heme environment and therefore can escape the erythrocyte and may then transduce $\cdot\text{NO}$ signal in the vessel or endothelial compartment via, for instance, its ability to nitrosate thiols [80]. Likewise, deoxymyoglobin acts as a nitrite reductase,

reducing nitrite to $\cdot\text{NO}$ faster than deoxyhemoglobin due to its lower heme redox potential. This mechanism underlies some of the mitochondrial effects of nitrite, as myoglobin:nitrite-derived $\cdot\text{NO}$, being produced in proximity to mitochondria, inhibits CcOX and regulates mitochondrial respiration [81]. Such an effect on oxygen bioavailability has been shown to positively impact in the performance of healthy volunteers under submaximal exercise upon dietary supplementation with nitrate [82]. In the heart, under hypoxic conditions, $\cdot\text{NO}$ production via deoxymyoglobin reduction of nitrite, downregulates the cardiac energy status, oxygen consumption and tissue contractility [83], highlighting the homeostatic effects of nitrite under particularly disfavored conditions.

Xanthine oxidase (XO)

Besides the catabolism of purines and pyrimidines, XO is also able to reduce oxygen to $\text{O}_2^{\cdot-}$ and hydrogen peroxide (H_2O_2). However, it can also reduce nitrite to $\cdot\text{NO}$ at the molybdenum site under hypoxic conditions in the presence of NADH as electron donor [84]. The reaction is obviously affected by the tissue levels of the enzyme but, additionally, its activity is critically controlled by nitrite that, given the high value of K_m ($k_m=2.5$ mM), acts as the limiting substrate [7]. Therefore, since the K_m for nitrite exceed the typical levels on nitrite by two orders of magnitude, it is likely that only when nitrite levels rise above basal (such as when the gastric mucosa is exposed to intense $\cdot\text{NO}$ fluxes upon a nitrate load) this reaction may acquire physiological relevance. Alternatively, under ischemic conditions, when the pH drops and nitrite is converted into HNO_2 , $\cdot\text{NO}$ production by this pathway is favored [7] and has been suggested to protect against myocardial infarction [85]. Interestingly, for 70 μM oxygen, XO can univalently reduce both, oxygen to $\text{O}_2^{\cdot-}$ and nitrite to $\cdot\text{NO}$, and further considering that both radicals can combine to produce ONOO^- under normoxic physiological conditions [86] one may envisage that nitrite may also be implicated in ONOO^- biological impact. Peroxynitrite is a short-lived oxidant implicated in a wide range of diseases [87] and therefore nitrite can be regarded as a Janus-faced molecule because depending on the local microenvironment (e.g., oxygen levels) it can activate protective pathways against ischemia or trigger the formation of deleterious oxidants.

The enteric nervous system

Gastrointestinal mucosa houses submucosal and myenteric plexuses that integrate gut and nervous functions, bridging the enteric and central nervous systems [88]. Diet composition modulates not only gastrointestinal homeostasis but also the gut-brain signaling through diverse mechanisms. For instance, there is an operative *amino acid sensing system* within the gastric mucosa that recognizes dietary glutamate [89]. Albeit being the major excitatory neurotransmitter in the central nervous system, this amino acid also modulates important gut functions such as secretion, motility and metabolism [90] via mechanisms that include the generation of $\cdot\text{NO}$ and serotonin [91]. Given the high steady state concentrations of $\cdot\text{NO}$ achieved in the gastric lumen and mucosa upon nitrite reduction, glutamate-dependent pathways in the gastrointestinal tract are likely to be modulated by dietary nitrate.

Nitrite and nitric oxide in the central nervous system

Presley et al. have shown that a high nitrate diet increases the blood flow in specific brain areas in older subjects [92], introducing the notion that nitrate available from the diet may impact on brain homeostasis. However, the mechanistic details that bridge dietary nitrate and brain function remain largely elusive. As

aforementioned, upon a nitrate load there is an increase of both nitrate and nitrite concentration in blood and tissues, but how (and if) is nitrite converted into $\cdot\text{NO}$ in the brain and what is the impact in brain functions such as cognition and memory formation remains to be elucidated.

Given the critical role of $\cdot\text{NO}$ in the brain, a particular challenging question is whether one can identify a brain selective or favored mechanism that might affect the redox conversion between nitrite and $\cdot\text{NO}$ and, thereby $\cdot\text{NO}$ -mediated process. When considering the extremely high concentration of ascorbate in neurons and the occurrence of glutamate:ascorbate exchangers it is tempting to speculate that ascorbate might, under conditions of excitatory stimuli, participate in nitrite-dependent $\cdot\text{NO}$ production in the brain. Before addressing this hypothesis we will briefly review biochemical and physiological features of $\cdot\text{NO}$ in the brain.

Basic tenets of nitric oxide in the brain: production, actions, diffusion and inactivation

Upon neuronal activation, $\cdot\text{NO}$ production in the brain is related to glutamatergic activity, involving the activation of ionotropic glutamate receptors (iGluR), particularly NMDA subtype, and the consequent influx of Ca^{2+} that activates neuronal isoform of NOS (nNOS), thus promoting the conversion of L-arginine to L-citrulline and $\cdot\text{NO}$, in an oxygen dependent way [93,94]. Following its production, $\cdot\text{NO}$, among other targets, may activate soluble guanylate cyclase, [95–98], triggering physiological responses that are tissue and cell specific and include neuronal excitability, synaptic plasticity, modulation of neurotransmitter release, learning and memory processes and neurovascular coupling [93,98–100].

It is of note that $\cdot\text{NO}$ rapidly diffuses in the brain across cell membranes with a coefficient of 2.2×10^{-5} cm^2/s [101], interacting with multiple targets, especially heme and other iron proteins at relatively fast reactions (in the case of myo and hemoglobin rate constants are of the order of 10^7 $\text{M}^{-1} \text{s}^{-1}$) [102]. In the absence of specific interaction with receptors, $\cdot\text{NO}$ actions are dependent on its spatiotemporal profile which, in turn, is dependent on the balance between its synthesis and inactivation. Regarding the latter process, on basis of selective microelectrodes stereotaxically inserted in the rat brain *in vivo* [103–107], we have recently shown that $\cdot\text{NO}$ has a half life of 0.64 s in rat cortex [101] and that scavenging by erythrocytes represents the major inactivation pathway in the brain.

Where nitric oxide and nitrite come together

Nitric oxide synthesized by the neuronal isoform of NOS (nNOS) can not only regulate oxygen availability in tissues (via mitochondrial actions described above) but also bridges neuronal activity with local changes of blood flow in the microcirculation of the brain, a process known as neurovascular coupling [108]. Under hypoxia, it is expectable that the activity of NOS (and, in particular nNOS) is highly attenuated and, consequently, we would expect that the decreased production of $\cdot\text{NO}$ translates into subsequent impairment of blood flow and compromised vasodilation. However, apparently, NOS inactivation during hypoxia does not block vasodilation [109]. This observation adds to several others, including the viability of mice lacking all three isoforms of NOS [110], to suggest the occurrence of alternative and/or complementary pathways to ensure that tissues get the needed amount of $\cdot\text{NO}$, regardless of the functional state of the enzymes usually involved in its synthesis. In this scenario, the possibility that dietary nitrite/nitrate may contribute to basal levels of $\cdot\text{NO}$ signaling emerges as

a plausible possibility yet poorly investigated in the brain. This interaction becomes particularly relevant in maintaining the balance between $\cdot\text{NO}$ production, oxygen consumption and neurovascular coupling in the brain. Nitrite has shown to increase blood flow preferentially in hypoxic conditions, allowing blood flow to increase precisely where it is needed most [111–120]. This shift from the classical view of nitrite as a simple by product of $\cdot\text{NO}$ oxidation to a molecule with physiological impact, as discussed ahead, is associated with redox mechanisms leading to $\cdot\text{NO}$ formation when enzymatic production is somehow compromised.

Based on preliminary data and on the rationale discussed ahead we will propose a critical role for ascorbate in $\cdot\text{NO}$ biochemistry in the brain via the univalent reduction of nitrite.

Nitric oxide/ nitrite/ ascorbate cycle in the brain

Ascorbate is a powerful reducing agent highly concentrated in the brain, both in the extracellular fluid and cells, namely neurons and astrocytes. Ascorbate is present at even higher concentrations in areas such as the hippocampus where $\cdot\text{NO}$ has been shown to play critical roles in learning and memory processes [121–123]. Ascorbate release from neuronal cells to the extracellular fluid occurs in connection with glutamatergic activity, and a putative glutamate/ascorbate heteroexchange mechanism has been proposed to regulate this flow outwards the cells following stimulation [124]. However, other reports failed to find a direct correlation between glutamatergic uptake and ascorbate release [125,126].

Nitric oxide undergoes oxidation to nitrite and this amount adds to nitrite received from the diet, yielding a reservoir in the extracellular fluid. As discussed before, the reduction of nitrite promoted by univalent reductants such as dietary polyphenols and ascorbate is now well established in the gastric environment where both reagents (nitrite and ascorbate) achieve optimal conditions (low pH, high concentration) for the reduction of nitrite to $\cdot\text{NO}$ occur.

Years ago, Millar elegantly described an alternative mechanism for $\cdot\text{NO}$ production in the brain completely independent of enzymatic control, involving the reduction of nitrite to $\cdot\text{NO}$ by ascorbate released from neurons during increased neuronal activity [109]. In this regard, it is of note the millimolar range of ascorbate concentration in neurons and also that ascorbate, which is released following neuronal activity, could contribute to convert nitrite to $\cdot\text{NO}$, thus promoting beneficial vasodilator actions on blood vessels in the brain in the absence of an enzymatic and oxygen requiring process, namely the NOS-catalyzing process. This would be particularly important in hypoxia, because (a) the low of oxygen compromises NOS activity and (b) during hypoxia, local pH may drop to values near 6.4 [127] and, as depicted in Fig. 2, when ascorbate and nitrite are in solution and pH drops from 7.4 to values near 6.5 $\cdot\text{NO}$ production occurs.

However, considering that under glutamatergic stimulus local oxygen drops to a very low tension, reaching a “hypoxia like” transient status, it can be hypothesized that during this time period pH will drop sufficiently to promote nitrite reduction to $\cdot\text{NO}$ by the ascorbate hence released during the stimulation. In agreement with this hypothesis preliminary experiments have shown a temporal correlation between $\cdot\text{NO}$ and ascorbate dynamics upon glutamate stimulation of rat hippocampus *in vivo* (unpublished data), suggesting that ascorbate is contributing to $\cdot\text{NO}$ signals upon glutamatergic stimulation.

The basal nitrite concentration in the extracellular space of hippocampus is not accurately known and is likely to vary as a function of the redox environment but on basis of a microdialysis approach in connection with a chemiluminescence measurement of nitrite-derived nitric oxide a range between 50 and 400




Fig. 2. Nitric oxide production from a mixture of ascorbate and nitrite as a function of pH. The reaction vessel was supplemented with 250 μM of ascorbate and 20 μM of nitrite at a pH of 7.4. The vessel was coupled to an $\text{amiNO}^{\text{®}}$ sensor (selective for $\cdot\text{NO}$ measurements) and to a pH meter. Current was allowed to stabilize. Then, pH was slowly acidified until it reached the value of 6.51 (a value that can be reached *in vivo* under hypoxic conditions). At this pH value, current increased indicating that $\cdot\text{NO}$ was being produced. Maximum concentration of $\cdot\text{NO}$ obtained in these conditions was 118 nM.

nanomolar was routinely found in male Wistar rats found [128]. On the other hand, nitrite concentration in the cerebrospinal fluid (of humans and rats), was reported to be *circa* 1 μM [129,130]. Under these conditions, the increase of $\cdot\text{NO}$ and ascorbate concentration in the extracellular space concurrently with a drop on the local pH (since oxygen concentration also drops) might set the conditions for nitrite reduction to $\cdot\text{NO}$. The physiological outcome is that nitrite/ $\cdot\text{NO}$ /ascorbate redox interplay could contribute to $\cdot\text{NO}$ -mediated processes such as the neurovascular coupling and the subsequent blood flow increase.

In summary, the redox couple nitrite: $\cdot\text{NO}$ in the brain might acquire relevance under physiological conditions in which the stimulation of glutamatergic terminals sets the environment and facilitates the chemical conditions for nitrite univalent reduction to $\cdot\text{NO}$ by ascorbate in a way reminiscent of what occurs in the stomach. The notion of nitrite as $\cdot\text{NO}$ reservoir acquires a particular interest in the brain because $\cdot\text{NO}$ production is achieved under physiological process, without the need to propose hypoxic or ischemic conditions for $\cdot\text{NO}$ production. However, although challenging, the hypothesis of the nitrite/ $\cdot\text{NO}$ /ascorbate redox interplay with functional consequences in the brain still needs to be robustly substantiated.

Acknowledgments

This work was supported by grants PTDC/AGR-ALI/115744/2009, PTDC/SAU-NEU/108992/2008 and PTDC/SAU-NEU/103538/2008 from Fundação para a Ciência e Tecnologia (Portugal) and COMPETE. Grant PEst-C/SAU/LA0001/2011 from CNC is also acknowledged.

References

- [1] S. Moncada, R.M. Palmer, E.A. Higgs, Nitric oxide: physiology, pathophysiology, and pharmacology, *Pharmacological Reviews* 43 (2) (1991) 109–142.
- [2] S. Moncada, Nitric oxide: discovery and impact on clinical medicine, *Journal of the Royal Society of Medicine* 92 (4) (1999) 164–169.
- [3] S. Moncada, A. Higgs, The L-arginine-nitric oxide pathway, *The New England Journal of Medicine* 329 (27) (1993) 2002–2012.
- [4] J.O. Lundberg, E. Weitzberg, M.T. Gladwin, The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics, *Nature Reviews Drug Discovery* 7 (2) (2008) 156–167.
- [5] J.O. Lundberg, E. Weitzberg, J.M. Lundberg, K. Alving, Intra-gastric nitric oxide production in humans: measurements in expelled air, *Gut* 35 (11) (1994) 1543–1546.

- [6] B. Gago, J.O. Lundberg, R.M. Barbosa, J. Laranjinha, Red wine-dependent reduction of nitrite to nitric oxide in the stomach, *Free Radical Biology and Medicine* 43 (9) (2007) 1233–1242.
- [7] E. van Faassen, S. Bahrami, M. Feelisch, N. Hogg, M. Kelm, D. Kim-Shapiro, A. Kozlov, H. Li, J. Lundberg, R. Mason, H. Nohl, T. Rassaf, A. Samouilov, A. Slama-Schwok, S. Shiva, A. Vanin, E. Weitzberg, J. Zweier, M. Gladwin, Nitrite as regulator of hypoxic signaling in mammalian physiology, *Medicinal Research Reviews* 29 (5) (2009) 683–741.
- [8] D.A. Wagner, D.S. Schultz, W.M. Deen, V.R. Young, S.R. Tannenbaum, Metabolic fate of an oral dose of ¹⁵N-labeled nitrate in humans: effect of diet supplementation with ascorbic acid, *Cancer Research* 43 (4) (1983) 1921–1925.
- [9] B. Spiegelhalter, G. Eisenbrand, R. Preussmann, Influence of dietary nitrate on nitrite content of human saliva: possible relevance to *in vivo* formation of N-nitroso compounds, *Food and Cosmetics Toxicology* 14 (6) (1976) 545–548.
- [10] B.S. Rocha, B. Gago, C. Pereira, R.M. Barbosa, S. Bartesaghi, J.O. Lundberg, R. Radi, J. Laranjinha, Dietary nitrite in nitric oxide biology: a redox interplay with implications for pathophysiology and therapeutics, *Current Drug Targets* 12 (9) (2011) 1351–1363.
- [11] J.O. Lundberg, E. Weitzberg, Biology of nitrogen oxides in the gastrointestinal tract, *Gut* (2012).
- [12] H. Rubbo, R. Radi, Protein and lipid nitration: role in redox signaling and injury, *Biochimica Biophysica Acta* 1780 (11) (2008) 1318–1324.
- [13] S. Archer, Measurement of nitric oxide in biological models, *FASEB Journal* 7 (2) (1993) 349–360.
- [14] J.L. Caulfield, S.P. Singh, J.S. Wishnok, W.M. Deen, S.R. Tannenbaum, Bicarbonate inhibits N-nitrosation in oxygenated nitric oxide solutions, *Journal of Biological Chemistry* 271 (42) (1996) 25859–25863.
- [15] G. Richardson, S.L. Hicks, S. O'Byrne, M.T. Frost, K. Moore, N. Benjamin, G. M. McKnight, The ingestion of inorganic nitrate increases gastric S-nitrosothiol levels and inhibits platelet function in humans, *Nitric Oxide* 7 (1) (2002) 24–29.
- [16] M.P. Doyle, J.W. Hoekstra, Oxidation of nitrogen oxides by bound dioxygen in hemoproteins, *Journal of Inorganic Biochemistry* 14 (4) (1981) 351–358.
- [17] S.R. Tannenbaum, J.S. Wishnok, C.D. Leaf, Inhibition of nitrosamine formation by ascorbic acid, *The American Journal of Clinical Nutrition* 53 (Suppl. 1) (1991) 247S–250S.
- [18] L. Peri, D. Pietraforte, G. Scorza, A. Napolitano, V. Fogliano, M. Minetti, Apples increase nitric oxide production by human saliva at the acidic pH of the stomach: a new biological function for polyphenols with a catechol group? *Free Radical Biology and Medicine* 39 (5) (2005) 668–681.
- [19] B.S. Rocha, B. Gago, R.M. Barbosa, J. Laranjinha, Dietary polyphenols generate nitric oxide from nitrite in the stomach and induce smooth muscle relaxation, *Toxicology* 265 (1–2) (2009) 41–48.
- [20] C.G. Fraga, Plant polyphenols: how to translate their *in vitro* antioxidant actions to *in vivo* conditions, *IUBMB Life* 59 (4–5) (2007) 308–315.
- [21] C.G. Fraga, M. Galleano, S.V. Verstraeten, P.I. Oteiza, Basic biochemical mechanisms behind the health benefits of polyphenols, *Molecular Aspects of Medicine* 31 (6) (2010) 435–445.
- [22] G. He, R.A. Shankar, M. Chzhan, A. Samouilov, P. Kuppasamy, J.L. Zweier, Noninvasive measurement of anatomic structure and intraluminal oxygenation in the gastrointestinal tract of living mice with spatial and spectral EPR imaging, *Proceedings of the National Academy of Sciences of the United States of America* 96 (8) (1999) 4586–4591.
- [23] A.S. Pannala, A.R. Mani, C.A. Rice-Evans, K.P. Moore, pH-dependent nitration of para-hydroxyphenylacetic acid in the stomach, *Free Radical Biology and Medicine* 41 (6) (2006) 896–901.
- [24] M.S. Ramezani, S. Padmaja, W.H. Koppenol, Nitration and hydroxylation of phenolic compounds by peroxynitrite, *Chemical Research in Toxicology* 9 (1) (1996) 232–240.
- [25] S.V.S.S. Jovanovic, M. Simic, Y. Hara, Antioxidant properties of flavonoids: reduction potentials and electron transfer reactions of flavonoid radicals, in: C.A. Rice-Evans, L. Packer (Eds.), *Flavonoids in Health and Disease*, Marcel Dekker, New York, 1998, pp. 137–161.
- [26] K. Asanuma, K. Iijima, H. Sugata, S. Ohara, T. Shimosegawa, T. Yoshimura, Diffusion of cytotoxic concentrations of nitric oxide generated luminally at the gastro-oesophageal junction of rats, *Gut* 54 (8) (2005) 1072–1077.
- [27] B.S. Rocha, B. Gago, R.M. Barbosa, J. Laranjinha, Diffusion of nitric oxide through the gastric wall upon reduction of nitrite by red wine: physiological impact, *Nitric Oxide* 22 (3) (2010) 235–241.
- [28] H.H. Bjorne, J. Petersson, M. Phillipson, E. Weitzberg, L. Holm, J.O. Lundberg, Nitrite in saliva increases gastric mucosal blood flow and mucus thickness, *Journal of Clinical Investigation* 113 (1) (2004) 106–114.
- [29] J. Petersson, M. Phillipson, E.A. Jansson, A. Patzak, J.O. Lundberg, L. Holm, Dietary nitrate increases gastric mucosal blood flow and mucosal defense, *American Journal of Physiology—Gastrointestinal and Liver Physiology* 292 (3) (2007) G718–G724.
- [30] R.S. Dykhuizen, A. Fraser, H. McKenzie, M. Golden, C. Leifert, N. Benjamin, *Helicobacter pylori* is killed by nitrite under acidic conditions, *Gut* 42 (3) (1998) 334–337.
- [31] B. Gago, T. Nystrom, C. Cavaleiro, B.S. Rocha, R.M. Barbosa, J. Laranjinha, J. O. Lundberg, The potent vasodilator ethyl nitrite is formed upon reaction of nitrite and ethanol under gastric conditions, *Free Radical Biology and Medicine* 45 (4) (2008) 404–412.
- [32] A.D. Allen, Studies in the hydrolysis and alcoholysis of some organic nitrites, *Journal of the Chemical Society* (1954) 1968–1974.
- [33] J.O. Lundberg, E. Weitzberg, S. Shiva, M.T. Gladwin, The Nitrate–Nitrite–Nitric Oxide Pathway in Mammals, in: N.S. Bryan, J. Loscalzo (Eds.), *Nitrite and Nitrate in Human Health and Disease*, Humana Press, 2011, pp. 21–48.
- [34] A.R. Butler, P. Rhodes, Chemistry, analysis, and biological roles of S-nitrosothiols, *Analytical Biochemistry* 249 (1) (1997) 1–9.
- [35] K.M. Wink DAM, M.G. Espey, J.B. Mitchell, M.B. Grisham, J. Fukuto, M. Feelisch, The Chemical Biology of Nitric Oxide. Balancing Nitric Oxide with Oxidative and Nitrosative Stress. In: S.B. Heidelberg, editor, *Mayer*, B2000, pp. 7–29.
- [36] J.R. Lancaster Jr., J.B. Hibbs Jr., EPR demonstration of iron-nitrosyl complex formation by cytotoxic activated macrophages, *Proceedings of the National Academy of Sciences USA* 87 (3) (1990) 1223–1227.
- [37] F.L. Jour'd'heil, A.M. Lowery, E.M. Melton, S. Mnaimneh, N.S. Bryan, B. O. Fernandez, J.H. Park, C.E. Ha, N.V. Bhagavan, M. Feelisch, D. Jour'd'heil, Redox-sensitivity and site-specificity of S- and N- denitrosation in proteins, *PLoS One* 5 (12) e14400 5 (12) (2010).
- [38] N.S. Bryan, D.D. Alexander, J.R. Coughlin, A.L. Milkowski, P. Boffetta, Ingested nitrate and nitrite and stomach cancer risk: An updated review, *Food and Chemical Toxicology* 50 (10) (2012) 3646–3665.
- [39] J.S. Stamler, S. Lamas, F.C. Fang, Nitrosylation, the prototypic redox-based signaling mechanism, *Cell* 106 (6) (2001) 675–683.
- [40] A. Martinez-Ruiz, S. Lamas, Signaling by NO-induced protein S-nitrosylation and S-glutathionylation: convergences and divergences, *Cardiovascular Research* 75 (2) (2007) 220–228.
- [41] D.A. Wink, J.F. Darbyshire, R.W. Nims, J.E. Saavedra, P.C. Ford, Reactions of the bioregulatory agent nitric oxide in oxygenated aqueous media: determination of the kinetics for oxidation and nitrosation by intermediates generated in the NO/O₂ reaction, *Chemical Research in Toxicology* 6 (1) (1993) 23–27.
- [42] R. Rossi, L. Lusini, F. Giannerini, D. Giustarini, G. Lungarella, P. Di Simplicio, A method to study kinetics of transnitrosation with nitrosoglutathione: reactions with hemoglobin and other thiols, *Analytical Biochemistry* 254 (2) (1997) 215–220.
- [43] Z. Liu, M.A. Rudd, J.E. Freedman, J. Loscalzo, S-Transnitrosation reactions are involved in the metabolic fate and biological actions of nitric oxide, *Journal of Pharmacology and Experimental Therapeutics* 284 (2) (1998) 526–534.
- [44] G.M. McKnight, L.M. Smith, R.S. Drummond, C.W. Duncan, M. Golden, N. Benjamin, Chemical synthesis of nitric oxide in the stomach from dietary nitrate in humans, *Gut* 40 (2) (1997) 211–214.
- [45] H. Ischiropoulos, Biological tyrosine nitration: a pathophysiological function of nitric oxide and reactive oxygen species, *Archives of Biochemistry and Biophysics* 356 (1) (1998) 1–11.
- [46] B.S. Rocha, B. Gago, R.M. Barbosa, J.O. Lundberg, R. Radi, J. Laranjinha, Intra-gastric nitration by dietary nitrite: implications for modulation of protein and lipid signaling, *Free Radical Biology and Medicine* 52 (3) (2012) 693–698.
- [47] J.S. Beckman, Oxidative damage and tyrosine nitration from peroxynitrite, *Chemical Research in Toxicology* 9 (5) (1996) 836–844.
- [48] R. Radi, J.S. Beckman, K.M. Bush, B.A. Freeman, Peroxynitrite oxidation of sulfhydryls. The cytotoxic potential of superoxide and nitric oxide, *Journal of Biological Chemistry* 266 (7) (1991) 4244–4250.
- [49] J.S. Beckman, T.W. Beckman, J. Chen, P.A. Marshall, B.A. Freeman, Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide, *Proceedings of the National Academy of USA* 87 (4) (1990) 1620–1624.
- [50] A. Denicola, B.A. Freeman, M. Trujillo, R. Radi, Peroxynitrite reaction with carbon dioxide/bicarbonate: kinetics and influence on peroxynitrite-mediated oxidations, *Archives of Biochemistry and Biophysics* 333 (1) (1996) 49–58.
- [51] R. Radi, Nitric oxide, oxidants, and protein tyrosine nitration, *Proceedings of the National Academy of USA* 101 (12) (2004) 4003–4008.
- [52] B.S. Rocha, B. Gago, R.M. Barbosa, J.O. Lundberg, G.E. Mann, R. Radi, J. Laranjinha, Pepsin is nitrated in the rat stomach, acquiring antiulcerogenic activity: a novel interaction between dietary nitrate and gut proteins, *Free Radical Biology and Medicine* 8 (2012) 26–34.
- [53] A. Napolitano, L. Panzella, M. Savarese, R. Sacchi, I. Giudicianni, L. Paolillo, M. d'Ischia, Acid-induced structural modifications of unsaturated fatty acids and phenolic olive oil constituents by nitrite ions: a chemical assessment, *Chemical Research in Toxicology* 17 (10) (2004) 1329–1337.
- [54] E.A. Jansson, J. Petersson, C. Reinders, T. Sobko, H. Bjorne, M. Phillipson, E. Weitzberg, L. Holm, J.O. Lundberg, Protection from nonsteroidal anti-inflammatory drug (NSAID)-induced gastric ulcers by dietary nitrate, *Free Radical Biology and Medicine* 42 (4) (2007) 510–518.
- [55] L.J. Ignarro, J.M. Fukuto, J.M. Griscavage, N.E. Rogers, R.E. Byrns, Oxidation of nitric oxide in aqueous solution to nitrite but not nitrate: comparison with enzymatically formed nitric oxide from L-arginine, *Proceedings of the National Academy of USA* 90 (17) (1993) 8103–8107.
- [56] M. Govoni, E.A. Jansson, E. Weitzberg, J.O. Lundberg, The increase in plasma nitrite after a dietary nitrate load is markedly attenuated by an antibacterial mouthwash, *Nitric Oxide* 19 (4) (2008) 333–337.
- [57] U. Burner, P.G. Furtmuller, A.J. Kettle, W.H. Koppenol, C. Obinger, Mechanism of reaction of myeloperoxidase with nitrite, *Journal of Biological Chemistry* 275 (27) (2000) 20597–20601.
- [58] J.L. Wallace, Prostaglandins, NSAIDs, and gastric mucosal protection: why doesn't the stomach digest itself? *Physiological Reviews* 88 (4) (2008) 1547–1565.
- [59] E. Nisoli, S. Falcone, C. Tonello, V. Cozzi, L. Palomba, M. Fiorani, A. Piscanti, S. Brunelli, A. Cardile, M. Francolini, O. Cantoni, M.O. Carruba, S. Moncada,

- E. Clementi, Mitochondrial biogenesis by NO yields functionally active mitochondria in mammals, *Proceedings of the National Academy of USA* 101 (47) (2004) 16507–16512.
- [60] J.J. Poderoso, M.C. Carreras, C. Lisdero, N. Riobo, F. Schopfer, A. Boveris, Nitric oxide inhibits electron transfer and increases superoxide radical production in rat heart mitochondria and submitochondrial particles, *Archives of Biochemistry and Biophysics* 328 (1) (1996) 85–92.
- [61] P.S. Brookes, E.P. Salinas, K. Darley-Usmar, J.P. Eiserich, B.A. Freeman, V.M. Darley-Usmar, P.G. Anderson, Concentration-dependent effects of nitric oxide on mitochondrial permeability transition and cytochrome c release, *Journal of Biological Chemistry* 275 (27) (2000) 20474–20479.
- [62] G.C. Brown, C.E. Cooper, Nanomolar concentrations of nitric oxide reversibly inhibit synaptosomal respiration by competing with oxygen at cytochrome oxidase, *FEBS Letters* 356 (2–3) (1994) 295–298.
- [63] P.R. Castello, D.K. Woo, K. Ball, J. Wojcik, L. Liu, R.O. Poyton, Oxygen-regulated isoforms of cytochrome c oxidase have differential effects on its nitric oxide production and on hypoxic signaling, *Proceedings of the National Academy of Sciences USA* 105 (24) (2008) 8203–8208.
- [64] R.O. Poyton, P.R. Castello, K.A. Ball, D.K. Woo, N. Pan, Mitochondria and hypoxic signaling, *Annals of the New York Academy of Sciences* 1177 (1) (2009) 48–56.
- [65] P. Sarti, E. Forte, D. Mastronicola, A. Giuffrè, M. Arese, Cytochrome c oxidase and nitric oxide in action: molecular mechanisms and pathophysiological implications, *Biochimica Biophysica Acta* 1817 (4) (2012) 610–619.
- [66] L. Mo, Y. Wang, L. Geary, C. Corey, M.J. Alef, D. Beer-Stolz, B.S. Zuckerbraun, S. Shiva, Nitrite activates AMP kinase to stimulate mitochondrial biogenesis independent of soluble guanylate cyclase, *Free Radical Biology and Medicine* 53 (7) (2012) 1440–1450.
- [67] E. Clementi, G.C. Brown, M. Feelisch, S. Moncada, Persistent inhibition of cell respiration by nitric oxide: crucial role of S-nitrosylation of mitochondrial complex I and protective action of glutathione, *Proceedings of the National Academy of Sciences USA* 95 (13) (1998) 7631–7636.
- [68] A. Galkin, S. Moncada, S-nitrosation of mitochondrial complex I depends on its structural conformation, *Journal of Biological Chemistry* 282 (52) (2007) 37448–37453.
- [69] E.V. Gavrikova, A.D. Vinogradov, Active/de-active state transition of the mitochondrial complex I as revealed by specific sulfhydryl group labeling, *FEBS Letters* 455 (1–2) (1999) 36–40.
- [70] J.L. Zweier, A. Samouilov, P. Kuppusamy, Non-enzymatic nitric oxide synthesis in biological systems, *Biochimica Biophysica Acta* 1411 (2–3) (1999) 250–262.
- [71] P. Sarti, A. Giuffrè, E. Forte, D. Mastronicola, M.C. Barone, M. Brunori, Nitric oxide and cytochrome c oxidase: mechanisms of inhibition and NO degradation, *Biochemical and Biophysical Research Communications* 274 (1) (2000) 183–187.
- [72] D. Mastronicola, M.L. Genova, M. Arese, M.C. Barone, A. Giuffrè, C. Bianchi, M. Brunori, G. Lenaz, P. Sarti, Control of respiration by nitric oxide in Keilin-Hartree particles, mitochondria and SH-SY5Y neuroblastoma cells, *Cellular and Molecular Life Sciences* 60 (8) (2003) 1752–1759.
- [73] P.R. Castello, P.S. David, T. McClure, Z. Crook, R.O. Poyton, Mitochondrial cytochrome oxidase produces nitric oxide under hypoxic conditions: implications for oxygen sensing and hypoxic signaling in eukaryotes, *Cell Metabolism* 3 (4) (2006) 277–287.
- [74] M.W. Cleeter, J.M. Cooper, V.M. Darley-Usmar, S. Moncada, A.H. Schapira, Reversible inhibition of cytochrome c oxidase, the terminal enzyme of the mitochondrial respiratory chain, by nitric oxide. Implications for neurodegenerative diseases, *FEBS Letters* 345 (1) (1994) 50–54.
- [75] S. Shiva, M.T. Gladwin, Nitrite mediates cytoprotection after ischemia/reperfusion by modulating mitochondrial function, *Basic Research in Cardiology* 104 (2) (2009) 113–119.
- [76] M.T. Gladwin, D.B. Kim-Shapiro, The functional nitrite reductase activity of the heme-globins, *Blood* 112 (7) (2008) 2636–2647.
- [77] M.P. Doyle, D.M. LePoire, R.A. Pickering, Oxidation of hemoglobin and myoglobin by alkyl nitrites inhibition by oxygen, *Journal of Biological Chemistry* 256 (23) (1981) 12399–12404.
- [78] J. Brooks, The action of nitrite on haemoglobin in the absence of oxygen, *Proceedings of the Royal Society of Medicine* 137 (1937) 368–382.
- [79] Z. Huang, S. Shiva, D.B. Kim-Shapiro, R.P. Patel, L.A. Ringwood, C.E. Irby, K. T. Huang, C. Ho, N. Hogg, A.N. Schechter, M.T. Gladwin, Enzymatic function of hemoglobin as a nitrite reductase that produces NO under allosteric control, *Journal of Clinical Investigation* 115 (8) (2005) 2099–2107.
- [80] S. Basu, R. Grubina, J. Huang, J. Conradie, Z. Huang, A. Jeffers, A. Jiang, X. He, I. Azarov, R. Seibert, A. Mehta, R. Patel, S.B. King, N. Hogg, A. Ghosh, M. T. Gladwin, D.B. Kim-Shapiro, Catalytic generation of N₂O₃ by the concerted nitrite reductase and anhydrase activity of hemoglobin, *Nature Chemical Biology* 3 (12) (2007) 785–794.
- [81] S. Shiva, Z. Huang, R. Grubina, J. Sun, L.A. Ringwood, P.H. MacArthur, X. Xu, E. Murphy, V.M. Darley-Usmar, M.T. Gladwin, Deoxyhemoglobin is a nitrite reductase that generates nitric oxide and regulates mitochondrial respiration, *Circulation Research* 100 (5) (2007) 654–661.
- [82] F.J. Larsen, T.A. Schiffer, S. Borniquel, K. Sahlin, B. Eklblom, J.O. Lundberg, E. Weitzberg, Dietary inorganic nitrate improves mitochondrial efficiency in humans, *Cell Metabolism* 13 (2) (2011) 149–159.
- [83] T. Rassaf, U. Flögel, C. Drexhage, U. Hendgen-Cotta, M. Kelm, J. Schrader, Nitrite reductase function of deoxyhemoglobin: oxygen sensor and regulator of cardiac energetics and function, *Circulation Research* 100 (12) (2007) 1749–1754.
- [84] T.M. Millar, C.R. Stevens, N. Benjamin, R. Eisenthal, R. Harrison, D.R. Blake, Xanthine oxidoreductase catalyses the reduction of nitrates and nitrite to nitric oxide under hypoxic conditions, *FEBS Letters* 427 (2) (1998) 225–228.
- [85] J.E. Baker, J. Su, X. Fu, A. Hsu, G.J. Gross, J.S. Tweddell, N. Hogg, Nitrite confers protection against myocardial infarction: role of xanthine oxidoreductase, NADPH oxidase and K(ATP) channels, *Journal of Molecular and Cellular Cardiology* 43 (4) (2007) 437–444.
- [86] T.M. Millar, Peroxynitrite formation from the simultaneous reduction of nitrite and oxygen by xanthine oxidase, *FEBS Letters* 562 (1–3) (2004) 129–133.
- [87] C. Szabo, H. Ischiropoulos, R. Radi, Peroxynitrite: biochemistry, pathophysiology and development of therapeutics, *Nature Reviews Drug Discovery* 6 (8) (2007) 662–680.
- [88] M.D. Gershon, The enteric nervous system: a second brain, *Hospital Practice (Minneapolis)* 34 (7) (1999) 31–32 5–8, 41–2 passim.
- [89] A.M. San Gabriel, T. Maekawa, H. Uneyama, S. Yoshie, K. Torii, mGluR1 in the fundic glands of rat stomach, *FEBS Letters* 581 (6) (2007) 1119–1123.
- [90] E.A. Mayer, The neurobiology of stress and gastrointestinal disease, *Gut* 47 (6) (2000) 861–869.
- [91] A. Kitamura, T. Tsurugizawa, A. Uematsu, K. Torii, H. Uneyama, New therapeutic strategy for amino acid medicine: effects of dietary glutamate on gut and brain function, *Journal of Pharmacological Sciences* 118 (2) (2012) 138–144.
- [92] T.D. Presley, A.R. Morgan, E. Bechtold, W. Clodfelter, R.W. Dove, J.M. Jennings, R.A. Kraft, S.B. King, P.J. Laurienti, W.J. Rejeski, J.H. Burdette, D.B. Kim-Shapiro, G.D. Miller, Acute effect of a high nitrate diet on brain perfusion in older adults, *Nitric Oxide* 24 (1) (2011) 34–42.
- [93] J. Garthwaite, C.L. Boulton, Nitric oxide signaling in the central nervous system, *Annual Review of Physiology* 57 (1995) 683–706.
- [94] D.S. Bredt, C.E. Glatt, P.M. Hwang, M. Fotuhi, T.M. Dawson, S.H. Snyder, Nitric oxide synthase protein and mRNA are discretely localized in neuronal populations of the mammalian CNS together with NADPH diaphorase, *Neuron* 7 (4) (1991) 615–624.
- [95] S.J. East, J. Garthwaite, NMDA receptor activation in rat hippocampus induces cyclic GMP formation through the L-arginine-nitric oxide pathway, *Neuroscience Letters* 123 (1) (1991) 17–19.
- [96] J. Garthwaite, Glutamate, nitric oxide and cell-cell signaling in the nervous system, *Trends in Neuroscience* 14 (2) (1991) 60–67.
- [97] E. Southam, J. Garthwaite, Nitric oxide-cyclic GMP pathway in brain slices, *Methods in Enzymology* 269 (1996) 129–133.
- [98] J. Garthwaite, Concepts of neural nitric oxide-mediated transmission, *European Journal of Neuroscience* 27 (11) (2008) 2783–2802.
- [99] F.X. Guix, I. Urbesalago, M. Coma, F.J. Muñoz, The physiology and pathophysiology of nitric oxide in the brain, *Progress in Neurobiology* 76 (2) (2005) 126–152.
- [100] J.R. Steinert, T. Chernova, I.D. Forsythe, Nitric oxide signaling in brain function, dysfunction, and dementia, *Neuroscientist* 16 (4) (2010) 435–452.
- [101] R.M. Santos, C.F. Lourenco, G.A. Gerhardt, E. Cadenas, J. Laranjinha, R. M. Barbosa, Evidence for a pathway that facilitates nitric oxide diffusion in the brain, *Neurochemistry International* 59 (1) (2011) 90–96.
- [102] C.E. Cooper, Nitric oxide and iron proteins, *Biochimica Biophysica Acta* 1411 (2–3) (1999) 290–309.
- [103] A. Ledo, R.M. Barbosa, J. Frade, J. Laranjinha, Nitric oxide monitoring in hippocampal brain slices using electrochemical methods, *Methods in Enzymology* 359 (2002) 111–125.
- [104] N.R. Ferreira, A. Ledo, J.G. Frade, G.A. Gerhardt, J. Laranjinha, R.M. Barbosa, Electrochemical measurement of endogenously produced nitric oxide in brain slices using Nafion/o-phenylenediamine modified carbon fiber microelectrodes, *Analytica Chimica Acta* 535 (1–2) (2005) 1–7.
- [105] R.M. Barbosa, C.F. Lourenco, R.M. Santos, F. Pomerleau, P. Huettl, G. A. Gerhardt, J. Laranjinha, *In vivo* real-time measurement of nitric oxide in anesthetized rat brain, *Methods in Enzymology* 441 (2008) 351–367.
- [106] R.M. Santos, C.F. Lourenco, A.P. Piedade, R. Andrews, F. Pomerleau, P. Huettl, G.A. Gerhardt, J. Laranjinha, R.M. Barbosa, A comparative study of carbon fiber-based microelectrodes for the measurement of nitric oxide in brain tissue, *Sensors and Bioelectronics* 24 (4) (2008) 704–709.
- [107] J.L. Peters, L.H. Miner, A.C. Michael, S.R. Sesack, Ultrastructure at carbon fiber microelectrode implantation sites after acute voltammetric measurements in the striatum of anesthetized rats, *Journal of Neuroscience Methods* 137 (1) (2004) 9–23.
- [108] D. Attwell, A.M. Buchan, S. Charpak, M. Lauritzen, B.A. Macvicar, E. A. Newman, Glial and neuronal control of brain blood flow, *Nature* 468 (7321) (2010) 232–243.
- [109] J. Millar, The nitric oxide/ascorbate cycle: how neurones may control their own oxygen supply, *Molecular Hypotheses* 45 (1) (1995) 21–26.
- [110] A.B. Milsom, B.O. Fernandez, M.F. Garcia-Saura, J. Rodriguez, M. Feelisch, Contributions of nitric oxide synthases, dietary nitrite/nitrate, and other sources to the formation of NO signaling products, *Antioxidants and Redox Signaling* 17 (3) (2012) 422–432.
- [111] M.T. Gladwin, N.J. Raat, S. Shiva, C. Dezfulian, N. Hogg, D.B. Kim-Shapiro, R. P. Patel, Nitrite as a vascular endocrine nitric oxide reservoir that contributes to hypoxic signaling, cytoprotection, and vasodilation, *American Journal of Physiology: Heart and Circulatory Physiology* 291 (5) (2006) H2026–H2035.

- [112] C.E. Sparacino-Watkins, Y.C. Lai, M.T. Gladwin, Nitrate-nitrite-nitric oxide pathway in pulmonary arterial hypertension therapeutics, *Circulation* 125 (23) (2012) 2824–2826.
- [113] M. Totzeck, U.B. Hendgen-Cotta, P. Luedike, M. Berenbrink, J.P. Klare, H. J. Steinhoff, D. Semmler, S. Shiva, D. Williams, A. Kipar, M.T. Gladwin, J. Schrader, M. Kelm, A.R. Cossins, T. Rassaf, Nitrite regulates hypoxic vasodilation via myoglobin-dependent nitric oxide generation, *Circulation* 126 (3) (2012) 325–334.
- [114] B.S. Zuckerbraun, S. Shiva, E. Ifedigbo, M.A. Mathier, K.P. Mollen, J. Rao, P. M. Bauer, J.J. Choi, E. Curtis, A.M. Choi, M.T. Gladwin, Nitrite potently inhibits hypoxic and inflammatory pulmonary arterial hypertension and smooth muscle proliferation via xanthine oxidoreductase-dependent nitric oxide generation, *Circulation* 121 (1) (2010) 98–109.
- [115] C.L. Pedersen, S. Faggiano, S. Helbo, H. Gesser, A. Fago, Roles of nitric oxide, nitrite and myoglobin on myocardial efficiency in trout (*Oncorhynchus mykiss*) and goldfish (*Carassius auratus*): implications for hypoxia tolerance, *Journal of Experimental Biology* 213 (Pt 16) (2010) 2755–2762.
- [116] I. Mikula, S. Durocher, P. Martasek, B. Mutus, A. Slama-Schwok, Isoform-specific differences in the nitrite reductase activity of nitric oxide synthases under hypoxia, *Biochemical Journal* 418 (3) (2009) 673–682.
- [117] A.J. Webb, A.B. Milsom, K.S. Rathod, W.L. Chu, S. Qureshi, M.J. Lovell, F. M. Lecomte, D. Perrett, C. Raimondo, E. Khoshbin, Z. Ahmed, R. Uppal, N. Benjamin, A.J. Hobbs, A. Ahluwalia, Mechanisms underlying erythrocyte and endothelial nitrite reduction to nitric oxide in hypoxia: role for xanthine oxidoreductase and endothelial nitric oxide synthase, *Circulation Research* 103 (9) (2008) 957–964.
- [118] M. Feelisch, B.O. Fernandez, N.S. Bryan, M.F. Garcia-Saura, S. Bauer, D. R. Whitlock, P.C. Ford, D.R. Janero, J. Rodriguez, H. Ashrafian, Tissue processing of nitrite in hypoxia: an intricate interplay of nitric oxide-generating and scavenging systems, *Journal of Biological Chemistry* 283 (49) (2008) 33927–33934.
- [119] B. Pisknova, A. Kocharyan, A.N. Schechter, A.C. Silva, The role of nitrite in neurovascular coupling, *Brain Research* 1407 (2011) 62–68.
- [120] J.M. Rifkind, E. Nagababu, E. Barbiro-Michaely, S. Ramasamy, R.M. Pluta, A. Mayevsky, Nitrite infusion increases cerebral blood flow and decreases mean arterial blood pressure in rats: a role for red cell NO, *Nitric Oxide* 16 (4) (2007) 448–456.
- [121] M.E. Rice, Ascorbate regulation and its neuroprotective role in the brain, *Trends in Neuroscience* 23 (5) (2000) 209–216.
- [122] R.A. Grunewald, Ascorbic acid in the brain, *Brain Research Reviews* 18 (1) (1993) 123–133.
- [123] A. Corti, A.F. Casini, A. Pompella, Cellular pathways for transport and efflux of ascorbate and dehydroascorbate, *Archives of Biochemistry and Biophysics* 500 (2) (2010) 107–115.
- [124] G.V. Rebec, R.C. Pierce, A vitamin as neuromodulator: ascorbate release into the extracellular fluid of the brain regulates dopaminergic and glutamatergic transmission, *Progress in Neurobiology* 43 (6) (1994) 537–565.
- [125] C.C. Portugal, V.S. Miya, C. Calaza Kda, R.A. Santos, R. Paes-de-Carvalho, Glutamate receptors modulate sodium-dependent and calcium-independent vitamin C bidirectional transport in cultured avian retinal cells, *Journal of Neurochemistry* 108 (2) (2009) 507–520.
- [126] N.V. Kulagina, L. Shankar, A.C. Michael, Monitoring glutamate and ascorbate in the extracellular space of brain tissue with electrochemical microsensors, *Analytical Chemistry* 71 (22) (1999) 5093–5100.
- [127] A. van der Toorn, E. Sykova, R.M. Dijkhuizen, I. Vorisek, L. Vargova, E. Skobisova, M. van Lookeren Campagne, T. Reese, K. Nicolay, Dynamic changes in water ADC, energy metabolism, extracellular space volume, and tortuosity in neonatal rat brain during global ischemia, *Magnetic Resonance in Medical Sciences* 36 (1) (1996) 52–60.
- [128] J. Watts, P.S. Whitton, B. Pearce, Unexpected effects of nitric oxide synthase inhibitors on extracellular nitrite levels in the hippocampus in vivo, *Pharmacology* 74 (3) (2005) 163–168.
- [129] A.H. Cross, P.T. Manning, R.M. Keeling, R.E. Schmidt, T.P. Misko, Peroxynitrite formation within the central nervous system in active multiple sclerosis, *Journal of Neuroimmunology* 88 (1–2) (1998) 45–56.
- [130] W.S. Jobgen, S.C. Jobgen, H. Li, C.J. Meininger, G. Wu, Analysis of nitrite and nitrate in biological samples using high-performance liquid chromatography, *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences* 851 (1–2) (2007) 71–82.