

A BIJECTION BETWEEN NONCROSSING AND NONNESTING PARTITIONS OF TYPES A, B AND C

RICARDO MAMEDE

ABSTRACT: The total number of noncrossing partitions of type Ψ is the n th Catalan number $\frac{1}{n+1}\binom{2n}{n}$ when $\Psi = A_{n-1}$, and the coefficient binomial $\binom{2n}{n}$ when $\Psi = B_n$ or C_n , and these numbers coincide with the correspondent number of nonnesting partitions. For type A, there are several bijective proofs of this equality; in particular, the intuitive map, which locally converts each crossing to a nesting, is one of them. In this paper we present a bijection between nonnesting and noncrossing partitions of types A, B and C that generalizes the type A bijection that locally converts each crossing to a nesting.

KEYWORDS: Root systems, noncrossing partitions, nonnesting partitions, bijection.

AMS SUBJECT CLASSIFICATION (2000): 05A10; 05A15.

1. Introduction

The poset of noncrossing partitions can be defined in a uniform way for any finite Coxeter group W . More precisely, for $u, w \in W$, let $u \leq w$ if there is a shortest factorization of w as a product of reflections in W having as prefix such a shortest factorization for u . This partial order turns W into a graded poset $Abs(W)$ having the identity 1 as its unique minimal element, where the rank of w is the length of the shortest factorization of w into reflections. Let c be a Coxeter element of W . Since all Coxeter elements in W are conjugate to each other, the interval $[1, c]$ in $Abs(W)$ is independent, up to isomorphism, of the choice of c . We denote this interval by $NC(W)$ or by $NC(\Psi)$, where Ψ is the Cartan-Killing type of W , and call it the *poset of noncrossing partitions* of W . It is a self-dual, graded lattice which reduces to the classical lattice of noncrossing partitions of the set $[n] = \{1, 2, \dots, n\}$ defined by Kreweras in [9] when W is the symmetric group \mathfrak{S}_n (the Coxeter group of type A_{n-1}), and to its types B and C analogues defined by Reiner [11] when W is the hyperoctahedral group. The elements in $NC(W)$ are

Received March 11, 2009.

This work was supported by CMUC - Centro de Matemática da Universidade de Coimbra.

counted by the generalized Catalan numbers,

$$Cat(W) = \prod_{i=1}^k \frac{d_i + h}{d_i},$$

where k is the number of simple reflections in W , h is the Coxeter number and d_1, \dots, d_k are the degrees of the fundamental invariants of W (see [1, 6, 7, 11] for details on the theory of Coxeter groups and noncrossing partitions). When W is the symmetric group \mathfrak{S}_n , the number $Cat(\mathfrak{S}_n)$ is just the usual n th Catalan number $\frac{1}{n+1} \binom{2n}{n}$, and in types B_n and C_n this number is the binomial coefficient $\binom{2n}{n}$.

Nonnesting partitions were defined by Postnikov (see [11, Remark 2]) in a uniform way for all irreducible root systems associated with Weyl groups. If Φ is such a system, Φ^+ is a choice of positive roots, and Δ is the simple system in Φ^+ , define the *root order* on Φ^+ by $\alpha \leq \beta$ if $\alpha, \beta \in \Phi^+$ and $\beta - \alpha$ is in the positive integer span of the simple roots in Δ . Equipped with this partial order, (Φ^+, \leq) is the *root poset* of the associated Weyl group W . A *nonnesting partition* on Φ is just an antichain in root poset (Φ^+, \leq) , that is, a set of pairwise incomparable elements. Denote by $NN(W)$ or by $NN(\Psi)$, where Ψ is the Cartan-Killing type of W , the set of all nonnesting partitions of W . Postnikov showed that the nonnesting partitions in $NN(W)$ are also counted by the generalized Catalan number $Cat(W)$.

In the case of the root systems of type A , different bijective proofs of the equality between the cardinals $|NN(A_{n-1})| = |NC(A_{n-1})|$ are known (see [1, 2, 3, 8, 12]). Recently, Christian Stump [12] described a bijection between nonnesting and noncrossing partitions for type B , and simultaneously with our work, Alex Fink and Benjamin Giraldo [5, 10] have presented a different bijection for each classical group. Our contribution in this paper is to present a uniform proof that $|NN(\Psi)| = |NC(\Psi)|$, for $\Psi = A_{n-1}, B_n$ or C_n that generalizes the bijection presented by Armstrong in [1].

2. Noncrossing and nonnesting set partitions

A *set partition* of $[n]$ is a collection of nonempty disjoint subsets of $[n]$, called *blocks*, whose union is $[n]$. The type of a set partition π of $[n]$ is the integer partition formed by the cardinals of the blocks of π . A set partition of $[n]$ type $(2, \dots, 2, 1, \dots, 1)$ is called a partial matching, and a set partition of $[2n]$ of type $(2, 2, \dots, 2)$ is said to be a (perfect) matching of $[2n]$.

A set partition can be graphically represented by placing the integers $1, 2, \dots, n$ along a line and drawing arcs above the line between i and j whenever i and j lie in the same block and no other element between them does so. A *singleton* of a set partition is a block which has only one element, so it corresponds to an isolated vertex in the graphical representation. For instance, the graphical representation of the set partition $\pi = \{\{1, 3, 4\}, \{2, 6\}, \{5\}\}$ of type $(3, 2, 1)$ is displayed below:



Given a set partition π , let

$$\begin{aligned} op(\pi) &= \{\text{least block elements of } \pi\} \\ cl(\pi) &= \{\text{greatest block elements of } \pi\} \\ tr(\pi) &= [n] \setminus (op(\pi) \cup cl(\pi)) \end{aligned}$$

The elements of $op(\pi)$, $cl(\pi)$, and $tr(\pi)$ are called openers, closers and transients, respectively. Graphically, the openers corresponds to singletons and to vertices from which one arc begin and no arc ends, the closers corresponds to singletons and to vertices to which one arc ends, and no arc begins, and the transients are the vertices from which one arc ends an another one begins. In the example above, $op(\pi) = \{1, 2, 5\}$, $cl(\pi) = \{4, 5, 6\}$ and $tr(\pi) = \{3\}$. The triples $\mathfrak{T}(\pi) = (op(\pi), tr(\pi), cl(\pi))$ encodes useful information about the partition π . For instance, the number of blocks is $|op(\pi)| = |cl(\pi)|$, the number of singletons is $|op(\pi) \cap cl(\pi)|$, π is a partial matching if and only if $tr(\pi) = \emptyset$, and π is a (perfect) matching if and only if $tr(\pi) = \emptyset$ and $op(\pi) \cap cl(\pi) = \emptyset$.


A noncrossing partition of the set $[n]$ is a set partition of $[n]$ such that there are no $a < b < c < d$, with a, c belonging to some block of the partition and b, d belonging to some other block. The set of noncrossing partitions of $[n]$, denoted by $\text{NC}(n)$, is a lattice for the refinement order. A nonnesting partition of the set $[n]$ is a partition of $[n]$ such that if $a < b < c < d$ and a, d are consecutive elements of a block, then b and c are not both contained in some other block. The set of nonnesting partitions of $[n]$ will be denoted by $\text{NN}(n)$. Graphically, the noncrossing condition means that no two of the arcs cross, while the nonnesting condition means that no two arcs are nested one within the other.

The partition $\pi = \{\{1, 3, 4\}, \{2, 6\}, \{5\}\}$ represented above is neither noncrossing nor nonnesting since two of the arcs cross, and two of the arcs are


nested one within the other. The partitions

$$\{\{2, 3\}, \{1, 4, 5\}\} \quad \text{and} \quad \{\{1, 3\}, \{2, 4, 5\}\},$$

represented below, are examples of noncrossing and nonnesting partitions of $[5]$:



As pointed out in [1], the intuitive map that locally converts each crossing to a nesting



defines a bijection between noncrossing and nonnesting set partitions that preserves the number of blocks. We will refer to this bijection as the L-map.

A B_n set partition π is a set partition of $[\pm n] := \{\pm 1, \pm 2, \dots, \pm n\}$ which has at most one block (called the zero block) fixed by negation and is such that for any block B of π , the set $-B$, obtained by negating the elements of B , is also a block of π . The type of a B_n set partition π is the integer partition whose parts are the cardinalities of the blocks of π , including one part for each pair of nonzero blocks $B, -B$. Given a B_n set partition π , let the set of openers $op(\pi)$ be formed by the least element of all blocks of π having only positive integers; let the set of closers $cl(\pi)$ be formed by the greatest element of all blocks of π having only positive integers and by the absolute values of the least and greatest elements of all blocks having positive and negative integers; and finally let the set of transients $tr(\pi)$ be formed by all elements of $[n]$ which are not in $op(\pi) \cup cl(\pi)$.


Identifying the sets $[\pm n]$ and $[2n]$ through the map $i \mapsto i$ for $i \in [n]$ and $i \mapsto n-i$ for $i \in \{-1, -2, \dots, -n\}$, we may represent B_n set partitions graphically using the conventions made for its type A analogs, placing the integers $-1, -2, \dots, -n, 1, 2, \dots, n$ along a line instead of the usual $1, 2, \dots, 2n$.

Example 2.1. The B_5 set partition

$$\pi = \{\{-1, 1\}, \{2, 3, 5\}, \{-2, -3, -5\}, \{4\}, \{-4\}\},$$

represented below, has type $(3, 2, 1)$, set of openers $op(\pi) = \{2, 4\}$, closers $cl(\pi) = \{1, 4, 5\}$ and transients $tr(\pi) = \{3\}$. The openers, closers and transients can be visualized as in type A by looking only at the arcs on the

positive half of the representation of π .



3. Noncrossing and nonnesting partitions of types A , B and C

We will now review the usual combinatorial realizations of the Coxeter groups of types A , B and C , referring to [7] for any undefined terminology. The Coxeter group W of type A_{n-1} is realized combinatorially as the symmetric group \mathfrak{S}_n . The permutations in \mathfrak{S}_n will be written in cycle notation. The simple generators of \mathfrak{S}_n are the transpositions of adjacent integers $(i \ i+1)$, for $i = 1, \dots, n-1$, and the reflections are the transpositions $(i \ j)$ for $1 \leq i < j \leq n$. To any permutation $\pi \in \mathfrak{S}_n$ we associate the partition of the set $[n]$ given by its cycle structure. This defines an isomorphism between the posets $\text{NC}(\mathfrak{S}_n)$ of noncrossing partitions of \mathfrak{S}_n , defined in the introduction, and $\text{NC}(n)$, with respect to the Coxeter element $c = (12 \cdots n)$ [4, Theorem 1].

Denoting by e_1, \dots, e_n the standard basis of \mathbb{R}^n , the root system of type A_{n-1} consists of the set of vectors

$$\Phi = \{e_i - e_j : i \neq j, 1 \leq i, j \leq n\},$$

each root $e_i - e_j$ defining a reflection that acts on \mathbb{R}^n as the transposition $(i \ j)$. We shall identify the root $e_i - e_j$ with the corresponding transposition $(i \ j)$. Take


$$\Phi^+ = \{e_i - e_j \in \Phi : i > j\}$$

for the set of positive roots and, defining $r_i := e_{i+1} - e_i$, $i = 1, \dots, n-1$, we obtain the simple system $\Delta = \{r_1, \dots, r_{n-1}\}$ for \mathfrak{S}_n . Note that

$$e_i - e_j = \sum_{k=j}^{i-1} r_k, \quad \text{if } i > j.$$

The correspondence between the antichains in the root poset (Φ^+, \leq) and the set of nonnesting partitions of $[n]$ is given by the bijection which sends the positive root $e_i - e_j$ to the set partition of $[n]$ having i and j in the same

block. For instance, consider the root poset (Φ^+, \leq) of type A_4 :



The antichain $r_1 + r_2 = e_3 - e_1$ corresponds to the transposition (13) in the symmetric group \mathfrak{S}_5 , and thus to the nonnesting set partition $\{\{1, 3\}, \{2\}, \{4\}, \{5\}\}$, while the antichain $\{r_1 + r_2, r_2 + r_3, r_4\}$ corresponds to the product of transpositions (13)(24)(45) = (13)(245) in \mathfrak{S}_5 , and thus to the nonnesting set partition $\{\{1, 3\}, \{2, 4, 5\}\}$.

Given a positive root $\alpha = r_i + r_{i+1} + \cdots + r_j \in \Phi^+$, define the *support* of α as the set $\text{supp}(\alpha) = \{r_i, r_{i+1}, \dots, r_j\}$. The integers i and j will be called, respectively, the *first* and *last indices* of α , and the roots r_i and r_j the first and last elements of α , respectively. We have the following lemma.

Lemma 3.1. *Let α_1, α_2 be two roots in Φ^+ with first and last indices i_1, j_1 and i_2, j_2 , respectively. Then, α_1, α_2 form an antichain if and only if $i_1 < i_2$ and $j_1 < j_2$.*

Consider now the Coxeter group W of type B_n , with its usual combinatorial realization as the hyperoctahedral group of signed permutations of

$$[\pm n] := \{\pm 1, \pm 2, \dots, \pm n\}.$$

These are permutations of $[\pm n]$ which commute with the involution $i \mapsto -i$. We will write the elements of W in cycle notation, using commas between elements. The simple generators of W are the transposition $(-1, 1)$ and the pairs $(-i-1, -i)(i, i+1)$ for $i = 1, \dots, n-1$. The reflections in W are the transpositions $(-i, i)$ for $i = 1, \dots, n$, and the pairs of transpositions $(i, j)(-j, -i)$ for $i \neq j$.

Identifying the sets $[\pm n]$ and $[2n]$ through the map $i \mapsto i$ for $i \in [n]$ and $i \mapsto n-i$ for $i \in \{-1, -2, \dots, -n\}$, allows us to identify the hyperoctahedral group W with the subgroup U of \mathfrak{S}_{2n} which commutes with the permutation $(1, n+1)(2, n+2) \cdots (n, 2n)$. For example, the signed permutations $(1, 3)$ and $(2, -3)(-2, 3)$ in the hyperoctahedral group of type B_3 correspond to

the permutations $(1\ 3)$ and $(2\ 6)(5\ 3)$ in the symmetric group \mathfrak{S}_6 . The set $\text{NC}(U)$ is a sublattice of $\text{NC}(\mathfrak{S}_{2n})$, isomorphic to $\text{NC}(W)$ (see [1]). It follows that the map sending a signed permutation of $[\pm n]$ to the B_n set partition of $[\pm n]$ given by its cycle structure defines an isomorphism between $\text{NC}(W)$ and the B_n set partitions in which no two arcs cross.

The type B_n root system consists on the set of $2n^2$ vectors

$$\Phi = \{\pm e_i : 1 \leq i \leq n\} \cup \{\pm e_i \pm e_j : i \neq j, 1 \leq i, j \leq n\},$$

and we take

$$\Phi^+ = \{e_i : 1 \leq i \leq n\} \cup \{e_i \pm e_j : 1 \leq j < i \leq n\}$$

as a choice of positive roots. Changing the notation slightly from the one used for \mathfrak{S}_n , let $r_1 := e_1$ and $r_i := e_i - e_{i-1}$, for $i = 2, \dots, n$. The set

$$\Delta := \{r_1, r_2, \dots, r_n\}$$

is a simple system for W , and easy computations show that


$$\begin{aligned} e_i &= \sum_{k=1}^i r_k, \\ e_i - e_j &= \sum_{k=j+1}^i r_k, & \text{if } i > j \\ e_i + e_j &= 2 \sum_{k=1}^j r_k + \sum_{k=j+1}^i r_k, & \text{if } i > j. \end{aligned}$$

Each root $e_i, e_i - e_j$ and $e_i + e_j$ defines a reflection that acts on \mathbb{R}^n as the permutation $(i, -i), (i, j)(-i, -j)$ and $(i, -j)(-i, j)$, respectively, and we will identify the roots with the corresponding permutations.


A graphical representation of a nonnesting partition $\pi \in \text{NN}(B_n)$ can be drawn by placing the integers $-n, \dots, -2, -1, 0, 1, 2, \dots, n$, in this order, along a line and arcs between them as follows: for $i, j \in [n]$, we include an arc between i and j , and between $-i$ and $-j$, if π contains the root $e_i - e_j$; an arc between i and $-j$, and between $-i$ and j if π contains the root $e_i + e_j$; and arcs between i and 0 and -1 and 0 if π contains the root e_i . The presence of 0 in the ground set for nonnesting partitions is necessary to correctly represent (when present) the arc between a positive number i and its negative (see [2]). The chains of successive arcs in the diagram become the blocks of a B_n set

partition, after dropping 0, which is the partition we associate to π . This map defines a bijection between nonnesting partitions of W and B_n set partitions whose diagrams, in the above sense, contain no two arcs nested one within the other. We call this diagram the nonnesting graphical representation of π , to distinguish it from the graphical representation of the B_n set partition associated to π .

Example 3.1. Consider the root poset of type B_3 is displayed below:



In this root poset, the antichain $\{2r_1 + r_2, r_1 + r_2 + r_3\}$ corresponds to the B_3 set partition $\pi = \{\{3, -3\}, \{1, -2\}, \{-1, 2\}\}$, and thus to signed permutation $(3, -3)(1, -2)(-1, 2)$. The nonnesting graphical representation of π is given below:



A factor $2r_i$ appearing in a positive root α will be called a *double root* of α . The *support* of a positive root $\alpha \in \Phi^+$ is the set of simple and double roots in α . Define also the set $\overline{\text{supp}}(\alpha)$ as the set formed by the simple roots appearing in alpha as simple or double roots. For instance, for $\alpha = 2r_1 + \dots + 2r_j + r_{j+1} + \dots + r_i$ and $\beta = r_\ell + \dots + r_k$ we have $\text{supp}(\alpha) = \{2r_1, \dots, 2r_j, r_{j+1}, \dots, r_i\}$, $\overline{\text{supp}}(\alpha) = \{r_1, \dots, r_j, r_{j+1}, \dots, r_i\}$, and $\text{supp}(\beta) = \overline{\text{supp}}(\beta) = \{r_\ell, \dots, r_k\}$. The first and last indices of α and β are, respectively, $1, i$ and ℓ, k . The integer j will be called the last double index in α . Define also the set $D_\alpha := \{r_2, \dots, r_j\}$ as the set of simple roots appearing in α as double roots, other than r_1 . We have the following lemma.

Lemma 3.2. *Let α and β be two roots in Φ^+ with first indices i, i' and last indices j, j' , respectively. If neither α nor β have double roots, then $\{\alpha, \beta\}$ is an antichain if and only if $i < i'$ and $j < j'$. If α has double roots, then*

$\{\alpha, \beta\}$ is an antichain if and only if $j < j'$ and the number of double roots in α is greater than the number of double roots in β .

The root system of type C_n is obtained from the root system of type B_n by replacing the roots e_i by $2e_i$,

$$\Phi = \{\pm 2e_i : 1 \leq i \leq n\} \cup \{\pm e_i \pm e_j : i \neq j, 1 \leq i, j \leq n\}.$$

For $n \geq 3$, the two root systems are not congruent, but their corresponding Weyl groups, generated by reflections orthogonal to the roots, are clearly the same. Thus, the relations between roots of type C_n and signed permutations of $[\pm n]$ is the same as in type B_n . We let

$$\Phi^+ = \{2e_i : 1 \leq i \leq n\} \cup \{e_i \pm e_j : 1 \leq j < i \leq n\}$$

as a choice of positive roots, and put $r_1 := 2e_1$ and $r_i := e_i - e_{i-1}$, for $i = 2, \dots, n$. The set


$$\Delta := \{r_1, r_2, \dots, r_n\}$$

is a simple system for W , and easy computations show that


$$\begin{aligned} 2e_i &= r_1 + \sum_{k=2}^i 2r_k, \\ e_i - e_j &= \sum_{k=j+1}^i r_k \quad (i > j) \\ e_i + e_j &= r_1 + \sum_{k=2}^j 2r_k + \sum_{k=j+1}^i r_k \quad (i > j), \\ e_i + e_1 &= \sum_{k=1}^i r_k. \end{aligned}$$

The notions of C_n set partition and noncrossing partitions of type C_n coincide with its type B_n analogs, as well as the notions of opener, closer and transient. Also, the nonnesting diagram associated to a nonnesting partition $\pi \in \text{NN}(C_n)$ is determined as in type B_n , except that i and $-i$ are connected by an arc if π contains $2e_i$ and that 0 does not appear in the diagram. Again, this map defines a bijection between nonnesting partitions of type C_n and C_n set partitions whose diagrams, in the above sense, contain no two arcs nested one within the other.

Example 3.2. Consider the root poset of type C_3 :



In this root poset, the antichain $\{r_1 + 2r_2, r_1 + r_2 + r_3\}$ corresponds to the C_3 set partition $\pi = \{\{2, -2\}, \{1, -3\}, \{-1, 3\}\}$, and thus to the signed permutation $(2, -2)(1, -3)(-1, 3)$. Its nonnesting diagram is represented below.



Note that this set partition π is not a nonnesting partition of type B_3 , since, considered in type B_3 , there would be an arc linking 0 to 2 which would be nested by the arc linking -1 to 3.

Import to type C the notions of double root and support of a positive root. We have the following lemma.

Lemma 3.3. *Let α and β be two roots in Φ^+ with first indices i, i' and last indices j, j' , respectively. If α has double roots, then $\{\alpha, \beta\}$ is an antichain if and only if the number of double roots in α is greater than the number of double roots in β and $j < j'$. If neither α nor β have double roots, then $\{\alpha, \beta\}$ is an antichain if and only if $i < i'$ and $j < j'$.*

Although the root systems of type B_n and C_n are not congruent, there is a simple connection between the antichains in $\text{NN}(B_n)$ and those in $\text{NN}(C_n)$.

Proposition 3.4. *There is a bijection τ between antichains in the root posets of types B_n and C_n which preserves the triples (op, cl, tr) .*

Proof: First, notice that if $\pi \in \text{NN}(B_n)$ does not have the positive root $r_1 + r_2 + \cdots + r_k$, for some $k \in [n]$, then the set $\tau(\pi)$, obtained by replacing each root $2r_1 + 2r_2 + \cdots + 2r_i + r_{i+1} + \cdots + r_j$, $i < j$, in π (if any) by the root $r_1 + 2r_2 + \cdots + 2r_i + r_{i+1} + \cdots + r_j$, is an antichain in $\text{NN}(C_n)$ which does not

have the positive root $r_1 + 2r_2 + \cdots + 2r_k$, and both π and $\tau(\pi)$ correspond to the same $B_n(C_n)$ set partition. In particular, $\mathfrak{T}(\pi) = \mathfrak{T}(\tau(\pi))$.

Similarly, if π possesses the root $r_1 + r_2 + \cdots + r_k$, but it does not have any root with double roots, then the set $\tau(\pi)$, obtained by replacing $r_1 + r_2 + \cdots + r_k$ by $r_1 + 2r_2 + \cdots + 2r_k$, is again an antichain in $\text{NN}(C_n)$ and both π and $\tau(\pi)$ correspond to the same $B_n(C_n)$ set partition. Again, we find that $\mathfrak{T}(\pi) = \mathfrak{T}(\tau(\pi))$.

Assume now that $\pi \in \text{NN}(B_n)$ has a positive root with double roots and also has the root $r_1 + r_2 + \cdots + r_k$, for some $k \in [n]$. Notice that in this case, by lemma 3.3, the $B_n(C_n)$ set partition associated with π does not corresponds to an antichain in $\text{NN}(C_n)$. Let $\pi' = \{\alpha_1, \alpha_2, \dots, \alpha_\ell\}$ be the antichain formed by all the roots in π having double roots and by the root $r_1 + r_2 + \cdots + r_k$. If

$$\begin{aligned}\alpha_1 &= 2r_1 + \cdots + 2r_a + r_{a+1} + \cdots + r_b, \\ \alpha_2 &= 2r_1 + \cdots + 2r_c + r_{c+1} + \cdots + r_d, \\ \alpha_3 &= 2r_1 + \cdots + 2r_e + r_{e+1} + \cdots + r_f, \\ &\vdots \\ \alpha_{\ell-1} &= 2r_1 + \cdots + 2r_g + r_{g+1} + \cdots + r_h, \\ \alpha_\ell &= r_1 + \cdots + r_k,\end{aligned}$$

with $g < \cdots < e < c < a < b < d < f < \cdots < h < k$, then define the following C_n roots:

$$\begin{aligned}\alpha'_1 &= r_1 + 2r_2 + \cdots + 2r_b, \\ \alpha'_2 &= r_1 + 2r_2 + \cdots + 2r_a + r_{a+1} + \cdots + r_d, \\ \alpha'_3 &= r_1 + 2r_2 + \cdots + 2r_c + r_{c+1} + \cdots + r_f, \\ &\vdots \\ \alpha'_\ell &= r_1 + 2r_2 + \cdots + 2r_g + r_{g+1} + \cdots + r_k.\end{aligned}$$

Let $\tau(\pi)$ be the set formed by replacing each root α_i , $i = 1, \dots, \ell$, in π by the C_n roots α'_i , $i = 1, \dots, \ell$. By lemma 3.3, $\tau(\pi)$ is an antichain in $\text{NN}(C_n)$ and it is straightforward to see that $\mathfrak{T}(\pi) = \mathfrak{T}(\tau(\pi))$. Also, notice that by lemma 3.2, the $B_n(C_n)$ set partition associated with $\tau(\pi)$ does not corresponds to an antichain in $\text{NN}(B_n)$.

This map τ is clearly injective, and since both $\text{NN}(B_n)$ and $\text{NN}(C_n)$ have the same cardinality, it establishes a bijection between antichains in the root posets of types B_n and C_n which preserves the triples (op, cl, tr) . ■


Example 3.3. Consider the antichain

$$\pi = \{2r_1 + 2r_2 + r_3, r_1 + r_2 + r_3 + r_4, r_3 + r_4 + r_5\}$$

is $\text{NN}(B_5)$, associated to the B_5 (C_5) set partition

$$\{\{4, -4\}, \{-3, 2, 5\}, \{-5, -2, 3\}, \{1\}, \{-1\}\}.$$

Its nonnesting diagram is represented below:




Note that this set partition does not corresponds to a nonnesting diagram of type C_5 , since in this case the arc linking -4 to 4 would nest the arc between -3 and 2 . Applying the construction given in the proposition above, we get the antichain

$$\tau(\pi) = \{r_1 + 2r_2 + 2r_3, r_1 + 2r_2 + r_3 + r_4, r_3 + r_4 + r_5\}$$

in $\text{NN}(C_5)$, corresponding to the B_5 (C_5) set partition

$$\{\{3, -3\}, \{-4, 2, 5\}, \{-5, -2, 4\}, \{1\}, \{-1\}\},$$

whose nonnesting diagram is represented below:



As before, the B_5 (C_5) set partition associated with $\tau(\pi)$ does not corresponds to a nonnesting diagram of type B_5 , since in this case the arc linking -3 to 0 would be nested by the arc linking -4 to 2 .

In what follows we will identify antichains in the root posets of types B_n and C_n by the bijection τ , *i.e.*, if π is an antichain in $\text{NN}(C_n)$, we will consider its image $\tau(\pi)$ in $\text{NN}(B_n)$.

4. Main result

Let Φ denote a root system of type A , B or C , and let Φ^+ and Δ be defined as above. In view of lemmas 3.1 and 3.2, we consider antichains $\{\alpha_1, \dots, \alpha_m\}$ in Φ^+ as ordered m -tuples numbered so that if i_ℓ is the last index of α_ℓ , then $i_1 < \dots < i_m$.

Definition 4.1. Given two positive roots α and β , with β having no double roots, and such that the intersection of their supports is nonempty, define their union $\alpha \cup \beta$ and their intersection $\alpha \cap \beta$ as the positive roots with supports

$$\text{supp}(\alpha \cup \beta) := \text{supp}(\alpha) \cup (\text{supp}(\beta) \setminus \overline{\text{supp}}(\alpha))$$

and

$$\text{supp}(\alpha \cap \beta) := \overline{\text{supp}}(\alpha) \cap \text{supp}(\beta),$$

respectively. If moreover α has double roots, define also their d-intersection $\alpha \cap^d \beta$ as the positive root with support $\text{supp}(\alpha \cap^d \beta) := D_\alpha \cap \text{supp}(\beta)$.

Example 4.1. The union and the intersections of the type B_3 positive roots $\alpha = 2r_1 + 2r_2 + r_3$ and $\beta = r_2 + r_3 + r_4$ are $\alpha \cup \beta = 2r_1 + 2r_2 + r_3 + r_4$, $\alpha \cap^d \beta = r_2$, and $\alpha \cap \beta = r_2 + r_3$.

An antichain $(\alpha_1, \dots, \alpha_m)$ is said to be *connected* if the intersection of the supports of any two adjacent roots α_i, α_{i+1} is non empty. The connected components

$$(\alpha_1, \dots, \alpha_i), (\alpha_{i+1}, \dots, \alpha_j), \dots, (\alpha_k, \dots, \alpha_m)$$

of an antichain $\alpha = (\alpha_1, \dots, \alpha_m)$ are the connected sub-antichains of α for which the supports of the union of the roots in any two distinct components are disjoint. For instance, the antichain $(r_1 + r_2, r_2 + r_3, r_4)$ has the connected components $(r_1 + r_2, r_2 + r_3)$ and r_4 . We will use lower and upper arcs to match two roots in a connected antichain in a geometric manner. Two roots linked by a lower [respectively upper] arc are said to be l-linked [respectively, u-linked]. In what follows we will identify each root with the correspondent permutation.

Definition 4.2. Define the map f from the set $\text{NN}(\Phi)$ into $\text{NC}(\Phi)$ recursively as follows. When α_1 is a positive root we set $f(\alpha_1) := \alpha_1$. If $\alpha = (\alpha_1, \dots, \alpha_m)$ is a connected antichain with $m > 2$, we have two cases:

(a) If there are no double roots in the antichain, define

$$f(\alpha) := \left(\bigcup_{k=1}^m \alpha_k \right) f(\bar{\alpha}_2, \dots, \bar{\alpha}_m),$$

where $\bar{\alpha}_k = \alpha_{k-1} \cap \alpha_k$ for $k = 2, \dots, m$.

(b) Assume now that $\alpha_1, \dots, \alpha_\ell$ have double roots, for some $\ell \geq 1$, and $\alpha_{\ell+1}, \dots, \alpha_m$ have none. Let $\Gamma_d := (\alpha_1, \dots, \alpha_\ell)$ and $\Gamma := (\alpha_{\ell+1}, \dots, \alpha_m)$. We start by introducing l-links as follows.

Let m' be the largest index of elements in Γ such that the following holds: $\alpha_{m'}$ has a first index $i \neq 1$, so that there is a rightmost element, say α_k , of Γ_d which has a term $2r_i$. If there is such an integer m' , l-link α_k with $\alpha_{m'}$. Then, ignore α_k and $\alpha_{m'}$ and proceed with the remaining roots as before. This procedure terminates after a finite number of steps (and not all elements of α need to be l-linked).

Next proceed by introducing u-links in α . The starting point of u-links, which we consider drawn from right to left, will be elements in Γ that have no first index 1 and are not l-linked. We will refer to these elements as admissible roots. So, let m' be the smallest integer such that the following holds: $\alpha_{m'}$ is an admissible root with first index $i \neq 1$ so that there is a leftmost element, say α_k which has r_i or $2r_i$ in its support and is not yet u-linked to an element on its right. If there is such an integer m' , u-link α_k with $\alpha_{m'}$. Remove $\alpha_{m'}$ from the set of admissible roots and proceed as before. Again this process terminates after a finite number of steps.

Finally, let $T = \{t_1 < \dots < t_p\}$ be the collection of all last double indices of the roots in Γ_d not l-linked, and all the last indices of the roots in α not u-linked to an element on its right. Then, define

$$f(\alpha) := \pi_1 \cdots \pi_\ell \pi_0 \theta_1 \cdots \theta_q f(\theta_{q+1}, \dots, \theta_s),$$

where for $j = 1, \dots, \ell$, $\pi_j = 2r_1 + \dots + 2r_{j'} + r_{j'+1} + \dots + r_{j''}$, with j' and j'' respectively the leftmost and rightmost integers in T not considered yet; π_0 is either the root $r_1 + \dots + r_{i_j}$, if the first index of $\alpha_{\ell+1}$ is 1, with i_j the only integer in T not used yet for defining the roots π_j , or the identity otherwise; each θ_j , $j = 1, \dots, q$ is the d-intersection of l-linked roots, starting from the rightmost one in Γ_d , and each θ_j , $j = q+1, \dots, s$ is the intersection of u-linked roots, starting from the leftmost one in Γ .

(c) For the general case, if $(\alpha_1, \dots, \alpha_i), (\alpha_{i+1}, \dots, \alpha_j), \dots, (\alpha_k, \dots, \alpha_m)$ are the connected components of $(\alpha_1, \dots, \alpha_m)$, let

$$f(\alpha_1, \dots, \alpha_m) := f(\alpha_1, \dots, \alpha_i) f(\alpha_{i+1}, \dots, \alpha_j) \cdots f(\alpha_k, \dots, \alpha_m).$$

Remark 4.1. (i) Notice that in the type A case, the map f is defined only by conditions (a) and (c) of the above definition. Also, note that if all roots in α have double roots then condition (b) is vacuous and the map f reduces to the identity map. We point out that the number of roots in $f(\alpha)$ is equal to the number of roots in the antichain α .


(ii) The sequence $(\bar{\alpha}_2, \dots, \bar{\alpha}_m)$ obtained in step (a) is a (not necessarily connected) antichain. It is easy to check that after all l-links and all u-links are settled, the set T has an odd number of elements if and only if the first index of $\alpha_{\ell+1}$ is 1. Thus, the root π_0 given in condition (b) is well defined.

We will show that f establishes a bijection between the sets $\text{NN}(\Psi)$ and $\text{NC}(\Psi)$, for $\Psi = A_{n-1}, B_n$ or C_n . Before, however, we present some examples.

Example 4.2. Consider the antichain $\alpha = (r_1 + r_2, r_2 + r_3, r_3 + r_4 + r_5, r_4 + r_5 + r_6, r_5 + r_6 + r_7)$ in the root poset of type A_7 , corresponding to the permutation (136)(247)(58) in the symmetric group \mathfrak{S}_8 . Applying the map f to α , we get the noncrossing partition

$$\begin{aligned} f(\alpha) &= (r_1 + \cdots + r_7) f(r_2, r_3, r_4 + r_5, r_5 + r_6) \\ &= (r_1 + \cdots + r_7) r_2 r_3 f(r_4 + r_5, r_5 + r_6) \\ &= (r_1 + \cdots + r_7) r_2 r_3 (r_4 + r_5 + r_6) r_5 \\ &\equiv (18)(2347)(56), \end{aligned}$$

whose graphical representation is given below:



Example 4.3. Consider now the antichain $\alpha = (\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5)$ in the root poset B_9 , where

$$\begin{aligned} \alpha_1 &= 2r_1 + 2r_2 + 2r_3 + 2r_4 + r_5, & \alpha_2 &= 2r_1 + 2r_2 + r_3 + r_4 + r_5 + r_6, \\ \alpha_3 &= r_1 + r_2 + r_3 + r_4 + r_5 + r_6 + r_7, & \alpha_4 &= r_3 + r_4 + r_5 + r_6 + r_7 + r_8, \\ \alpha_5 &= r_4 + r_5 + r_6 + r_7 + r_8 + r_9. \end{aligned}$$


Following definition 4.2, we get the l-links and the u-links shown below:

$$\alpha = (\alpha_1, \overbrace{\alpha_2, \alpha_3, \alpha_4}^{\quad}, \alpha_5) \quad .$$

Therefore, $T = \{2, 6, 7, 8, 9\}$ and the application of f to α yields:

$$\begin{aligned} f(\alpha) &= (2r_1 + 2r_2 + r_3 + \cdots + r_9)(2r_1 + \cdots + 2r_6 + r_7 + r_8) \cdot \\ &\quad \cdot (r_1 + \cdots + r_7)r_4 f(r_3 + r_4 + r_5) \\ &\equiv (2, -9)(-2, 9)(6, -8)(-6, 8)(7, -7)(3, 4)(-3, -4)(2, 5)(-2, -5) \\ &= (2, 5, -9)(-2, -5, 9)(6, -8)(-6, 8)(7, -7)(3, 4)(-3, -4). \end{aligned}$$

The image $f(\alpha)$ is a noncrossing partition in $[\pm 9]$, as we may check in its representation:



Example 4.4. For a final example, consider the antichain $\alpha = (\alpha_1, \dots, \alpha_6)$ in the root poset of type B_{11} , where

$$\begin{aligned} \alpha_1 &= 2r_1 + 2r_2 + 2r_3 + 2r_4 + 2r_5 + r_6, \\ \alpha_2 &= 2r_1 + 2r_2 + 2r_3 + 2r_4 + r_5 + r_6 + r_7, \\ \alpha_3 &= 2r_1 + 2r_2 + 2r_3 + r_4 + r_5 + r_6 + r_7 + r_8, \\ \alpha_4 &= r_2 + r_3 + r_4 + r_5 + r_6 + r_7 + r_8 + r_9, \\ \alpha_5 &= r_5 + r_6 + r_7 + r_8 + r_9 + r_{10}, \text{ and} \\ \alpha_6 &= r_7 + r_8 + r_9 + r_{10} + r_{11}. \end{aligned}$$

The l-links and u-links are shown below, so that $T = \{4, 6, 8, 9, 10, 11\}$:

$$\alpha = (\alpha_1, \overbrace{\alpha_2, \alpha_3, \alpha_4}^{\quad}, \underbrace{\alpha_5, \alpha_6}_{\quad}) \quad .$$

Therefore, the application of f to α gives $f(\alpha) = \pi_1\pi_2\pi_3(r_2 + r_3)(r_5)f(r_7)$, where

$$\pi_1 = 2r_1 + 2r_2 + 2r_3 + 2r_4 + r_5 + r_6 + r_7 + r_8 + r_9 + r_{10} + r_{11},$$


$$\pi_2 = 2r_1 + 2r_2 + 2r_3 + 2r_4 + 2r_5 + 2r_6 + r_7 + r_8 + r_9 + r_{10}, \text{ and}$$

$$\pi_3 = 2r_1 + 2r_2 + 2r_3 + 2r_4 + 2r_5 + 2r_6 + 2r_7 + 2r_8 + r_9.$$

Thus, $f(\alpha)$ is the noncrossing partition

$$\begin{aligned} f(\alpha) &= (4, -11)(-4, 11)(6, -10)(-6, 10)(8, -9)(-8, 9)(1, 3)(-1, -3)(4, 5) \cdot \\ &\quad \cdot (-4, -5)(6, 7)(-6, -7) \\ &= (4, 5, -11)(-4, -5, 11)(6, 7, -10)(-6, -7, 10)(8, -9)(-8, 9)(1, 3)(-1, -3) \end{aligned}$$

represented below:



Lemma 4.1. *If $\alpha \in \text{NN}(B_n)$ then $f(\alpha) \in \text{NC}(B_n)$, and $\mathfrak{T}(\alpha) = \mathfrak{T}(f(\alpha))$.*

Proof: Let $\alpha = (\alpha_1, \dots, \alpha_m)$ be an antichain in the root poset of type B_n , and let $(\alpha_1, \dots, \alpha_w)$ be its first connected component. Start by assuming that none of the positive roots in α have the simple root r_1 nor the double root $2r_1$. We will use induction on $m \geq 1$ to show that in this case $f(\alpha)$ is a noncrossing partition on the set $\{i - 1, \dots, q, -(i - 1), \dots, -q\}$, where i is the first index of α_1 and q is the last index of α_m , and such that each positive integer is sent to another positive integer. The result is clear when $m = 1$. So, let $m \geq 2$ and assume the result for antichains of length less than, or equal to $m - 1$. Then, we may write

$$f(\alpha) = \left(\bigcup_{k=1}^w \alpha_k \right) f(\bar{\alpha}_2, \dots, \bar{\alpha}_w) f(\alpha_{w+1}, \dots, \alpha_m),$$

where each $\bar{\alpha}_k = \alpha_{k-1} \cap \alpha_k$, for $k = 2, \dots, w$. By the inductive step, $f(\bar{\alpha}_2, \dots, \bar{\alpha}_w) \equiv \pi_1$ and $f(\alpha_{w+1}, \dots, \alpha_m) \equiv \pi_2$ are noncrossing partitions on the sets

$$\{a-1, \dots, b, -(a-1), \dots, -b\} \text{ and } \{p-1, \dots, q, -(p-1), \dots, -q\},$$

respectively, where a and p are the first indices of $\bar{\alpha}_2$ and α_{w+1} , respectively, and b and q are the last indices of $\bar{\alpha}_w$ and α_m , respectively. Moreover, all positive integers are sent to positive ones by π_1 and π_2 . Denoting by j the last index of α_w , we get $\bigcup_{k=1}^w \alpha_k = r_i + \dots + r_j \equiv (i-1, j)(-(i-1), -j)$ with $i-1 < a-1 < b < j \leq p-1 < q$. Therefore

$$f(\alpha) \equiv (i-1, j)(-(i-1), -j)\pi_1\pi_2$$

is a noncrossing partition on the set $\{i-1, \dots, q, -(i-1), \dots, -q\}$ sending each positive integer to another positive integer.

Note that for the rest of the proof, we may assume without loss of generality that α is connected, since none of the connected components of an antichain, except possible for the first one, have double roots, and therefore their images are noncrossing partitions sending each positive integer to another positive integer.

Suppose now that the first element of α_1 is r_1 . We will show that $f(\alpha)$ is a noncrossing partition on the set $\{i-1, \dots, q, -(i-1), \dots, -q\}$, where i is the first index of α_2 and q is the last index of α_m , and such that one and only one positive integer is sent to a negative one. The result is certainly true for $m = 1$, and when $m > 1$ we have

$$f(\alpha) = \left(\bigcup_{k=1}^m \alpha_k \right) f(\bar{\alpha}_2, \dots, \bar{\alpha}_m),$$

where $\bigcup_{k=1}^m \alpha_k \equiv (q, -q)$, and $\bar{\alpha}_k = \alpha_{k-1} \cap \alpha_k$ for $k = 2, \dots, m$. By the previous case, $f(\bar{\alpha}_2, \dots, \bar{\alpha}_m) \equiv \pi$ is a noncrossing partition on the set $\{i-1, \dots, j, -(i-1), \dots, -j\}$, with i the first index of α_2 and $j < q$ the last index of α_{m-1} . Therefore, $f(\alpha) \equiv (q, -q)\pi$ is a noncrossing partition satisfying the desired conditions.

Next, assume that α satisfies condition (b) of definition 4.2, and consider its image

$$f(\alpha) = \pi_1 \cdots \pi_\ell \pi_0 \theta_1 \cdots \theta_q f(\theta_{q+1}, \dots, \theta_s).$$

By the construction of the set T , it follows that each D_{α_j} , $j = 1, \dots, \ell$, is contained in D_{π_i} , for some $i = 1, \dots, \ell$, and that $\pi_1 \cdots \pi_\ell \pi_0$ is a noncrossing partition, sending each nonfixed positive integer to a negative one. Note also that the support of each θ_j , $j = 1, \dots, q$, is contained in some D_{α_i} , $i = 1, \dots, \ell$, and therefore, in some D_{π_i} , $i = 1, \dots, \ell$. Moreover, the supports of any two roots θ_i and θ_j , $1 \leq i, j \leq q$, are either disjoint, or one of them is contained into the other one. Therefore $\theta_1 \cdots \theta_q$ is a noncrossing partition sending each nonfixed positive integer into another positive integer. By the previous cases, $f(\theta_{q+1}, \dots, \theta_s)$ is also a noncrossing partition sending each nonfixed positive integer into another positive integer. Again by the construction of the set T , we find that the support of each θ_j , $j = q+1, \dots, s$, is either contained in some D_{π_i} , or it does not intersect D_{π_ℓ} . For each $j = 1, \dots, q$ and $i = q+1, \dots, s$, either we have $\text{supp}(\theta_i) \cap \text{supp}(\theta_j) = \emptyset$, or $\text{supp}(\theta_i) \supseteq \text{supp}(\theta_j)$, this last case happening when θ_i arises from the intersection of two u-linked roots $\alpha_u \in \Gamma_d$ and $\alpha_v \in \Gamma$, and there is some $\alpha_{v+k} \in \Gamma$, $k \geq 1$, l-linked to α_u , whose d-intersection gives θ_j . Therefore, it follows that $f(\alpha)$ is noncrossing.

To see that $\mathfrak{T}(\alpha) = \mathfrak{T}(f(\alpha))$, notice that if $a \in \text{op}(\alpha)$, then α must have a root $r_{a+1} + \cdots + r_b$, for some $b > a + 1$, and cannot have neither a root with last double index equal to a nor a root with last index equal to a . By its construction, the same is true for the sequence $f(\alpha)$, and therefore $a \in \text{op}(f(\alpha))$. Assume now that $a \in \text{tr}(\alpha)$. Then, a must appear either as the last index or last double index of a root and $a + 1$ as the first index of another root. Again the same will happen in the sequence $f(\alpha)$, and thus $a \in \text{tr}(f(\alpha))$. It follows that $\mathfrak{T}(\alpha) = \mathfrak{T}(f(\alpha))$. ■

With some minor adaptations, the proof of lemma 4.1, in the case where neither the simple root r_1 nor the double root $2r_1$ are present in α , gives the type A analog of the previous result, and its type C analog is obtained through the bijection τ .

Corollary 4.2. *If $\alpha \in \text{NN}(\Psi)$ then $f(\alpha) \in \text{NC}(\Psi)$, for $\Psi = A_{n-1}$ or C_n , and $\mathfrak{T}(\alpha) = \mathfrak{T}(f(\alpha))$.*

We will now construct the inverse function of f , thus showing that f establishes a bijection between the sets $\text{NN}(\Psi)$ and $\text{NC}(\Psi)$, for $\Psi = A_{n-1}, B_n$ or C_n . For that propose, recall the following property.

Lemma 4.3. *Two distinct transpositions (a, b) and (i, j) in \mathfrak{S}_n commute if and only if the sets $\{i, j\}$ and $\{a, b\}$ are disjoint.*

If $\pi_1 \cdots \pi_p$ is the cycle structure of a signed permutation π , then for each cycle $\pi_i = (i_j \cdots k)$ there is another cycle $\pi_j = (-i - j \cdots -k)$. Denote by π'_i the cycle in $\{\pi_i, \pi_j\}$ having the smallest positive integer (when $\pi_i = \pi_j$ then π'_i is just π_i), and call positive cycle structure to the subword of $\pi_1 \cdots \pi_p$ formed by the cycles π'_i . Extend this definition to permutations in \mathfrak{S}_n by identifying positive cycle structure with cycle structure.

Theorem 4.4. *The map f is a bijection between the sets $\text{NN}(\Psi)$ and $\text{NC}(\Psi)$, for $\Psi = A_{n-1}, B_n$ or C_n , which preserves the triples $(\text{op}(\pi), \text{cl}(\pi), \text{tr}(\pi))$.*

Proof: We will construct the inverse map $g : \text{NC}(\Psi) \rightarrow \text{NN}(\Psi)$ of f . Given $\pi \in \text{NC}(\Psi)$, let $\pi_1 \cdots \pi_s$ be its positive cycle structure. Replace each cycle $\pi_i = (i_1 i_2 \cdots i_k)$ by $(i_1 i_2)(i_2 i_3) \cdots (i_{k-1} i_k)$, if $i_\ell > 0$ for $\ell = 1, \dots, k$, or by

$$\pi_i = (i_1 i_{j+1})(i_1 i_2)(i_2 i_3) \cdots (i_{j-1} i_j)(i_{j+1} i_{j+2}) \cdots (i_{k-1} i_k),$$

if $i_\ell > 0$ for $\ell = 1, \dots, j$, and $i_\ell < 0$ for $\ell = j+1, \dots, k$. Next, baring in mind lemma 4.3 and recalling that π is noncrossing, move all transpositions (i, j) , with $i > 0$ and $j < 0$ (if any), to the leftmost positions and order them by its least positive element, and order all remaining transpositions (i, j) , with $i, j > 0$, by its least positive integer. Replace each transposition (ij) by its correspondent root in the root system of type Ψ , and let

$$(\alpha_1, \dots, \alpha_k)(\alpha_{k+1}, \dots, \alpha_\ell) \cdots (\alpha_m, \dots, \alpha_n) \quad (4.1)$$

be the correspondent sequence of roots, divided by its connected components. Note that given two distinct roots in (4.1), the sets formed by the first and last indices, if there are no double roots, or by the last and last double indices, otherwise, are clearly disjoint.

We start by considering that the sequence (4.1) has only one connected component $(\alpha_1, \dots, \alpha_k)$. Let $\Gamma_d = (\alpha_1, \dots, \alpha_r)$ be the subsequence formed by the roots having double roots, and denote by $\Gamma = (\alpha_{r+1}, \dots, \alpha_k)$ the remaining subsequence. Define $\Gamma' = \Gamma'_d = \emptyset$. If Γ_d is not empty and $r \neq k$, apply the following algorithm:

Let $\bar{\Gamma}$ be the subsequence of Γ obtained by striking out the root α_{r+1} if its first index is 1. While $\bar{\Gamma} \neq \emptyset$, repeat the following steps:

(i) Let α_i be the leftmost root in $\bar{\Gamma}$ and check if $\text{supp}(\alpha_i) \subseteq D_{\alpha_j}$, for some $\alpha_j \in \Gamma_d \setminus \Gamma'_d$.

(ii) If so, let α_{i_j} be the rightmost root in $\Gamma_d \setminus \Gamma'_d$ with this property. Update Γ' by including in it the rightmost root $\bar{\alpha}$ of $\bar{\Gamma}$ whose support is contained in

$\text{supp}(\alpha_i)$. Update $\bar{\Gamma}$ by striking out the root $\bar{\alpha}$ and update Γ'_d by including in this set the root α_{i_j} .

(iii) Otherwise, update $\bar{\Gamma}$ by striking out the root α_i .

Next, let $T = \{t_1 > \cdots > t_r\}$ be the set formed by all last double indices of the roots in $\Gamma_d \setminus \Gamma'_d$ and by the last indices of the roots in Γ' ; let $F_{st} = \{f_{r+1} < \cdots < f_k\}$ be the set formed by the first indices of the roots in Γ , and let $L_{st} = \{\ell_1 < \cdots < \ell_k\}$ be the set formed by the last indices of the roots in $(\Gamma \setminus \Gamma') \cup \Gamma_d$ and by the last double indices of the roots in Γ'_d . By this construction, we have $f_i < \ell_i$ for $i = 1, \dots, r$, and $f_i < \ell_i$, for $i = r+1, \dots, k$. Then, define

$$g(\pi) = (\bar{\alpha}_1, \dots, \bar{\alpha}_k),$$

where for $i = 1, \dots, r$, $\bar{\alpha}_i = 2r_1 + \cdots + 2r_{t_i} + r_{t_i+1} + \cdots + r_{\ell_i}$, and for $i = r+1, \dots, k$, $\bar{\alpha}_i = r_{f_i} + \cdots + r_{\ell_i}$.

For the general case define

$$g(\pi) = g(\alpha_1, \dots, \alpha_k)g(\alpha_{k+1}, \dots, \alpha_\ell) \cdots g(\alpha_m, \dots, \alpha_n).$$

It is clear from this construction that $g(\pi)$ is an antichain in the root poset of type Ψ . Moreover, a closer look at the construction of the map f shows that g is the inverse of f . Thus, f (and g) establishes a bijection between nonnesting and noncrossing partitions of types A, B and C . ■

In the following examples we illustrate the application of the map g .

Example 4.5. Consider the cycle structure of the noncrossing partition $\pi = (18)(2347)(56)$ in the symmetric group \mathfrak{S}_8 used in example 4.2. Following the proof of theorem 4.4, write

$$\begin{aligned} \pi &\equiv (18)(2347)(56) \\ &= (18)(23)(34)(47)(56) \\ &\equiv (r_1 + \cdots + r_7)r_2r_3(r_4 + r_5 + r_6)r_5 \end{aligned}$$

Note that π has only one connected component, and there are no double roots. Next define the sets

$$F_{st} = \{1, 2, 3, 4, 5\}, \text{ and } L_{st} = \{2, 3, 5, 6, 7\}.$$

Thus, we find that the image of π by the map g is the antichain

$$g(\pi) = (r_1 + r_2, r_2 + r_3, r_3 + r_4 + r_5, r_4 + r_5 + r_6, r_5 + r_6 + r_7).$$

Example 4.6. Consider now the noncrossing partition

$$\pi = (2, 5, -9)(-2, -5, 9)(6, -8)(-6, 8)(7, -7)(3, 4)(-3, -4)$$

obtained in example 4.3. Its positive cycle structure is

$$(2, -9)(2, 5)(6, -8)(7, -7)(3, 4) = (2, -9)(6, -8)(7, -7)(2, 5)(3, 4),$$

and thus we get

$$\pi \equiv (2r_1 + 2r_2 + r_3 + \cdots + r_9, 2r_1 + \cdots + 2r_6 + r_7 + r_8, r_1 + \cdots + r_7, r_3 + r_4 + r_5, r_4).$$

Next, construct the sets

$$\begin{aligned} T &= \{4, 2\}, & F_{st} &= \{1, 3, 4\} \\ L_{st} &= \{5, 6, 7, 8, 9\}. \end{aligned}$$

Therefore, the image of π by the map g is the antichain

$$(2r_1 + \cdots + 2r_4 + r_5, 2r_1 + 2r_2 + r_3 + \cdots + r_6, r_1 + \cdots + r_7, r_3 + \cdots + r_8, r_4 + \cdots + r_9).$$

Finally, in the next result we prove that the map f generalizes the bijection that locally converts each crossing to a nesting.

Theorem 4.5. *When restricted to the type A_{n-1} case, the map f coincides with the L -map.*

Proof: Let $\alpha = (\alpha_1, \dots, \alpha_m)$ be an antichain in the root poset of type A_{n-1} . The result will be handled by induction over $m \geq 1$. Without loss of generality, we may assume that α is connected, since otherwise there is an integer $1 < k < n - 1$ such that each integer less (resp. greater) than k is sent by α to an integer that still is less (resp. greater) than k . Therefore, the same happens with the image of α by either the map f or the L -map.

The result is vacuous when $m = 1$, and when $m = 2$, the only connected nonnesting partition which does not stay invariant under the maps f and L is $\alpha = (r_i + \cdots + r_{i'})(r_j + \cdots + r_{j'})$, for some integers $1 \leq i < j < i' < j' \leq n - 1$. In this case, the equality between f and the L -map is obvious. So, let $m > 2$ and assume the result for antichains of length $\leq m - 1$. Let i and j be, respectively, the first and last indices of α_1 and α_m . Then,

$$f(\alpha) = (r_i + \cdots + r_j)f(\bar{\alpha}_2, \dots, \bar{\alpha}_m),$$

where each $\bar{\alpha}_k = \alpha_{k-1} \cap \alpha_k$ for $k \geq 2$, and the antichain $(\bar{\alpha}_2, \dots, \bar{\alpha}_m)$ is clearly nonnesting, and not necessarily connected. By the inductive step, $f(\bar{\alpha}_2, \dots, \bar{\alpha}_m) = L(\bar{\alpha}_2, \dots, \bar{\alpha}_m)$. Moreover, note that converting, from left to

right, each local crossing between the first root and the leftmost root in α whose arcs cross, into a nesting gives, precisely,


$$(r_i + \cdots + r_j)L(\bar{\alpha}_2, \dots, \bar{\alpha}_m),$$

and this operation may be considered the first step of the L-map. Thus, we find that $f(\alpha) = L(\alpha)$. ■


Example 4.7. Consider the antichain $\alpha = (r_1 + r_2 + r_3, r_2 + r_3 + r_4 + r_5, r_3 + r_4 + r_5 + r_6, r_5 + r_6 + r_7)$ in the root poset of type A_7 . Applying the map f we get

$$\begin{aligned} f(\alpha) &= (r_1 + \cdots + r_7)f(r_2 + r_3, r_3 + r_4 + r_5, r_5 + r_6) \\ &= (r_1 + \cdots + r_7)(r_2 + r_3 + r_4 + r_5 + r_6)f(r_3, r_5) \\ &= (r_1 + \cdots + r_7)(r_2 + r_3 + r_4 + r_5 + r_6)r_3r_5 \equiv (18)(27)(34)(56). \end{aligned}$$

On the other hand, applying the L-map to each crossing between the first root and the leftmost root in α whose arcs cross, we get successively



Thus, in the first step of the L-map, we get $L(\alpha) = (r_1 + \cdots + r_7)L(r_2 + r_3, r_3+r_4+r_5, r_5+r_6)$. Continuing the application of the L-map, now replacing, by a nesting, each crossing between the second root and the leftmost root in α whose arcs cross, we get



and therefore, we have $L(\alpha) = (r_1 + \cdots + r_7)(r_2 + r_3 + r_4 + r_5 + r_6)r_3r_5 = f(\alpha)$.

5. Concluding remarks

The existing three bijections [12, 10, 5] between noncrossing and nonnesting partitions are all distinct, and preserves different statistics. While our bijection preserves the triples (op, cl, tr) formed by the openers, closers and transients of the partitions, for the types A, B and C , and therefore also the number of blocks, the one by Alex Fink and Benjamin Giraldo preserves

the type of the partitions but not the triples (op, cl, tr) . For the types A and B , Stump's bijection does not preserve neither the type nor the triples (op, cl, tr) .

Acknowledgements. The author thanks Christian Krattenthaler and Alexander Kovacec for useful comments and suggestions.

References

- [1] D. Armstrong, *Generalized Noncrossing Partitions and Combinatorics of Coxeter Groups*, preprint, arXiv:math/0611106v2 [math.CO].
- [2] C.A. Athanasiadis, *On noncrossing and nonnesting partitions for classical reflection groups*, Elec. J. Comb. 5 (1998), Research Paper 42, 16pp (electronic).
- [3] C.A. Athanasiadis and V. Reiner, *Noncrossing partitions for the group D_n* , SIAM J. Discrete Math. 18 (2004), 397-417.
- [4] P. Biane, *Some properties of crossings and partitions*, Discrete Math. 175 (1997), 41-53.
- [5] A. Fink and B. I. Giraldo, *Bijections between noncrossing and nonnesting partitions for classical reflection groups*, preprint, arXiv:math/0810.2613v1 [math.CO].
- [6] L.C. Grove and C.T. Benson, *Finite Reflection Groups*, Springer-Verlag, New York, 1996.
- [7] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics, vol. 29, Cambridge Univ. Press, Cambridge, 1990.
- [8] A. Kasraoui and J. Zeng, *Distribution of crossings, nestings and alignments of two edges in matchings and partitions*, preprint, arXiv:math/0601081v1 [math.CO].
- [9] G. Kreweras, *Sur les partitions non-croisées d'un cycle*, Discrete Math., 1 (1972), pp. 333-350.
- [10] R. Mamede, *A bijection between noncrossing and nonnesting partitions of types A and B* , preprint, arXiv:0810.1422v1 [math.CO].
- [11] V. Reiner, *Non-crossing partitions for classical reflection groups*, Discrete Math. 177 (1997), 195-222.
- [12] C. Stump, *Non-crossing partitions, non-nesting partitions and Coxeter sortable elements in types A and B* , preprint, arXiv:0808.2822v1 [math.CO].

RICARDO MAMEDE

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, 3001-454 COIMBRA, PORTUGAL

E-mail address: mamede@mat.uc.pt