

INVARIANT STAR PRODUCTS ON NONDEGENERATE TRIANGULAR LIE BIALGEBRAS OVER POLYNOMIAL RINGS

CARLOS MORENO AND JOANA TELES

ABSTRACT: There is a bijection between the set of equivalent classes of Invariant Star Products on a nondegenerate triangular finite dimensional Lie bialgebra $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t}, r_t)$ over the ring $\mathbb{K}[[t]]$ and the set $\hbar H^2(\mathfrak{a}_t)[[\hbar]]$. As a consequence, in case $t = \hbar$ we obtain a set of triangular Hopf Quantized Universal enveloping algebras which are quantizations of a deformation algebra $(\mathfrak{a}_\hbar, [;]_{\mathfrak{a}_\hbar}, r_\hbar)$. These algebras are isomorphic if and only if the corresponding star products are equivalent.

KEYWORDS: Quantization of Lie bialgebras, Star Products, Quantized Universal Enveloping algebras.

AMS SUBJECT CLASSIFICATION (2000): 16W30; 17B62; 17B37; 46L65; 53D55.

1. Introduction

1) Let $(\mathfrak{a}, [;]_{\mathfrak{a}})$ be a *finite dimensional* Lie algebra over a field \mathbb{K} of characteristic zero. Let $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t})$ be a Lie algebra over the ring $\mathbb{K}_t \equiv \mathbb{K}[[t]]$ of formal power series in the indeterminate t which is a *deformation algebra* [5] of $(\mathfrak{a}, [;]_{\mathfrak{a}})$, i.e., as a \mathbb{K}_t -module \mathfrak{a}_t is $\mathfrak{a}[[t]]$ and $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t})$ is $(\mathfrak{a}, [;]_{\mathfrak{a}})$ modulo t . Let u be another indeterminate. Let $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}})$, $\mathfrak{a}_{t,u} = \mathfrak{a}_t[[u]]$, be the Lie algebra obtained by extension of the ring of scalars $\mathbb{K}[[t]] \longrightarrow \mathbb{K}_t[[u]]$ ($\mathbb{K}_{t,u} \equiv \mathbb{K}_t[[u]] \equiv \mathbb{K}[[t, u]]$) from the Lie algebra $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t})$. Let \hbar be a third indeterminate and consider the ring $\mathbb{K}_{t,u}[[\hbar]]$. Let $(\mathfrak{a}_t[[\hbar]], [;]_{\mathfrak{a}_t[[\hbar]]})$ be the Lie algebra over $\mathbb{K}_t[[\hbar]]$ defined as before in case of the indeterminate u .

$r_1 \in \mathfrak{a} \wedge \mathfrak{a}$ will be a given nondegenerate solution of the Yang Baxter Equation (YBE), i.e. $[r_1, r_1]_{\mathfrak{a}} = 0$, on the Lie algebra $(\mathfrak{a}, [;]_{\mathfrak{a}})$. By nondegenerate we mean $\text{rang}(r_1) = \dim \mathfrak{a}$.

The symbol $r_t = \sum_{l \geq 1} r_l \cdot t^{(l-1)} \in \mathfrak{a}_t \wedge \mathfrak{a}_t$, $r_l \in \mathfrak{a} \wedge \mathfrak{a}$, $l \in \mathbb{N}$, will denote a nondegenerate solution of YBE on the Lie algebra $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t})$.

Received January 16, 2008.

The second author has been partially supported by CMUC and FCT - project POCI/MAT/58452/2004.

The symbol $r_{t,u} = \sum_{l \geq 1} r_{t,l} \cdot u^{(l-1)} \in \mathfrak{a}_t[[u]] \wedge \mathfrak{a}_t[[u]]$, $r_{t,l} \in \mathfrak{a}_t \wedge \mathfrak{a}_t$, $l \in \mathbb{N}$, and $r_{t,1} = r_t$ will denote a nondegenerate solution of YBE on the Lie algebra $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}})$ over the ring $\mathbb{K}_t[[u]]$.

As $r_t \in \mathfrak{a}_t \wedge \mathfrak{a}_t$ is a solution of YBE on $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t})$ it defines the corresponding Poisson cohomology spaces $H_{P,r_t}^k(\mathfrak{a}_t)$. As r_t is nondegenerate, let $\mu_{r_t} : \Lambda(\mathfrak{a}_t) \longrightarrow \Lambda(\mathfrak{a}_t^*)$ be the corresponding isomorphism. Let $\mu_{r_t}(r_t) = \beta_t \in \mathfrak{a}_t^* \wedge \mathfrak{a}_t^*$ be the corresponding 2-cocycle in the Chevalley cohomology of $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t})$ with the trivial action of \mathfrak{a}_t on K_t and $H^l(\mathfrak{a}_t)$ be the corresponding cohomological modules. Let $\bar{\mu}_{r_t} : H_{P,r_t}^l(\mathfrak{a}_t) \longrightarrow H^l(\mathfrak{a}_t)$ be the induced mapping on cohomology spaces. Similar meanings have the symbols $\mu_{r_{t,u}}$, μ_{r_1} .

2) $r \in (\mathfrak{a} \oplus \mathfrak{a}^*) \otimes (\mathfrak{a} \oplus \mathfrak{a}^*)$ will denote the *canonical element*. Then $r = (e_i, 0) \otimes (0, e^i)$ in any pair of dual basis. The symbol d_c will denote the coboundary of the Chevalley cohomology of any Lie algebra *with values in the adjoint representation*. The symbol $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*, [;]_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*}, \varepsilon_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*} = d_c(t, u)r)$ will denote the *quasitriangular double Lie bialgebra* of the *nondegenerate triangular Lie bialgebra* $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$. The element r is a solution of the YBE on $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}})$, and defines the symmetric element $\Omega = r + \sigma(r)$ where σ is the cycle (12). Ω is $ad_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*}$ -invariant and it satisfies the usual infinitesimal tress relations.

3) We fix a Lie associator $\Phi = \exp P(\hbar t_{12}, \hbar t_{23})$ over \mathbb{K} , [6, 7].

4) $(\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*), \Delta_0^{r_{t,u}})$ will denote the universal enveloping algebra of the Lie algebra $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*, [;]_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*})$. We do not specify its product, unit or antipode.

A theorem in [6] allow us to prove the existence of a quasitriangular quasi-Hopf algebra $(\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)[[\hbar]], \Delta_0^{r_{t,u}}, \Phi_{r_{t,u}}, R_0^{r_{t,u}} = e^{\frac{\hbar}{2}\Omega})$ over the ring $\mathbb{K}_{t,u}[[\hbar]]$. See theorem 2.1 bellow.

Etingof-Kazhdan theory of quantization of Lie bialgebras [7] allows us to obtain an element $J_{r_{t,u}} \in (\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*) \otimes \mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*))[[\hbar]]$, verifying $J_{r_{t,u}} = 1 \otimes 1 + \frac{1}{2}r\hbar$ modulo \hbar^2 such that when twisting [5] the above quasitriangular Quasi-Hopf algebra via the element $J_{r_{t,u}}^{-1}$ we obtain a *quasitriangular Hopf algebra* $(\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)[[\hbar]], \Delta^{r_{t,u}}, R^{r_{t,u}})$ over the ring $\mathbb{K}_{t,u}[[\hbar]]$. We also write this algebra as $A_{(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)[[\hbar]], \Omega, J_{r_{t,u}}^{-1}}$.

5) From a theorem by Etingof-Kazhdan we can obtain an element $\tilde{J}_{r_{t,u}} = (\tilde{\pi}_{t,u} \otimes \tilde{\pi}_{t,u})J_{r_{t,u}} \in (\mathcal{U}\mathfrak{a}_{t,u} \otimes \mathcal{U}\mathfrak{a}_{t,u})[[\hbar]]$ verifying the condition $\tilde{J}_{r_{t,u}} = 1 \otimes 1 +$

$\frac{1}{2}r_{t,u}\hbar$ modulo \hbar^2 such that when twisting via the element $\tilde{J}_{r_{t,u}}^{-1}$ the *trivial* triangular Hopf algebra $(\mathcal{U}\mathfrak{a}_{t,u}[[\hbar]], \Delta_{\mathfrak{a}_{t,u}}, R_{\mathfrak{a}_{t,u}} = 1 \otimes 1)$ over the ring $\mathbb{K}_{t,u}[[\hbar]]$ one obtains a *triangular Hopf algebra* $(\mathcal{U}\mathfrak{a}_{t,u}[[\hbar]], \tilde{\Delta}_{\mathfrak{a}_{t,u}}, \tilde{R}_{\mathfrak{a}_{t,u}})$ over the ring $\mathbb{K}_{t,u}[[\hbar]]$. We will denote this algebra as $A_{\mathfrak{a}_{t,u}[[\hbar]], \tilde{J}_{r_{t,u}}^{-1}}$. The element $\tilde{J}_{r_{t,u}}$ is an Invariant Star Product on the nondegenerate triangular Lie bialgebra $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$ over the ring $\mathbb{K}_{t,u}$, see [1, 4, 11].

6) Now, *it has a meaning to put* $u = \hbar$ in every element appearing in the definition of the quasitriangular quasi-Hopf algebra, quasitriangular Hopf algebra, or triangular Hopf algebra over $\mathbb{K}_{t,u}[[\hbar]]$ considered in 4) and 5) above. In this way, we obtain, respectively, *i)* a quasitriangular quasi-Hopf algebra $(\mathcal{U}(\mathfrak{a}_{t,\hbar} \oplus \mathfrak{a}_{r_{t,\hbar}}^*)[[\hbar]], \Delta_0^{r_{t,\hbar}}, \Phi_{r_{t,\hbar}}, R_0^{r_{t,\hbar}} = e^{\frac{\hbar}{2}\Omega})$ over the ring $\mathbb{K}_t[[\hbar]]$; *ii)* a quasitriangular Hopf algebra $(\mathcal{U}(\mathfrak{a}_{t,\hbar} \oplus \mathfrak{a}_{r_{t,\hbar}}^*)[[\hbar]], \Delta^{r_{t,\hbar}}, R^{r_{t,\hbar}})$ over the ring $\mathbb{K}_t[[\hbar]]$; it can be obtained by a twist via the element $J_{r_{t,\hbar}} \in (\mathcal{U}(\mathfrak{a}_{t,\hbar} \oplus \mathfrak{a}_{r_{t,\hbar}}^*) \otimes \mathcal{U}(\mathfrak{a}_{t,\hbar} \oplus \mathfrak{a}_{r_{t,\hbar}}^*))[[\hbar]]$ from the one obtained in *i)*; and *iii)* a triangular Hopf algebra $(\mathfrak{a}_{t,\hbar}, \tilde{\Delta}_{\mathfrak{a}_{t,\hbar}}, \tilde{R}_{\mathfrak{a}_{t,\hbar}})$ over the ring $\mathfrak{a}_t[[\hbar]]$. We say that this algebra is a quantization of the pair (\mathfrak{a}_t, r_t) . It can be obtained by a twist via the element $\tilde{J}_{r_{t,\hbar}}^{-1} \in (\mathcal{U}\mathfrak{a}_{t,\hbar} \otimes \mathcal{U}\mathfrak{a}_{t,\hbar})[[\hbar]]$ from the *trivial* triangular Hopf algebra $(\mathcal{U}\mathfrak{a}_{t,\hbar}[[\hbar]], \Delta_{\mathfrak{a}_{t,\hbar}}, R_{\mathfrak{a}_{t,\hbar}} = 1 \otimes 1)$ over the ring $\mathbb{K}_t[[\hbar]]$.

7) The adjoint representation of a Lie group \mathbf{G} with Lie algebra $(\mathfrak{a}, [;]_{\mathfrak{a}})$ and $\mathbb{K} = \mathbb{R}$ induces a representation on the Chevalley complex $H^*(\mathfrak{a})$ that is *trivial*. This classical theorem inspired us for considering the Lie algebra isomorphisms in Section 5 that allow us to obtain, in Sections 6 and 7, the equivalence of Invariant Star Products. This equivalence allows us to obtain the corresponding isomorphisms for Hopf algebras.

No proofs of these results are given here. They will appear in a forthcoming article. References [15] and [14] are related with the subject of this paper.

2. Quantization of the quasitriangular Lie bialgebra $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*, [;]_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*}, \varepsilon_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*} = d_c(t, u)r)$ over the ring $\mathbb{K}_{t,u}$

1) Pentagon, hexagon properties of associators [6, 7] and the *ad*-invariance of Ω allow us [6] to obtain the following:

Theorem 2.1. *Let $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$ and $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*, [;]_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*}, \varepsilon_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*} = d_c(t,u)r)$ be as in 2) of Section 1. Consider the $\mathbb{K}_{t,u}[[\hbar]]$ -module $\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)[[\hbar]]$. The set*

$$\left(\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)[[\hbar]], \Delta_0^{r_{t,u}}, \Phi_{r_{t,u}}, R_0^{r_{t,u}} = e^{\frac{\hbar}{2}\Omega} \right)$$

is then a quasitriangular quasi-Hopf algebra over $\mathbb{K}_{t,u}[[\hbar]]$.

We do not specify the corresponding antipode. Its existence follows from Theorem 1.6 in [5] and specific forms for it may be obtained from Propositions 1.1 and 1.3 in [5]. We also do not specify product, unity or counity.

Definition 2.2. *We say that the quasitriangular quasi-Hopf algebra over $\mathbb{K}_{t,u}[[\hbar]]$ of Theorem 2.1 is a quantization of the pair $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*, \Omega)$ or that this pair is the classical limit of the quasitriangular quasi-Hopf algebra.*

2) Part 2) of the following theorem can be proved analogously to the corresponding one in [7] Part I. We only need to remark that $\mathbb{K}_{t,u}$ is a \mathbb{Q} -algebra and that the symmetric algebras of the $\mathbb{K}_{t,u}$ -modules $\mathfrak{a}_{t,u}$, $\mathfrak{a}_{r_{t,u}}^*$, $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)$ are isomorphic to the corresponding algebras of symmetric tensors [2]. Then we apply Corollary 3 of Theorem 1 of §2.8, Chapter III, of [3].

Theorem 2.3. *Let $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$ and $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*, [;]_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*}, \varepsilon_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*} = d_c(t,u)r)$ be as in the above theorem. Let $M(t,u)_{\pm}$ be the $\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*$ -modules with one generator 1_{\pm} and defined as follows: $M(t,u)_{+} = \mathcal{U}\mathfrak{a}_{r_{t,u}}^* \cdot 1_{+}$; $\mathcal{U}\mathfrak{a}_{t,u} \cdot 1_{+} = 0$ and $M(t,u)_{-} = \mathcal{U}\mathfrak{a}_{t,u} \cdot 1_{-}$; $\mathcal{U}\mathfrak{a}_{r_{t,u}}^* \cdot 1_{-} = 0$. Then*

1) *The equalities $i_{\pm}(1_{\pm}) = 1_{\pm} \otimes 1_{\pm}$ define unique $\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*$ -module morphisms $i_{\pm} : M(t,u)_{\pm} \longrightarrow M(t,u)_{\pm} \otimes_{\mathbb{K}_{t,u}} M(t,u)_{\pm}$.*

2) *The equality $\phi_{r_{t,u}}(1) = 1_{+} \otimes 1_{-}$ defines a unique $\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*$ -module morphism $\phi_{r_{t,u}} : \mathcal{U}(\mathfrak{a}_{r_{t,u}} \oplus \mathfrak{a}_{r_{t,u}}^*) \longrightarrow M(t,u)_{+} \otimes M(t,u)_{-}$. Moreover $\phi_{r_{t,u}}$ is an isomorphism.*

3) *There exists an element $J_{r_{t,u}} \in (\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)[[\hbar]])^{\hat{\otimes} 2}$ such that, when twisting via $J_{r_{t,u}}^{-1}$ the quasitriangular quasi-Hopf algebra considered in Theorem 2.1, one obtains a quasitriangular Hopf algebra, $(\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)[[\hbar]], \Delta^{r_{t,u}}, R^{r_{t,u}})$, over the ring $\mathbb{K}_{t,u}[[\hbar]]$. The element $J_{r_{t,u}}$ is given by*

$$J_{r_{t,u}} = (\phi_{r_{t,u}}^{-1} \otimes \phi_{r_{t,u}}^{-1})(\Phi_{1,2,34}^{-1} \circ \Phi_{2,3,4} \circ \sigma_{23} \circ e^{\frac{\hbar}{2}\Omega_{23}} \circ \Phi_{2,3,4}^{-1})$$

$$\Phi_{1,2,34} \circ (i_+ \otimes i_-)(\phi_{t,u}(1)),$$

and

$$\Delta^{r_{t,u}}(b) = J_{r_{t,u}}^{-1} \cdot_{r_{t,u}} \Delta_0^{r_{t,u}}(b) \cdot_{r_{t,u}} J_{r_{t,u}}; \quad R^{r_{t,u}} = \sigma(J_{r_{t,u}}^{-1}) \cdot_{r_{t,u}} e^{\frac{\hbar}{2}\Omega} \cdot_{r_{t,u}} J_{r_{t,u}}.$$

We also have $J_{r_{t,u}} = 1 \otimes 1 + \frac{1}{2}r\hbar \text{ mod } \hbar^2$ and $R^{r_{t,u}} = 1 \otimes 1 + r\hbar \text{ mod } \hbar^2$. The isomorphism $\Phi_{r_{t,u}}$ verifies the following equality

$$\Phi_{r_{t,u}} \cdot_{r_{t,u}} (\Delta_0^{r_{t,u}} \otimes id)(J_{r_{t,u}}) \cdot_{r_{t,u}} (J_{r_{t,u}} \otimes 1) = (1 \otimes \Delta_0^{r_{t,u}})(J_{r_{t,u}}) \cdot_{r_{t,u}} (1 \otimes J_{r_{t,u}}).$$

The products in these expressions are those of the enveloping algebra $\mathcal{U}\left((\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*) \otimes_{\mathbb{K}_{t,u}} \mathbb{K}_{t,u}[[\hbar]]\right) \equiv \mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*) \otimes_{\mathbb{K}_{t,u}} \mathbb{K}_{t,u}[[\hbar]]$ defined by extension of scalars $\mathbb{K}_{t,u} \longrightarrow \mathbb{K}_{t,u}[[\hbar]]$. This quasitriangular Hopf algebra over $\mathbb{K}_{t,u}[[\hbar]]$ will be denoted by $A_{(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)[[\hbar]], \Omega, J_{r_{t,u}}^{-1}}$.

Definition 2.4. We say that the quasitriangular Hopf algebra over $\mathbb{K}_{t,u}[[\hbar]]$ considered in 3) of Theorem 2.3 is a quantization of the pair $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*, r)$ or that the pair $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*, r)$ is the classical limit of the quasitriangular Hopf algebra over $\mathbb{K}_{t,u}[[\hbar]]$.

Fix an ordered basis $\{e_a\}$ in $\mathfrak{a}_{t,u}$, and its dual basis $\{e^a\}$ in $\mathfrak{a}_{r_{t,u}}^*$. Then we may construct ordered bases in $\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*$, $\mathcal{U}\mathfrak{a}_{t,u}$, $\mathcal{U}\mathfrak{a}_{r_{t,u}}^*$ and $\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)^{\otimes 2}$.

Lemma 2.5. The element $J_{r_{t,u}} \in \mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)^{\otimes 2}[[\hbar]]$ considered in theorem 2.3, 3) has the form

$$J_{r_{t,u}} = 1 \otimes 1 + \frac{1}{2}r\hbar + \sum_{k \geq 2} \left(r_{t,u}^{i_1 j_1} \dots r_{t,u}^{i_{l(k)} j_{l(k)}} Q_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k} \right) \hbar^k,$$

where $Q_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k} \in \mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)^{\otimes 2}$ are linear combinations of elements in the ordered basis fixed above. The coefficients of these linear combinations are \mathbb{K} -linear combinations of elements determined by the structure constants of the Lie algebra $(\mathfrak{a}_{t,u}, [\cdot, \cdot]_{\mathfrak{a}_{t,u}})$. The element $r_{t,u}$ is present in every element of the ordered basis through the product in $\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)$, but it does not occur in the coefficients defining $Q_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k}$.

3. Quantization of the nondegenerate triangular Lie bialgebra $(\mathfrak{a}_{t,u}, [\cdot, \cdot]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$

As in [8, 7], we define the mapping $\chi_{r_{t,u}} : \mathfrak{a}_{r_{t,u}}^* \longrightarrow \mathfrak{a}_{t,u}$ by $\chi_{r_{t,u}}(\xi) = (\xi \otimes 1)r_{t,u}$.

Proposition 3.1. *The mapping $\tilde{\pi}_{t,u} : \mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^* \longrightarrow \mathfrak{a}_{t,u}$, defined by $\tilde{\pi}_{t,u}(x; \xi) = x + \chi_{r_{t,u}}(\xi)$, is a Lie-bialgebra-morphism. That is, a Lie-algebra morphism verifying $d_c(t)r_{t,u} \circ \tilde{\pi}_{t,u} = (\tilde{\pi}_{t,u} \otimes \tilde{\pi}_{t,u}) \circ d_c(t, u)r$. Moreover $(\tilde{\pi}_{t,u} \otimes \tilde{\pi}_{t,u})r = r_{t,u}$. The symbol $\tilde{\pi}_{t,u}$ will also denote the unique algebra morphism $\tilde{\pi}_{t,u} : \mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*) \longrightarrow \mathcal{U}\mathfrak{a}_t$ defined by the Lie algebra morphism $\tilde{\pi}_{t,u}$.*

Theorem 3.2. *Consider the quasitriangular double Lie bialgebra $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*, [;]_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*}, \varepsilon_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*} = d_c(t, u)r)$ over $\mathbb{K}_{t,u}$. Let $(\mathcal{U}\mathfrak{a}_{t,u}, \Delta_{\mathfrak{a}_{t,u}})$ be the usual Hopf universal enveloping algebra. Let*

$$\left(\mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)[[\hbar]], \Delta_0^{r_{t,u}}, \Phi_{r_{t,u}}, R_0^{r_{t,u}} = e^{\frac{\hbar}{2}\Omega} \right)$$

be the quasitriangular quasi-Hopf algebra, considered in Theorem 2.1, whose classical limit is the pair $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^, \Omega)$. Then we have $(\tilde{\pi}_{t,u} \otimes \tilde{\pi}_{t,u}) \circ \Delta_0^{r_{t,u}} = \Delta_{\mathfrak{a}_{t,u}} \circ \tilde{\pi}_{t,u}$. Defining $\tilde{\Phi}_{r_{t,u}} = (\tilde{\pi}_{t,u} \otimes \tilde{\pi}_{t,u} \otimes \tilde{\pi}_{t,u})\Phi_{r_{t,u}}$ and $R_{\mathfrak{a}_{t,u}} = (\tilde{\pi}_{t,u} \otimes \tilde{\pi}_{t,u})R_0^{r_{t,u}}$, we get $\tilde{\Phi}_{r_{t,u}} = 1 \otimes 1$ and $R_{\mathfrak{a}_{t,u}} = 1 \otimes 1$. In this way, we obtain the (trivial) triangular Hopf algebra $(\mathcal{U}\mathfrak{a}_{t,u}[[\hbar]], \Delta_{\mathfrak{a}_{t,u}}, \tilde{\Phi}_{r_{t,u}} = 1 \otimes 1 \otimes 1, R_{\mathfrak{a}_{t,u}} = 1 \otimes 1)$ over the ring $\mathbb{K}_{t,u}[[\hbar]]$. We call this algebra a quantization of the pair $(\mathfrak{a}_{t,u}, 0)$, see [5].*

From Proposition 3.1, Theorem 3.2 and 3) of Theorem 2.3 we obtain

Corollary 3.3. *Write $\tilde{J}_{r_{t,u}} = (\tilde{\pi}_{t,u} \otimes \tilde{\pi}_{t,u})J_{r_{t,u}} \in (\mathcal{U}\mathfrak{a}_{t,u} \otimes \mathcal{U}(\mathfrak{a}_{t,u})[[\hbar]])$. Then*

- 1) $\tilde{J}_{r_{t,u}} = 1 \otimes 1 + \frac{1}{2}r_{t,u}\hbar + \dots$
- 2) $(\Delta_{\mathfrak{a}_{t,u}} \otimes 1)\tilde{J}_{r_{t,u}} \cdot (\tilde{J}_{r_{t,u}} \otimes 1) = (1 \otimes \Delta_{\mathfrak{a}_{t,u}})\tilde{J}_{r_{t,u}} \cdot (1 \otimes \tilde{J}_{r_{t,u}})$.
- 3) $\tilde{R}_{\mathfrak{a}_{t,u}} = (\tilde{\pi}_{t,u} \otimes \tilde{\pi}_{t,u})R^{r_{t,u}} = \sigma(\tilde{J}_{r_{t,u}}^{-1}) \cdot (1 \otimes 1) \cdot \tilde{J}_{r_{t,u}} = 1 \otimes 1 + r_{t,u}\hbar + \dots$

The products in these expressions coincide with the products of the enveloping algebra $\mathcal{U}\mathfrak{a}_{t,u}[[\hbar]] \equiv \mathcal{U}\mathfrak{a}_t[[u, \hbar]]$. The set $(\mathcal{U}\mathfrak{a}_{t,u}[[\hbar]], \tilde{\Delta}_{\mathfrak{a}_{t,u}}, \tilde{R}_{\mathfrak{a}_{t,u}})$, denoted by $A_{\mathfrak{a}_{t,u}[[\hbar]], \tilde{J}_{r_{t,u}}^{-1}}$, is a triangular Hopf algebra over $\mathbb{K}_{t,u}[[\hbar]]$. This algebra can be obtained by a twist via the element $\tilde{J}_{r_{t,u}}^{-1}$ from the trivial triangular Hopf algebra $(\mathcal{U}\mathfrak{a}_{t,u}\Delta_{\mathfrak{a}_{t,u}}, R_{\mathfrak{a}_{t,u}} = 1 \otimes 1)$ considered in Theorem 3.2. It is a quantization of the pair $(\mathfrak{a}_{t,u}; r_{t,u})$.

From Lemma 2.5 we obtain

Lemma 3.4. *The element $\tilde{J}_{r_{t,u}}$ has the form*

$$\tilde{J}_{r_{t,u}} = 1 \otimes 1 + \frac{1}{2}r_{t,u}\hbar + \sum_{k \geq 2} \left(r_{t,u}^{i_1 j_1} \dots r_{t,u}^{i_{l(k)} j_{l(k)}} M_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k} \right) \hbar^k,$$

where $M_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k}$ is a linear combination of elements in the ordered basis chosen in $\mathcal{U}\mathfrak{a}_{t,u}$ and whose coefficients are \mathbb{K} -linear combinations of elements (polynomials) determined by the structure constants of the Lie algebra $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}})$. The element $r_{t,u}$ does not appear in $M_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k} \in (\mathcal{U}\mathfrak{a}_{t,u})^{\otimes 2}$.

We now define Invariant Star Products.

Definition 3.5. [1, 4, 12] An Invariant Star Product on a nondegenerate triangular Lie bialgebra $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$ over the ring $\mathbb{K}_{t,u}$ is any element $F(t, u) = \sum_0^\infty F_k(t, u) \cdot \hbar^k \in (\mathcal{U}\mathfrak{a}_{t,u} \otimes \mathcal{U}\mathfrak{a}_{t,u})[[\hbar]]$ verifying the following equalities:

- 1) $F(t, u) = 1 \otimes 1 \text{ mod } \hbar$;
- 2) $F(t, u) - \sigma(F(t, u)) = r_{t,u} \hbar \text{ mod } \hbar^2$;
- 3) $(\Delta_{\mathfrak{a}_{t,u}} \otimes 1)F(t, u) \cdot (F(t, u) \otimes 1) = (1 \otimes \Delta_{\mathfrak{a}_{t,u}})F(t, u) \cdot (1 \otimes F(t, u))$.

The products in 3) coincide with the products of the enveloping algebra $\mathcal{U}\mathfrak{a}_{t,u}[[\hbar]] \equiv \mathcal{U}\mathfrak{a}_t[[u, \hbar]]$, that is, they coincide with the $\mathbb{K}[[u, \hbar]]$ linear extension of the product of the enveloping algebra $\mathcal{U}\mathfrak{a}_t$.

Then we have

Proposition 3.6. The element $\tilde{J}_{r_{t,u}} \in (\mathcal{U}\mathfrak{a}_{t,u})^{\otimes 2}[[\hbar]]$, considered in Corollary 3.3, is an Invariant Star Product on the nondegenerate triangular Lie bialgebra $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$ over the ring $\mathbb{K}_{t,u}$.

Definition 3.7. [1, 4, 12] An Invariant Star Product on a nondegenerate triangular Lie bialgebra $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_t)$ over the ring \mathbb{K}_t is any element $F(t) = \sum_0^\infty F_k(t) \cdot \hbar^k \in (\mathcal{U}\mathfrak{a}_t \otimes \mathcal{U}\mathfrak{a}_t)[[\hbar]]$ verifying the following equalities:

- 1) $F(t) = 1 \otimes 1 \text{ mod } \hbar$;
- 2) $F(t) - \sigma(F(t)) = r_t \hbar \text{ mod } \hbar^2$;
- 3) $(\Delta_{\mathfrak{a}_t} \otimes 1)F(t) \cdot (F(t) \otimes 1) = (1 \otimes \Delta_{\mathfrak{a}_t})F(t) \cdot (1 \otimes F(t))$.

The products in 3) coincide with the products of the enveloping algebra $\mathcal{U}(\mathfrak{a}_t \otimes_{\mathbb{K}_t} \mathbb{K}[[t, \hbar]])$, that is, they coincide with the $\mathbb{K}[[\hbar]]$ linear extension of the product of the enveloping algebra $\mathcal{U}\mathfrak{a}_t$.

Proposition 3.8. Let $F(t, u) \in (\mathcal{U}\mathfrak{a}_{t,u} \otimes \mathcal{U}\mathfrak{a}_{t,u})[[\hbar]]$ be an Invariant Star Product on the nondegenerate triangular Lie bialgebra $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$ over the ring $\mathbb{K}_{t,u}$. Consider the element $F(t) \in (\mathcal{U}\mathfrak{a}_t \otimes \mathcal{U}\mathfrak{a}_t)[[\hbar]]$ obtained from $F(t, u)$ by setting $u = \hbar$ in all the elements defining $F(t, u)$;

in particular by setting $r_{t,\hbar} = \sum_{l \geq 1} r_{t,l} \hbar^l \in \mathfrak{a}_t[[\hbar]] \wedge \mathfrak{a}_t[[\hbar]]$. Then the element $F(t) \in (\mathcal{U}\mathfrak{a}_t \otimes \mathcal{U}\mathfrak{a}_t)[[\hbar]]$ is an Invariant Star product on the triangular nondegenerate Lie bialgebra $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_t)$ over the ring \mathbb{K}_t .

Obviously we have

Corollary 3.9. *Let $\tilde{J}_{r_{t,\hbar}} \in (\mathcal{U}\mathfrak{a}_t \otimes \mathcal{U}\mathfrak{a}_t)[[\hbar]]$ be the element as in Proposition 3.8 obtained from the element $\tilde{J}_{r_{t,u}} \in (\mathcal{U}\mathfrak{a}_{t,u} \otimes \mathcal{U}\mathfrak{a}_{t,u})[[\hbar]]$ considered in Corollary 3.3. Then $\tilde{J}_{r_{t,\hbar}}$ is an Invariant Star Product on the triangular nondegenerate Lie bialgebra $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_t)$ over the ring \mathbb{K}_t .*

4. An Invariant Star Product $F(t) \in (\mathcal{U}\mathfrak{a}_t \otimes \mathcal{U}\mathfrak{a}_t)[[\hbar]]$ on $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_t)$ determines an element $r_{t,\hbar} \in (\mathfrak{a}_t \wedge \mathfrak{a}_t)[[\hbar]]$ such that $F(t)$ and $\tilde{J}_{r_{t,\hbar}} \in (\mathcal{U}\mathfrak{a}_t \otimes \mathcal{U}\mathfrak{a}_t)[[\hbar]]$ are equivalent

Let $F(t) \in \mathcal{U}\mathfrak{a}_t[[\hbar]] \hat{\otimes} \mathcal{U}\mathfrak{a}_t[[\hbar]]$ be an Invariant Star Product on the nondegenerate triangular Lie bialgebra $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_t)$ over \mathbb{K}_t . Let $A_{\mathfrak{a}_t[[\hbar]], F^{-1}(t)}$ be the triangular Hopf QUE algebra obtained by a twist via $F^{-1}(t)$ from the trivial triangular Hopf QUE algebra $(\mathcal{U}\mathfrak{a}_t[[\hbar]], \Delta_{\mathfrak{a}_t}, R_{\mathfrak{a}_t} = 1 \otimes 1)$. Then, this algebra is a quantization of the pair (\mathfrak{a}_t, r_t) .

The following proposition does not depend on any specific context of quantization but only on: i) the notion of deformation of associative algebras; ii) the fact that the Hochschild cohomology of the bialgebra $\mathcal{U}\mathfrak{a}_t$ is $H^k(\mathcal{U}\mathfrak{a}_t) = \Lambda^k \mathfrak{a}_t$, $k \in \mathbb{N}$, see [2]; iii) the Hochschild cohomological interpretation of Quantum Yang Baxter equation [12, 13].

Proposition 4.1. [12] *Let $F(t) = \sum_i^\infty F_i(t) \hbar^i$ and $F'(t) = \sum_i^\infty F'_i(t) \hbar^i$ be Invariant Star Products on $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_t)$. Let $A_{\mathfrak{a}_t[[\hbar]], F^{-1}(t)}$ and $A_{\mathfrak{a}_t[[\hbar]], F'^{-1}(t)}$ be as above in this Section. Suppose that $F(t)$ and $F'(t)$ coincide up to order k , i.e. $F_l(t) = F'_l(t)$, $l = 1, 2, \dots, k$. Then: a) there exist $h_{k+1} \in \mathfrak{a}_t \wedge \mathfrak{a}_t$ and $E_{k+1}(t) \in \mathcal{U}\mathfrak{a}_t$ such that $F'_{k+1}(t) - F_{k+1}(t) = h_{k+1} + d_H E_{k+1}(t)$ where d_H is the coboundary operator in the Hochschild cohomology of $\mathcal{U}\mathfrak{a}_t$; b) h_{k+1} is not only a Hochschild 2-cocycle but also a Poisson 2-cocycle relatively to the invariant Poisson structure defined by the element $r_t \in \mathfrak{a}_t \wedge \mathfrak{a}_t$.*

Again, the above Hochschild cohomology spaces and proposition 4.1 play a central role in the proof of next theorem. In the context of quantification in

[7] this theorem corresponds to a main theorem by Drinfeld in the context of quantification in [4]. In [12, 13] there is a proof of this Drinfeld theorem. See the References in [13] for a similar theorem about Star Products on general symplectic manifolds and on Poisson manifolds.

Theorem 4.2. [14, 15] *Fix a Lie associator Φ . Let $A_{\mathfrak{a}_t[[\hbar]], F^{-1}(t)}$ be defined at the beginning of this section. We have: (a) There exist elements $r_{t,\hbar} = r_t + r_{t,2}\hbar + r_{t,3}\hbar^2 + \dots \in (\wedge^2 \mathfrak{a}_t)[[\hbar]]$ and $E^{r_{t,\hbar}} = 1 + E_1^{r_{t,\hbar}}\hbar + \dots + E_n^{r_{t,\hbar}}\hbar^n + \dots \in \mathcal{U}\mathfrak{a}_t[[\hbar]]$ such that*

$$F(t) = \Delta_{\mathfrak{a}_t}((E^{r_{t,\hbar}})^{-1}) \cdot_t \tilde{J}_{r_{t,\hbar}}^\Phi \cdot_t (E^{r_{t,\hbar}} \otimes E^{r_{t,\hbar}});$$

i.e., $F(t)$ and $\tilde{J}_{r_{t,\hbar}}^\Phi \in (\mathcal{U}\mathfrak{a}_t \otimes \mathcal{U}\mathfrak{a}_t)[[\hbar]]$ are equivalent Invariant Star Products over the nondegenerate triangular Lie bialgebra $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_t)$ over the ring \mathbb{K}_t .

(b) The triangular Hopf QUE algebras $A_{\mathfrak{a}_t[[\hbar]], F^{-1}(t)}$ and $A_{\mathfrak{a}_t[[\hbar]], (\tilde{J}_{r_{t,\hbar}}^\Phi)^{-1}}$ are isomorphic.

As a consequence we have the following isomorphisms:

Corollary 4.3. *Let Φ, Φ' be two Lie associators. Let $A_{\mathfrak{a}_t[[\hbar]], F^{-1}(t)}$ be given as in the theorem. Let $r_{t,\hbar}, r'_{t,\hbar} \in (\wedge^2 \mathfrak{a}_t)[[\hbar]]$ be the elements determined in the theorem by the pairs $(\Phi; A_{\mathfrak{a}_t[[\hbar]], F^{-1}(t)})$ and $(\Phi'; A_{\mathfrak{a}_t[[\hbar]], F^{-1}(t)})$, respectively. Then we have*

$$A_{\mathfrak{a}_t[[\hbar]], F^{-1}(t)} \stackrel{isom}{\approx} A_{\mathfrak{a}_t[[\hbar]], (\tilde{J}_{r_{t,\hbar}}^\Phi)^{-1}} \stackrel{isom}{\approx} A_{\mathfrak{a}_t[[\hbar]], (\tilde{J}_{r'_{t,\hbar}}^{\Phi'})^{-1}}$$

5. Some properties of nondegenerate triangular Lie bialgebras $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$ over $\mathbb{K}_t[[u]]$

1) We now develop what we wrote in Section 1, 8) in the Introduction. We need the following:

Proposition 5.1. *Let $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$ be a nondegenerate triangular Lie bialgebra over $\mathbb{K}_{t,u}$. Let $\varphi_{t,u}^1 : \mathfrak{a}_{t,u} \longrightarrow \mathfrak{a}_{t,u}$ be a Lie algebra isomorphism. Let $r'_{t,u}$ be the element in $\mathfrak{a}_t[[u]] \wedge \mathfrak{a}_t[[u]]$ defined as $r'_{t,u} = (\varphi_{t,u}^1 \otimes \varphi_{t,u}^1)r_{t,u}$.*

a) The set $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon'_{\mathfrak{a}_{t,u}} = d_c(t)r'_{t,u})$ is a nondegenerate triangular Lie bialgebra.

b) The transposed map $(\varphi_{t,u}^1)^t : \mathfrak{a}_{r'_{t,u}}^ \longrightarrow \mathfrak{a}_{r_{t,u}}^*$ is a Lie algebra isomorphism.*

c) The pair $(\varphi_{t,u}^1; \varphi_{t,u}^2 = ((\varphi_{t,u}^1)^t)^{-1})$ defines a Lie bialgebra isomorphism between the Lie bialgebra $(\mathfrak{a}_t \oplus \mathfrak{a}_{r_{t,u}}^*, [,]_{\mathfrak{a}_t \oplus \mathfrak{a}_{r_{t,u}}^*}, \varepsilon_{\mathfrak{a}_t \oplus \mathfrak{a}_{r_{t,u}}^*} = d_c(t, u)r)$ and the Lie bialgebra $(\mathfrak{a}_t \oplus \mathfrak{a}_{r'_{t,u}}^*, [,]_{\mathfrak{a}_t \oplus \mathfrak{a}_{r'_{t,u}}^*}, \varepsilon_{\mathfrak{a}_t \oplus \mathfrak{a}_{r'_{t,u}}^*} = d_c(t, u)r)$. Furthermore, this isomorphism sends the canonical element $r \in (\mathfrak{a} \oplus \mathfrak{a}^*)^{\otimes 2}$ into itself.

Corollary 5.2. a) Under the hypotheses of the proposition let $\beta_{t,u} = \mu_{r_{t,u}}(r_{t,u})$ and $\beta'_{r,u} = \mu_{r'_{t,u}}(r'_{t,u})$ be elements of $\mathfrak{a}_t^* \wedge \mathfrak{a}_t^*[[u]]$. Then $(\varphi_{t,u}^2 \otimes \varphi_{t,u}^2)\beta_{t,u} = \beta'_{t,u}$.

b) Conversely, let $\beta_{t,u}$ and $\beta'_{t,u}$ as considered in a). Let $\varphi_{t,u}^1 : \mathfrak{a}_{t,u} \longrightarrow \mathfrak{a}_{t,u}$ be a Lie algebra isomorphism and $\varphi_{t,u}^2 = ((\varphi_{t,u}^1)^t)^{-1}$. Suppose that $(\varphi_{t,u}^2 \otimes \varphi_{t,u}^2)\beta_{t,u} = \beta'_{t,u}$. Then, $(\varphi_{t,u}^1 \otimes \varphi_{t,u}^1)r_{t,u} = r'_{t,u}$.

2) In the Lie algebra $(\mathfrak{a}_{t,u}, [,]_{\mathfrak{a}_{t,u}})$ over $\mathbb{K}_{t,u}$ consider the following Lie-algebra isomorphisms: $\varphi_{t,u}^1 = \exp(t \cdot \text{ad}_{X_{t,u}})$ where $X_{t,u} = X_{1,u} + X_{2,u}t + X_{3,u}t^2 + \dots \in \mathfrak{a}_u[[t]]$. Then $\varphi_{t,u}^2 = \exp(-t \cdot \text{ad}_{X_{t,u}}^t) = \exp(t \cdot \text{ad}_{X_{t,u}}^*)$. Our interest is in the map $\varphi_{t,u}^2 \otimes \varphi_{t,u}^2 = \exp(\text{ad}_{tX_{t,u}}^* \otimes 1 + 1 \otimes \text{ad}_{tX_{t,u}}^*)$.

Proposition 5.3. Let $\beta_{t,1} = \beta_t$ and let $\beta_{t,u} = \beta_{t,1} + \beta_{t,2}u + \beta_{t,3}u^2 + \dots \in \wedge^2(\mathfrak{a}_t^*[[u]])$, or equivalently $\beta_{t,u} = \beta_{1,u} + \beta_{2,u}t + \beta_{3,u}t^2 + \dots \in \wedge^2(\mathfrak{a}_u^*[[t]])$, be a nondegenerate 2-cocycle on the Lie algebra $(\mathfrak{a}_{t,u}, [,]_{\mathfrak{a}_{t,u}})$. The elements $\beta_{t,1}, \beta_{t,2}, \dots \in \wedge^2(\mathfrak{a}_t^*)$ are then 2-cocycles on the Lie algebra $(\mathfrak{a}_t, [,]_{\mathfrak{a}_t})$, with the trivial action, and $\beta_{t,1}$ is nondegenerate. Let $X_{t,u}$ be as considered before. Then

$$\exp(\text{ad}_{tX_{t,u}}^*)^{\otimes 2}(\beta_{t,u}) = \exp(\text{ad}_{tX_{t,u}}^* \otimes 1 + 1 \otimes \text{ad}_{tX_{t,u}}^*)(\beta_{t,u}) = \beta_{t,u} + d_R(t)\gamma_{t,u},$$

where $\gamma_{t,u} = t \sum_{l \geq 1} \gamma_{l,u}(t)t^{l-1} \in t\mathfrak{a}_u^*[[t]]$ and

$$\gamma_{1,u}(t) = (i(X_{t,u})\beta_{t,u})$$

$$\gamma_{2,u}(t) = \left(\frac{1}{2!} (i(X_{t,u})\beta_{t,u}) \circ \text{ad}_{X_{t,u}} \right)$$

$$\gamma_{3,u}(t) = ((i(X_{t,u})\beta_{t,u}) \circ \text{ad}_{X_{t,u}} \circ \text{ad}_{X_{t,u}}), \text{ etc}$$

A converse of proposition 5.3 is:

Proposition 5.4. Let $\beta_{t,u}$ be as considered in proposition 5.3. Let $\gamma_{t,u} = \alpha_{1,u}t + \alpha_{2,u}t^2 + \alpha_{3,u}t^3 + \dots \in \mathfrak{a}_u^*[[t]]$; $\alpha_{l,u} \in \mathfrak{a}_u^*$, $l = 1, 2, \dots$. Define $\beta'_{t,u} =$

$\beta_{t,u} + d_R(t)\gamma_{t,u}$. Then, there exists a unique $X_{t,u} = X_{1,u} + X_{2,u}t + X_{3,u}t^2 + \dots \in \mathfrak{a}_u[[t]]$ such that $\exp(\text{ad}_{tX_{t,u}}^*)^{\otimes 2}(\beta_{t,u}) = \beta'_{t,u}$. It is given by

$$\begin{aligned} i(X_{1,u})\beta_{1,u} &= \alpha_{1,u}, \\ i(X_{2,u})\beta_{1,u} + i(X_{1,u})\beta_{2,u} + \frac{1}{2!}(i(X_{1,u})\beta_1) \circ (i(X_{1,u})B_{1,u}) &= \alpha_{2,u}, \\ i(X_{1,u})\beta_{3,u} + i(X_{2,u})\beta_{2,u} + i(X_{3,u})\beta_{1,u} + \frac{1}{2!} &\left((i(X_{2,u})\beta_{1,u}) \circ (i(X_{1,u})B_{1,u}) \right. \\ + (i(X_{1,u})\beta_{2,u}) \circ (i(X_{1,u})B_{1,u}) + (i(X_{1,u})\beta_{1,u}) \circ (i(X_{2,u})B_{1,u}) &+ (i(X_{1,u})\beta_{1,u}) \circ (i(X_{1,u})B_{2,u}) \\ \left. + (i(X_{1,u})\beta_{1,u}) \circ (i(X_{1,u})B_{1,u}) \circ (i(X_{1,u})B_{1,u}) \right) &= \alpha_{3,u}, \quad \text{etc.} \end{aligned}$$

where $B_{l,u} \in L(\mathfrak{a}_u, \mathfrak{a}_u; \mathfrak{a}_u)$, $l = 1, 2, \dots$ are some well determined bilinear mappings.

As $\beta_{1,u}$ is invertible the first equation allows us to compute $X_{1,u}$. Analogously the second equation allows us to obtain $X_{2,u}$ etc. It is easy to obtain a general form for $X_{l,u}$ as a function of $X_{k,u}$, $1 \leq k < l$.

3) The following property is needed:

Proposition 5.5. *Let $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t)r_{t,u})$ and $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon'_{\mathfrak{a}_{t,u}} = d_c(t)r'_{t,u})$ be nondegenerate triangular Lie bialgebras as considered in Section 2.*

Let $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}, [;]_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}}, \varepsilon_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}} = d_c(t, u)r)$ and $(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r'_{t,u}}, [;]_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r'_{t,u}}}, \varepsilon_{\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r'_{t,u}}} = d_c(t, u)r)$ be the corresponding quasitriangular doubles. Let $(\varphi_{t,u}^1; \psi_{t,u}) : \mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^ \longrightarrow \mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r'_{t,u}}^*$ be a Lie algebra isomorphism such that $\varphi_{t,u}^1 : \mathfrak{a}_{t,u} \longrightarrow \mathfrak{a}_{t,u}$ and $\psi_{t,u} : \mathfrak{a}_{r_{t,u}}^* \longrightarrow \mathfrak{a}_{r'_{t,u}}^*$ are Lie algebra isomorphisms. Let $\tilde{\varphi}_{t,u}^1, \tilde{\psi}_{t,u}$ be the extensions of $\varphi_{t,u}^1$ and $\psi_{t,u}$ to homomorphisms $\mathcal{U}\mathfrak{a}_{t,u} \longrightarrow \mathcal{U}\mathfrak{a}_{t,u}$ and $\mathcal{U}\mathfrak{a}_{r_{t,u}}^* \longrightarrow \mathcal{U}\mathfrak{a}_{r'_{t,u}}^*$. Let $X(t, u) \in \mathcal{U}(\mathfrak{a}_{t,u} \oplus \mathfrak{a}_{r_{t,u}}^*)^{\otimes 2}$. Let $\phi_{r_{t,u}}$ and $\phi_{r'_{t,u}}$ be the Lie algebra-module isomorphisms defined in Theorem 2.3, 2). Then we have*

$$\begin{aligned} \phi_{r'_{t,u}}^{-1} \left[\left((\tilde{\varphi}_{t,u}^1; \tilde{\psi}_{t,u})^{\otimes 2}(X(t, u)) \right) \cdot (1_+^{r'_{t,u}} \otimes 1_-^{r'_{t,u}}) \right] &= \\ = \left((\tilde{\varphi}_{t,u}^1; \tilde{\psi}_{t,u}) \circ \phi_{r_{t,u}}^{-1} \right) (X(t, u) \cdot (1_+^{r_{t,u}} \otimes 1_-^{r_{t,u}})). & \end{aligned}$$

We can also prove the following:

Proposition 5.6. *Let $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon_{\mathfrak{a}_{t,u}} = d_c(t, u)r_{t,u})$ and $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}}, \varepsilon'_{\mathfrak{a}_{t,u}} = d_c(t, u)r'_{t,u})$ be nondegenerate triangular Lie bialgebras over $\mathbb{K}_{t,u}$. Let $\varphi_{t,u}^1 : \mathfrak{a}_{t,u} \rightarrow \mathfrak{a}_{t,u}$ be a Lie algebra isomorphism such that $r'_{t,u} = (\varphi_{t,u}^1 \otimes \varphi_{t,u}^1)r_{t,u}$ and let $(\varphi_{t,u}^1; \varphi_{t,u}^2)$ be the Lie bialgebra isomorphism between the corresponding classical doubles constructed in proposition 5.1. Then, we have*

$$\tilde{\pi}'_{t,u} \circ (\varphi_{t,u}^1; \varphi_{t,u}^2) = \varphi_{t,u}^1 \circ \tilde{\pi}_{t,u},$$

where $\tilde{\pi}'_{t,u}$ and $\tilde{\pi}_{t,u}$ are defined in Proposition 3.1.

6. A necessary and sufficient condition to be isomorphic two triangular Hopf algebras $A_{\mathfrak{a}_t[[\hbar]], \tilde{J}_{r_{t,\hbar}}^{-1}}$ and $A_{\mathfrak{a}_t[[\hbar]], \tilde{J}'_{r'_{t,\hbar}}^{-1}}$ over $\mathbb{K}_t[[\hbar]]$

If on the expression of $J_{r_{t,u}}$ given in theorem 2.3 we take into account Proposition 5.5 and also the form of a Lie associator $\Phi = e^{P(\hbar\Omega_{12}, \hbar\Omega_{23})}$ we arrive to:

Proposition 6.1. *Hypotheses are as in Proposition 5.5. Let $J_{r'_{t,u}}$ and $J_{r_{t,u}}$ be the corresponding elements in Theorem 2.3, 3). Suppose moreover that $(\varphi_{t,u}^1; \psi_{t,u}) \otimes (\varphi_{t,u}^1; \psi_{t,u}) \Omega = \Omega$. Then $J_{r'_{t,u}} = (\tilde{\varphi}_{t,u}^1; \tilde{\psi}_{t,u})^{\otimes 2} J_{r_{t,u}}$. In particular this proposition is valid for the Lie bialgebra isomorphism $(\varphi_{t,u}^1; \varphi_{t,u}^2)$ considered in propositions 5.3 and 5.4.*

4) Using Propositions 5.1, 6.1, 5.6 and Corollary 5.2 we can prove:

Proposition 6.2. *a) Let $\tilde{J}_{r_{t,\hbar}}$ and $\tilde{J}'_{r'_{t,\hbar}}$ be elements $\in (\mathcal{U}\mathfrak{a}_t)^{\otimes 2}[[\hbar]]$ which are Invariant Star Products on a nondegenerate triangular Lie bialgebra $(\mathfrak{a}_t, [;]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_t)$ over \mathbb{K}_t as in Definition 3.5, and obtained as in Corollary 3.3 and Corollary 3.9 respectively from the nondegenerate solutions $r_{t,u}, r'_{t,u} \in (\mathfrak{a}_t \wedge \mathfrak{a}_t)[[u]]$ of the YBE on the Lie-algebra $(\mathfrak{a}_{t,u}, [;]_{\mathfrak{a}_{t,u}})$ as in Section 1. Let $\mu_{r_{t,u}}(r_{t,u}) = \beta_{t,u} = \beta_{t,1} + \beta_{t,2}u + \beta_{t,3}u^2 \cdots \in (\mathfrak{a}_t^* \wedge \mathfrak{a}_t^*)[[u]]$ and $\mu_{r'_{t,u}}(r'_{t,u}) = \beta'_{t,u} = \beta_{t,1} + \beta'_{t,2}u + \beta'_{t,3}u^2 \cdots \in (\mathfrak{a}_t^* \wedge \mathfrak{a}_t^*)[[u]]$. b) Suppose that the cocycles $\beta_{t,u}$ and $\beta'_{t,u}$ belong to the same cohomological class in $H(\mathfrak{a}_{t,u}) \equiv H^2(\mathfrak{a}_t)[[u]]$, i.e., $\beta'_{t,u} = \beta_{t,u} + d_R(t)\gamma_{t,u}$ for some 1-cochain $\gamma_{t,u} = \gamma_{t,1}u + \gamma_{t,2}u^2 + \gamma_{t,3}u^3 \cdots \in \mathfrak{a}_t^*[[u]]$. Then, $\tilde{J}_{r_{t,\hbar}}$ and $\tilde{J}'_{r'_{t,\hbar}}$ are equivalent Invariant Star Products.*

To prove the converse we need the following lemma:

Lemma 6.3. *Suppose that in Proposition 6.2*

$$\beta_{t,u} = \beta_{t,1} + \beta_{t,2}u + \beta_{t,3}u^2 + \cdots + \beta_{t,R-1}u^{R-2} + \beta_{t,R}u^{R-1} + \cdots$$

$$\beta'_{t,u} = \beta_{t,1} + \beta_{t,2}u + \beta_{t,3}u^2 + \cdots + \beta_{t,R-1}u^{R-2} + (\beta_{t,R} + d_R\alpha_{t,(R-1)})u^{R-1} + \cdots$$

where $\alpha_{t,(R-1)}$ is an element in \mathfrak{a}_t^* , that is, a 1-cochain on the Lie algebra $(\mathfrak{a}_t, [\cdot, \cdot]_{\mathfrak{a}_t})$ over the ring \mathbb{K}_t . This means that $\beta'_{t,u}$ and $\beta_{t,u}$ are equal except in the term of order $R-1$. Then, $\tilde{J}_{r_t, \hbar}$ and $\tilde{J}_{r'_t, \hbar}$ are equivalent,

$$\tilde{J}_{r'_t, \hbar} = \Delta_{\mathfrak{a}_t}(E)^{-1} \cdot_t \tilde{J}_{r_t, \hbar} \cdot_t (E \otimes E),$$

and the element $E = 1 + E_{t,1}\hbar + E_{t,2}\hbar^2 + \cdots + E_{t,(R-1)}\hbar^{R-1} + \cdots \in \mathcal{U}\mathfrak{a}_t[[\hbar]]$ which defines this equivalence verifies

$$E_{t,1} = 0, \quad E_{t,2}, \dots, \quad E_{t,(R-2)} = 0, \quad E_{t,(R-1)} = \chi_{r_t}(\alpha_{t,(R-1)}) = \mu_{r_t}^{-1}(\alpha_{t,(R-1)}).$$

Lemma 6.3 and Hochschild cohomology properties allow us to prove

Proposition 6.4. *Let $\tilde{J}_{r_t, \hbar}$ and $\tilde{J}_{r'_t, \hbar}$ as considered in Theorem 6.2. Suppose that $\tilde{J}_{r_t, \hbar}$ and $\tilde{J}_{r'_t, \hbar}$ are equivalent. Then $\beta'_{t,u}$ and $\beta_{t,u}$ belong to the same cohomological class, i.e., $\beta'_{t,u} = \beta_{t,u} + d_R(t)\gamma_{t,u}$ for some 1-cochain $\gamma_{t,u} \cdots \in \mathfrak{a}_t^*[[u]]$.*

Combining the last two theorems we obtain the following result, similar in Etingof-Kazhdan quantization theory to the one by Drinfeld in [4]:

Proposition 6.5. *Let $\tilde{J}_{r_t, \hbar}$ and $\tilde{J}_{r'_t, \hbar}$ be elements in $(\mathcal{U}\mathfrak{a}_t)^{\otimes 2}[[\hbar]]$ which are Invariant Star Products on a nondegenerate triangular Lie bialgebra $(\mathfrak{a}_t, [\cdot, \cdot]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_t)$ over \mathbb{K}_t as in Theorem 6.2. Then $\tilde{J}_{r_t, \hbar}$ and $\tilde{J}_{r'_t, \hbar}$ are equivalent Invariant Star Products if, and only if, $\mu_{r_t, u}(r_{t,u}) = \beta_{t,u}$ and $\mu_{r'_t, u}(r'_{t,u}) = \beta'_{t,u}$ belong to the same cohomological class in $H^2(\mathfrak{a}_t)[[u]]$. In other words, $\tilde{J}_{r_t, \hbar}$ and $\tilde{J}_{r'_t, \hbar}$ are equivalent Invariant Star Products if, and only if, there exists a 1-cochain $\gamma_{t,u} \in \mathfrak{a}_{t,u}^*$ such that $\beta'_{t,u} = \beta_{t,u} + d_R(t)\gamma_{t,u}$.*

Theorem 6.5 and Remark 2) in page 841 of [6] allow us to obtain

Proposition 6.6. *Two triangular Hopf algebras $A_{\mathfrak{a}_t[[\hbar]], \tilde{J}_{r_t, \hbar}^{-1}}$ and $A_{\mathfrak{a}_t[[\hbar]], \tilde{J}_{r'_t, \hbar}^{-1}}$ over $\mathbb{K}_t[[\hbar]]$ defined as in Corollary 3.9 and Section 4 are isomorphic if, and only if, there exists an isomorphism of Lie algebras $\lambda_t : \mathfrak{a}_t[[\hbar]] \longrightarrow \mathfrak{a}_t[[\hbar]]$ over $\mathbb{K}_t[[\hbar]]$ such that $(\lambda_t^2 \otimes \lambda_t^2)\beta_{t, \hbar}$ and $\beta'_{t, \hbar}$ belong to the same cohomological class.*

5) From the above results and Remark 2) in page 841 of [6] we may also prove:

Proposition 6.7. *Let $A_{\mathfrak{a}_{t,\hbar} \oplus \mathfrak{a}_{r_{t,\hbar}}^*, \Omega, J_{r_{t,\hbar}}^{-1}}$ and $A_{\mathfrak{a}_{t,\hbar} \oplus \mathfrak{a}_{r'_{t,\hbar}}^*, \Omega, J_{r'_{t,\hbar}}^{-1}}$ be quasitriangular Hopf QUE algebras over $\mathbb{K}_t[[\hbar]]$ which are quantizations, as in Theorem 2.3 after putting $u = \hbar$, of the quasitriangular Lie bialgebra $(\mathfrak{a}_t \oplus \mathfrak{a}_{r_t}^*, [,]_{\mathfrak{a}_t \oplus \mathfrak{a}_{r_t}^*}, \varepsilon_{\mathfrak{a}_t \oplus \mathfrak{a}_{r_t}^*} = d_c(t)r)$ over the ring \mathbb{K}_t . Suppose that the cocycles $\mu_{r_{t,u}}(r_{t,u}) = \beta_{t,u}$ and $\mu_{r'_{t,u}}(r'_{t,u})$ as in Proposition 6.2 define the same cohomological class in $H^2(\mathfrak{a}_t)[[u]]$. Then the quasitriangular Hopf QUE algebras $A_{\mathfrak{a}_{t,\hbar} \oplus \mathfrak{a}_{r_{t,\hbar}}^*, \Omega, J_{r_{t,\hbar}}^{-1}}$ and $A_{\mathfrak{a}_{t,\hbar} \oplus \mathfrak{a}_{r'_{t,\hbar}}^*, \Omega, J_{r'_{t,\hbar}}^{-1}}$ over the ring $\mathbb{K}_t[[\hbar]]$ are isomorphic.*

7. Isomorphic triangular Hopf algebras over $\mathbb{K}[[\hbar]]$ of type

$$A_{\mathfrak{a}_{\hbar}, \tilde{J}_{r_{\hbar}, \hbar}}$$

We start from a deformation algebra $(\mathfrak{a}_{\hbar}, [,]_{\mathfrak{a}_{\hbar}})$ of the Lie algebra $(\mathfrak{a}, [,]_{\mathfrak{a}})$ over the field \mathbb{K} and from an element $r_{\hbar} = \sum_{l \geq 1} r_l \hbar^{l-1} \in \mathfrak{a}_{\hbar} \wedge \mathfrak{a}_{\hbar}$ which is nondegenerate (r_1 is invertible) and a solution of the YBE ($[r_{\hbar}, r_{\hbar}]_{\mathfrak{a}_{\hbar}} = 0$) on the Lie algebra $(\mathfrak{a}_{\hbar}, [,]_{\mathfrak{a}_{\hbar}})$ over $\mathbb{K}[[\hbar]]$. We call the set $(\mathfrak{a}_{\hbar}, [,]_{\mathfrak{a}_{\hbar}}, r_{\hbar})$ a nondegenerate triangular Lie bialgebra deformation of the nondegenerate triangular Lie bialgebra $(\mathfrak{a}, [,]_{\mathfrak{a}}, r_1)$ over \mathbb{K} . These elements are just the elements $(\mathfrak{a}_t, [,]_{\mathfrak{a}_t})$, $r_t = \sum_{l \geq 1} r_l t^{l-1} \in \mathfrak{a}_t \wedge \mathfrak{a}_t$, $[r_t, r_t]_{\mathfrak{a}_t} = 0$ considered before setting $t = \hbar$. Consider two elements $r_{t,u}$ and $r'_{t,u}$ as in Section 6. We set $u = \hbar$ and obtain $r_{t,\hbar}$ and $r'_{t,\hbar}$, as in that section, and the corresponding propositions there. We set moreover $t = \hbar$ and we get $r_{\hbar,\hbar}, r'_{\hbar,\hbar} \in \mathfrak{a}_{\hbar} \wedge \mathfrak{a}_{\hbar}$. The corresponding elements $\tilde{J}_{r_{\hbar,\hbar}}^{-1}, \tilde{J}_{r'_{\hbar,\hbar}}^{-1} \in \mathcal{U}\mathfrak{a}_{\hbar} \hat{\otimes} \mathcal{U}\mathfrak{a}_{\hbar}$ will also be called Invariant Star Products on the deformation algebra $(\mathfrak{a}_{\hbar}, [,]_{\mathfrak{a}_{\hbar}})$. From Proposition 6 we obtain

Proposition 7.1. *Let $\tilde{J}_{r_{\hbar,\hbar}}$ and $\tilde{J}_{r'_{\hbar,\hbar}} \in \mathcal{U}\mathfrak{a}_{\hbar} \hat{\otimes} \mathcal{U}\mathfrak{a}_{\hbar}$ be the above Invariant Star Products on a nondegenerate triangular Lie bialgebra deformation $(\mathfrak{a}_{\hbar}, [,]_{\mathfrak{a}_{\hbar}}, \varepsilon_{\mathfrak{a}_{\hbar}} = d_c(\hbar)r_{\hbar})$ of the nondegenerate triangular Lie bialgebra $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$ over \mathbb{K} . Then $\tilde{J}_{r_{\hbar,\hbar}}$ and $\tilde{J}_{r'_{\hbar,\hbar}}$ are equivalent Invariant Star Products if, and only if, $\mu_{r_{\hbar,\hbar}}(r_{\hbar,\hbar}) = \beta_{\hbar,\hbar}$ and $\mu_{r'_{\hbar,\hbar}}(r'_{\hbar,\hbar}) = \beta'_{\hbar,\hbar}$ belong to the same cohomological class in $H^2_R(\mathfrak{a}_{\hbar})[[\hbar]]$. In other words, $\tilde{J}_{r_{\hbar,\hbar}}$ and $\tilde{J}_{r'_{\hbar,\hbar}}$ are equivalent Invariant Star Products on $(\mathfrak{a}_{\hbar}, [,]_{\mathfrak{a}_{\hbar}})$ if, and only if, there exists a 1-cochain $\gamma_{\hbar,\hbar} \in \mathfrak{a}_{\hbar}^*$ such that $\beta'_{\hbar,\hbar} = \beta_{\hbar,\hbar} + d_R(\hbar)\gamma_{\hbar,\hbar}$.*

As a consequence we obtain

Corollary 7.2. *The triangular Quantized Universal Enveloping algebras $A_{\mathfrak{a}_{\hbar}, \tilde{J}_{r_{\hbar}, \hbar}}$ and $A_{\mathfrak{a}_{\hbar}, \tilde{J}'_{r'_{\hbar}, \hbar}}$ over $\mathbb{K}[[\hbar]]$ are isomorphic, if and only if, the elements $\beta_{\hbar, \hbar}$ and $\beta'_{\hbar, \hbar} \in (\mathfrak{a}_{\hbar}^* \wedge \mathfrak{a}_{\hbar}^*)[[\hbar]]$ are in the same cohomological class in $H^2_{\mathbb{R}}(\mathfrak{a}_{\hbar})[[\hbar]]$.*

Also, we obtain

Corollary 7.3. *Suppose that the deformation algebra $(\mathfrak{a}_{\hbar}, [;]_{\mathfrak{a}_{\hbar}})$ is a trivial one, that is, it is the one obtained just by extension of scalars $\mathbb{K} \longrightarrow \mathbb{K}[[\hbar]]$. Suppose also that $r_{\hbar} = r_1$ and $r_{\hbar, \hbar} = r_1 + r_2\hbar + r_3\hbar^2 + \dots$. Write it as $r_{0, \hbar}$. Similarly suppose $r'_{\hbar} = r_1$ and $r'_{\hbar, \hbar} = r_1 + r'_2\hbar + r'_3\hbar^2 + \dots$ and write it as $r'_{0, \hbar}$. Let $\beta_{0, \hbar} = \beta_1 + \beta_2\hbar + \beta_3\hbar^2 + \dots$ and $\beta'_{0, \hbar} = \beta_1 + \beta'_2\hbar + \beta'_3\hbar^2 + \dots$ be the corresponding elements in $(\mathfrak{a}^* \wedge \mathfrak{a}^*)[[\hbar]]$. The triangular Hopf Quantized Universal enveloping algebras $A_{\mathfrak{a}_{\hbar}, \tilde{J}_{r_{0, \hbar}}}$ and $A_{\mathfrak{a}_{\hbar}, \tilde{J}'_{r'_{0, \hbar}}}$ are isomorphic if, and only if, β_k and β'_k are, for any $k = 2, 3, 4, \dots$, in the same cohomological class in $H^2(\mathfrak{a})$. In the particular case when \mathbb{K} is the field \mathbb{R} , what we get is that the set of equivalent classes of quantizations (usual term) of the Lie group \mathbb{G} with Lie algebra $(\mathfrak{a}, [,]_{\mathfrak{a}})$ and endowed with a left invariant symplectic structure $\beta_1 \in \mathfrak{a}^* \wedge \mathfrak{a}^*$ is in a bijective correspondence with the set $\beta_1 + \hbar H^2(\mathfrak{a})[[\hbar]]$. A theorem given by Drinfeld in the setting of the quantization in [4].*

References

- [1] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, and D. Sternheimer, *Deformation theory and quantization*, Ann. Phys. **111** (1978), 61 – 151.
- [2] P. Cartier, *Hyperalgèbres et Groups de Lie formels*, Sèminaire Sophus Lie, 2ème anné, Faculté des Sciences de Paris, 19.
- [3] N. Bourbaki, *Algèbre Commutative, chapitres 3 et 4*, Actualités Scientifiques et Industrielles, vol. 1293, Hermann, Paris, 1961.
- [4] V.G. Drinfeld, *On constant, quasiclassical solutions of the Yang-Baxter equations*, Sov. Math. Dokl. **28** (1983), 667 – 671.
- [5] V.G. Drinfeld, *Quasi-Hopf algebras*, Leningrad Math. J. **1** (6) (1990), 1419 – 1457.
- [6] V.G. Drinfeld, *On quasitriangular quasi-Hopf algebras and a group closely connected with $Gal(\overline{\mathbb{Q}}/\mathbb{Q})$* , Leningrad Math. J. **2** (4) (1991), 829 – 860.
- [7] P. Etingof and D. Kazhdan, *Quantization of Lie Bialgebras, I*, Selecta Mathematica **2** (1) (1996), 1 – 41.
- [8] N.Yu. Reshetikhin and M.A. Semenov-Tian-Shansky, *Quantum R-matrices and factorization problems*, J. Geom. Phys. **5** (4) (1988), 533 – 550.
- [9] Y. Kosmann-Schwarzbach, *Integrability of Nonlinear Systems*, Lectures Notes in Physics 495, Springer, 1997, pp. 104-170.
- [10] C. Moreno and L. Valero, in Y.Choquet-Bruhat and C. De-Witt-Morette, Morette, *Analysis, Manifolds and Physics. Part II*, North-Holland, Amsterdam, 2000.
- [11] C. Moreno and L. Valero, *Star products and quantization of Poisson-Lie groups*, J. Geom. Phys. **9** (4) (1992), 369 – 402.

- [12] C. Moreno and L. Valero, *Star products and quantum groups*, in Physics and Manifolds, Mathematical Physics Studies, vol. **15**, Kluwer Academic Publishers, 1994, pp. 203–234.
- [13] C. Moreno and L. Valero, *The set of equivalent classes of invariant star products on $(G; \beta_1)$* , J. Geom. Phys. **23** (4) (1997), 360 – 378.
- [14] C. Moreno and J. Teles, *Preferred quantizations of nondegenerate triangular Lie bialgebras and Drinfeld associators*, in Recent Advances in Lie Theory, Research and Exposition in Mathematics, vol. **25**, Heldermann Verlag, 2002, pp. 346-365.
- [15] J. Teles, *The formal Chevalley cohomology in the quantization by Etingof and Kazhdan of non-degenerate triangular Lie bialgebras and their classical doubles*, Ph.D. thesis, Departamento de Matemática, Universidade de Coimbra, 2003.

CARLOS MORENO

DEPARTAMENTO DE FÍSICA TEÓRICA II, UNIVERSIDAD COMPLUTENSE, E-28040 MADRID, SPAIN.

JOANA TELES

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, COIMBRA, PORTUGAL

E-mail address: `jteles@mat.uc.pt`