

## ON THE UNIFORMIZATION OF $L$ -VALUED FRAMES

J. GUTIÉRREZ GARCÍA, I. MARDONES-PÉREZ, JORGE PICADO  
AND M. A. DE PRADA VICENTE

ABSTRACT: This note discusses the appropriate way of uniformizing the notion of an  $L$ -valued frame introduced by A. Pultr and S. Rodabaugh in [*Lattice-valued frames, functor categories, and classes of sober spaces*, Chapter 6 of *Topological and Algebraic Structures in Fuzzy Sets: A Handbook of Recent Developments in the Mathematics of Fuzzy Sets*, Kluwer, 2003]. It covers the case of a completely distributive lattice  $L$  (which is, in a certain sense, the most general one) and studies the corresponding category of uniform  $L$ -valued frames.

KEYWORDS:  $L$ -valued frames;  $L$ -Frm;  $L$ -topological spaces;  $L$ -Top; uniform  $L$ -valued frames; iota functors; upper/lower forgetful functors.

AMS SUBJECT CLASSIFICATION (2000): 54A40, 54E15, 06D22, 03E72.

### 1. Introduction

$L$ -valued frames are structures of increasing interest for fuzzy topology (see [14], [8], [15], [16], [17], [6], [7]). They were introduced by Pultr and Rodabaugh in [14] for the case of a complete chain  $L$  and were recently extended for the more general case of a completely distributive lattice  $L$  by Gutiérrez García, Höhle and de Prada Vicente [7]. This paper is a continuation of our previous paper [6], and has its roots in the convenience of finding (after [7]) an appropriate notion of a uniform  $L$ -valued frame. There are two obvious candidates for it: the most direct one, provided by the direct approach of uniformizing the  $L$ -topologies as a frame, and the one suggested by [14], based on the concept of an  $L$ -valued frame. In the previous [6] we chose the latter (that would eventually provide a nice categorical picture of the categories at hand) since the former reveals to make no sense after the following observation of Pultr and Rodabaugh in [14]:

---

Received November 6, 2009.

The first, the second and the fourth named authors acknowledge financial support from the Ministry of Education and Science of Spain and FEDER under grant MTM2006-14925-C02-02 and from the University of the Basque Country under grant GIU07/27. The third named author acknowledges financial support from the Centre for Mathematics of the University of Coimbra/FCT.

(...) *we envisage potential applications of the notion of an  $L$ -valued frame in the linear case. One of the questions of interest in fuzzy topology concerns well-founded definitions of the structures of the uniformity type. The case of a complete chain  $L$ , important historically and still very important for applications, does not allow the direct uniformization of an  $L$ -topology  $\tau$  as a frame: a uniformity on  $\tau$  induces a uniformity on  $L$  (as observed by Banaschewski); and since the only linearly ordered frame admitting a uniformity is the two point Boolean algebra  $2 = \{0 < 1\}$  we would be left with the crisp case. One can, however, think of definitions based on the concept of an  $L$ -valued frame which would be more satisfactory.*

However, as we now have found with the following example, this observation is not true in general (it holds only for the stratified case): Let  $L$  be a non-uniformizable frame, for example  $L = [0, 1]$  or any linearly ordered frame different from  $2 = \{0 < 1\}$ . Let  $1_\emptyset$  and  $1_X$  denote, respectively, the bottom and top elements in  $L^X$ . Clearly enough  $\tau = \{1_\emptyset, 1_X\}$  is a uniformizable  $L$ -topology on  $X$ . If the observation above would be true, then there would exist a uniformity on the frame  $L$ , a contradiction.

So, this places again the former alternative as the most natural candidate for a good definition of a uniform  $L$ -valued frame and prompts for its study. This is the problem that we address in this paper, having in mind the treatment of the case of a completely distributive lattice  $L$ , following the lines of [7]. As it is shown in [7], the case of a completely distributive lattice cannot, in a certain sense, be weakened.

## 2. Preliminaries and notation

For standard notions and facts from category theory used here we refer to [1]. As a general reference to frames we suggest [11] or [13].

**2.1. Uniform frames.** A *frame* is a complete lattice  $A$  satisfying the distributive law

$$\forall a \in A, \forall S \subseteq A, \quad a \wedge (\bigvee S) = \bigvee \{a \wedge b \mid b \in S\}.$$

Given two frames  $A$  and  $B$ , a *frame homomorphism*  $h : A \rightarrow B$  is a mapping preserving all joins and finite meets. The category of frames and frame homomorphism will be denoted by  $\mathbf{Frm}$ .

Let  $A$  be a frame. A set  $U \subseteq A$  is a *cover* of  $A$  if  $\bigvee U = 1$ . The set of covers of  $A$  can be preordered: a cover  $U$  *refines* a cover  $V$ , written  $U \leq V$ , if for each  $a \in U$  there is a  $b \in V$  with  $a \leq b$ .

For  $a \in A$ , the element  $st(a, U) = \bigvee \{b \in U \mid b \wedge a \neq 0\}$  is called the *star of  $a$  in  $U$* . Further, for a family  $\mathcal{U}$  of covers of  $A$ , we write

$$a \overset{\mathcal{U}}{\triangleleft} b \text{ if there exists } U \in \mathcal{U} \text{ such that } st(a, U) \leq b.$$

**Remark 2.1.** The following useful properties are easy to check:

- (i)  $a \overset{\mathcal{U}}{\triangleleft} b \implies a \leq b$ ,
- (ii)  $a \leq b \overset{\mathcal{U}}{\triangleleft} c \leq d \implies a \overset{\mathcal{U}}{\triangleleft} d$ ,
- (iii)  $a \overset{\mathcal{U}}{\triangleleft} b, c \overset{\mathcal{U}}{\triangleleft} d \implies a \wedge c \overset{\mathcal{U}}{\triangleleft} b \wedge d$ .

A family  $\mathcal{U}$  of covers of  $A$  is a *uniformity basis* [12] provided that:

- (U1)  $\mathcal{U}$  is a filter basis of the preordered set  $(Cov(A), \leq)$  of all covers of  $A$ .
- (U2) Every  $U \in \mathcal{U}$  has a star-refinement, i.e., for every  $U \in \mathcal{U}$  there is a  $V \in \mathcal{U}$  with

$$st(V) := \{st(a, V) \mid a \in V\} \leq U.$$

- (U3) For every  $a \in A$ ,  $a = \bigvee \{b \in A \mid b \overset{\mathcal{U}}{\triangleleft} a\}$ .

Also, we understand by a *uniformity subbasis* a family of covers of  $A$  such that finite meets of elements of the family constitute a uniformity basis.

A *uniformity* on  $A$  is a filter  $\mathcal{U}$  of covers of  $A$  generated by some uniformity basis. The pair  $(A, \mathcal{U})$  is then called a *uniform frame*. Let  $(A, \mathcal{U})$  and  $(B, \mathcal{V})$  be uniform frames. A frame homomorphism  $h : A \rightarrow B$  is a *uniform homomorphism* if, for every  $U \in \mathcal{U}$ ,  $h[U] = \{h(a) \mid a \in U\} \in \mathcal{V}$ . We denote by  $\mathbf{UFrm}$  the category of uniform frames and uniform homomorphisms.

**2.2. The iota functor  $\iota_L^\Gamma : L\text{-Top} \rightarrow \mathbf{Top}$ .** Let  $L$  denote a complete lattice. An element  $p \in L$  is called *prime* if for each  $a, b \in L$  with  $a \wedge b \leq p$  either  $a \leq p$  or  $b \leq p$ . As in [4], we denote by  $\text{PRIME } L$  the set of all prime elements of  $L$  and  $\text{Spec } L = \text{PRIME } L \setminus \{1\}$ .

Following [7], for each  $\alpha \in L$  we denote by  $\uparrow\alpha$  the set  $\{\beta \in L : \alpha \ll \beta\}$ , where  $\ll$  is the opposite relation of the way-below relation in the lattice  $L^{op}$ :

$$\alpha \ll \beta \equiv \text{for all } S \subseteq L \text{ such that } \bigwedge S \leq \alpha \text{ there exist } \gamma_1, \dots, \gamma_n \in S : \bigwedge_{i=1}^n \gamma_i \leq \beta.$$

An  $L$ -valued topological space [2, 5] (shortly, an  $L$ -topological space) is a pair  $(X, \tau)$  consisting of a set  $X$  and a subset  $\tau$  of  $L^X$  (the  $L$ -valued topology or  $L$ -topology on the set  $X$ ), containing  $1_\emptyset$  and  $1_X$  and closed under finite meets and arbitrary joins (where meets and joins in  $L^X$  are defined pointwisely).

Given two  $L$ -valued topological spaces  $(X, \tau_1), (Y, \tau_2)$  a map  $f : X \rightarrow Y$  is an  $L$ -continuous map if the correspondence  $b \mapsto f^\leftarrow(b) = b \circ f$  maps  $\tau_2$  into  $\tau_1$ . The resulting category will be denoted by  $L\text{-Top}$ .

Of course, when  $L = 2$ , an  $L$ -topological space is precisely a topological space and there is an isomorphism between  $\mathbf{Top}$  and  $L\text{-Top}$ , via the characteristic functor (the one associating to each subset its characteristic function and leaving morphisms unchanged). If  $L$  is a frame then the  $L$ -topologies, being subframes of the frame  $L^X$ , are frames as well.

The well-known iota functor  $\iota_L$ , originally introduced by Lowen [10] for  $L = [0, 1]$  and later on extended by Kubiak [9] to an arbitrary complete lattice, was the original motivation to define chain-valued frames and the corresponding category in [14]. It constructs at the fibre level, for each  $L$ -topology, a traditional topology with subbasis the family of all level sets for all members of the  $L$ -topology:

For each set  $X$  and each  $\alpha \in L$ , the  $\alpha$ -level mapping  $\iota_\alpha : L^X \rightarrow 2^X$  is defined by

$$\iota_\alpha(a) = [a \not\leq \alpha] := \{x \in X \mid a(x) \not\leq \alpha\}, \quad \text{for each } a \in L^X.$$

Now, given an  $L$ -topology  $\tau$  on  $X$ , we consider the associated crisp topology

$$\iota_L^\top(\tau) = \langle \{\iota_\alpha(\tau) \mid \alpha \in L\} \rangle = \langle \{\iota_\alpha(a) \mid a \in \tau, \alpha \in L\} \rangle.$$

**Remark 2.2.** Notice that when  $L$  is a completely distributive lattice, the collection  $\{\iota_p(a) \mid a \in \tau, p \in \text{Spec } L\}$  is also a subbase of the topology  $\iota_L^\top(\tau)$ .

The correspondence  $(X, \tau) \mapsto (X, \iota_L^\top(\tau))$  defines a functor  $\iota_L^\top : L\text{-Top} \rightarrow \mathbf{Top}$ : for each  $L$ -continuous map  $f : (X, \tau_1) \rightarrow (Y, \tau_2)$ ,  $f : (X, \iota_L^\top(\tau_1)) \rightarrow (Y, \iota_L^\top(\tau_2))$  is continuous, since  $b \circ f \in \tau_1$  and  $f^{-1}[\iota_\alpha(b)] = \iota_\alpha(b \circ f)$  for every  $b \in \tau_2$ .

Note that whenever  $L$  is a frame the mapping  $\iota_p : \tau \rightarrow \iota_L^\top(\tau)$  is a frame homomorphism for each  $p \in \text{Spec } L$  (see Corollary 3.5 below; this is not true in general if  $p$  fails to be prime). Consequently we can consider the system of frame homomorphisms

$$(\iota_p : \tau \rightarrow \iota_L^\top(\tau) \mid p \in \text{Spec } L). \quad (2.2.1)$$

**2.3.  $L$ -valued frames.** From now on let  $L$  denote a completely distributive lattice. Recall that any completely distributive lattice is a spatial frame (i.e. a frame isomorphic to the lattice of open sets of some topological space) and therefore in any completely distributive lattice each element is a meet of primes.

An  $L$ -valued frame (shortly, an  $L$ -frame) is a system

$$\mathfrak{A} = (\varphi_p^{\mathfrak{A}} : A^u \rightarrow A^l \mid p \in \text{Spec } L)$$

of frame homomorphisms ( $A^u$  is the *upper frame* and  $A^l$  is the *lower frame*) satisfying the following conditions:

- (F0) For every  $p \in \text{Spec } L$ ,  $\varphi_p^{\mathfrak{A}} = \bigvee \{\varphi_q^{\mathfrak{A}} \mid q \in \uparrow p \cap \text{Spec } L\}$ .
- (F1)  $A^l = \langle \bigcup_{p \in \text{Spec } L} \varphi_p^{\mathfrak{A}}(A^u) \rangle$ . (collectionwise extremally epimorphic)
- (F2) If  $a \neq b$  then  $\varphi_p^{\mathfrak{A}}(a) \neq \varphi_p^{\mathfrak{A}}(b)$  for some  $p \in \text{Spec } L$ . (collectionwise monomorphic)

$L$ -frames were introduced by Pultr and Rodabaugh [14] for the case of a complete chain  $L$  and extended by Gutiérrez García, Höhle and de Prada Vicente [7] for the case of a completely distributive lattice.

Given two  $L$ -frames  $\mathfrak{A}$  and  $\mathfrak{B}$  an  $L$ -frame homomorphism  $h : \mathfrak{A} \rightarrow \mathfrak{B}$  is an ordered pair of frame morphisms

$$(h^u : A^u \rightarrow B^u, h^l : A^l \rightarrow B^l)$$

satisfying

$$\forall p \in \text{Spec } L, h^l \circ \varphi_p^{\mathfrak{A}} = \varphi_p^{\mathfrak{B}} \circ h^u.$$

The resulting category, with composition and identities component-wise in  $\text{Frm}$ , is denoted by  $L\text{-Frm}$ .

Note that for each  $L$ -topological space  $(X, \tau)$  with  $L$  completely distributive, the system (2.2.1) of frame homomorphisms defines an  $L$ -frame [7].

This was the motivating example for the notion of an  $L$ -frame.

We recall also the *upper*, resp. *lower*, forgetful functors

$$\mathcal{U}^u : L\text{-Frm} \rightarrow \text{Frm}, \quad \mathcal{U}^l : L\text{-Frm} \rightarrow \text{Frm}$$

defined by sending  $(\varphi_p^{\mathfrak{A}} : A^u \rightarrow A^l \mid p \in \text{Spec } L)$  to  $A^u$ , resp.  $A^l$ , and  $(h^u, h^l)$  to  $h^u$ , resp.  $h^l$ .

### 3. An extension of the iota functor

Let  $L\text{-UTop}$  denote the category whose objects are of the form  $(X, \tau_X, \mathcal{U}_X)$  where  $\tau_X$  is an  $L$ -topology on  $X$  and  $\mathcal{U}_X$  is a uniformity on the frame  $\tau_X$ , and morphisms  $f : (X, \tau_X, \mathcal{U}_X) \rightarrow (Y, \tau_Y, \mathcal{U}_Y)$  are the  $L$ -continuous functions  $f : (X, \tau_X) \rightarrow (Y, \tau_Y)$  for which  $f^{\leftarrow} : (\tau_Y, \mathcal{U}_Y) \rightarrow (\tau_X, \mathcal{U}_X)$  is a uniform homomorphism.

In the particular case  $L = 2$  the objects of  $2\text{-UTop}$  are topological spaces  $(X, \mathcal{T}_X)$  endowed with a uniformity on the spatial frame  $\mathcal{T}_X$ .

One can then try to extend the iota functor for  $L\text{-UTop}$ :

$$\iota_L^{\text{TUF}} : L\text{-UTop} \rightarrow 2\text{-UTop}.$$

We will do that by defining how it acts on the additional structure (the uniformity on the  $L$ -topology) since in the  $L$ -topology it will act precisely as the iota functor already defined in Subsection (2.2).

For each  $p \in \text{Spec } L$  and each  $\mathcal{A} \subseteq L^X$ , let  $\iota_p[\mathcal{A}] = \{\iota_p(a) : a \in \mathcal{A}\} \subseteq 2^X$ .

We state without proof some basic facts satisfied by the maps  $\{\iota_p \mid p \in \text{Spec } L\}$ :

**Lemma 3.1.** *Let  $\mathcal{A} \subseteq L^X$ ,  $f : X \rightarrow Y$ ,  $a \in L^X$ ,  $b \in L^Y$  and  $p, q \in \text{Spec } L$ . Then:*

- (1)  $\iota_p(\bigvee \mathcal{A}) = \bigcup_{a \in \mathcal{A}} \iota_p(a)$ .
- (2)  $\iota_p(\bigwedge \mathcal{A}) = \bigcap_{a \in \mathcal{A}} \iota_p(a)$  whenever  $\mathcal{A}$  is finite.
- (3)  $p \leq q \Rightarrow \iota_q(a) \subseteq \iota_p(a)$ .
- (4)  $f^{-1}(\iota_p(b)) = \iota_p(f^{\leftarrow}(b))$ .

We have now the following, which is easy to check (cf. [6]):

**Proposition 3.2.** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be covers of the frame  $L^X$  and  $p \in \text{Spec } L$ . Then:*

- (1)  $\iota_p[\mathcal{A}]$  is a cover of  $X$ .
- (2) If  $\mathcal{A} \leq \mathcal{B}$  then  $\iota_p[\mathcal{A}] \leq \iota_p[\mathcal{B}]$ . Hence  $\iota_p[\mathcal{A} \wedge \mathcal{B}] \leq \iota_p[\mathcal{A}]$  and  $\iota_p[\mathcal{A} \wedge \mathcal{B}] \leq \iota_p[\mathcal{B}]$ .
- (3)  $\text{st}(\iota_p(a), \iota_p[\mathcal{A}]) \subseteq \iota_p[\text{st}(a, \mathcal{A})]$ .
- (4) If  $\text{st}(\mathcal{A}) \leq \mathcal{B}$  then  $\text{st}(\iota_p[\mathcal{A}]) \leq \iota_p[\mathcal{B}]$ .

**Corollary 3.3.** *Let  $(X, \tau_X, \mathcal{U}_X) \in L\text{-UTop}$ . The family*

$$\{\iota_p[U] \mid U \in \mathcal{U}_X \text{ and } p \in \text{Spec } L\}$$

*generates (as a subbase) a uniformity  $\iota_L^{\text{UF}}(\mathcal{U}_X)$  on the frame  $\iota_L^{\text{T}}(\tau_X)$  whose base is*

$$\{\iota_p[U] \wedge \iota_q[U] \mid U \in \mathcal{U}_X \text{ and } p, q \in \text{Spec } L\}.$$

*Proof:* It follows from Lemma 3.1 and the properties in Proposition 3.2. Regarding condition (U3) notice that, by Proposition 3.2(3),  $b \stackrel{\mathcal{U}_X}{\triangleleft} a$  implies that  $\iota_p(b) \stackrel{\iota_L^{\text{UF}}(\mathcal{U}_X)}{\triangleleft} \iota_p(a)$ , thus we have  $\iota_p(a) = \bigcup \{\iota_p(b) \mid \iota_p(b) \stackrel{\iota_L^{\text{UF}}(\mathcal{U}_X)}{\triangleleft} \iota_p(a)\}$  for every  $p \in \text{Spec } L$  and every  $a \in \tau_X$ , which suffices to conclude (U3). Indeed: for any element  $H$  of  $\iota_L^{\text{T}}(\tau_X)$ , there are  $a_i, b_i \in \tau_X$  and  $p_i, q_i \in \text{Spec } L$  such that (cf. Remark 2.2)

$$\begin{aligned} H &= \bigcup_{i \in I} (\iota_{p_i}(a_i) \cap \iota_{q_i}(b_i)) \\ &= \bigcup_{i \in I} \left( \bigcup \{ \iota_{p_i}(c) \mid \iota_{p_i}(c) \stackrel{\iota_L^{\text{UF}}(\mathcal{U}_X)}{\triangleleft} \iota_{p_i}(a_i) \} \cap \bigcup \{ \iota_{q_i}(d) \mid \iota_{q_i}(d) \stackrel{\iota_L^{\text{UF}}(\mathcal{U}_X)}{\triangleleft} \iota_{q_i}(b_i) \} \right) \\ &\subseteq \bigcup_{i \in I} \bigcup \{ \iota_{p_i}(c) \cap \iota_{q_i}(d) \mid \iota_{p_i}(c) \cap \iota_{q_i}(d) \stackrel{\iota_L^{\text{UF}}(\mathcal{U}_X)}{\triangleleft} \iota_{p_i}(a_i) \cap \iota_{q_i}(b_i) \} \\ &\subseteq \bigcup \{ G \in \iota_L^{\text{T}}(\tau_X) \mid G \stackrel{\iota_L^{\text{UF}}(\mathcal{U}_X)}{\triangleleft} H \} \subseteq H, \end{aligned}$$

where the last inclusions follow from Remark 2.1.  $\square$

The following is obvious:

**Corollary 3.4.** *If  $f : (X, \tau_X, \mathcal{U}_X) \rightarrow (Y, \tau_Y, \mathcal{U}_Y)$  is a morphism of  $L\text{-UTop}$  then  $f : (X, \iota_L^{\text{T}}(\tau_X), \iota_L^{\text{UF}}(\mathcal{U}_X)) \rightarrow (Y, \iota_L^{\text{T}}(\tau_Y), \iota_L^{\text{UF}}(\mathcal{U}_Y))$  is a morphism of  $2\text{-UTop}$ .  $\square$*

Consequently, we have a functor  $\iota_L^{\text{TUF}} : L\text{-UTop} \rightarrow 2\text{-UTop}$  given on objects by

$$\iota_L^{\text{TUF}}(X, \tau_X, \mathcal{U}_X) = (X, \iota_L^{\text{T}}(\tau_X), \iota_L^{\text{UF}}(\mathcal{U}_X)) \quad \text{for each } (X, \tau_X, \mathcal{U}_X) \in L\text{-UTop}$$

and, on morphisms, for each  $f : (X, \tau_X, \mathcal{U}_X) \rightarrow (Y, \tau_Y, \mathcal{U}_Y)$ ,  $\iota_L^{\text{TUF}}(f) = f$ .

Hence the diagram

$$\begin{array}{ccc}
L\text{-UTop} & \xrightarrow{\iota_L^{\text{UF}}} & 2\text{-UTop} \\
F_L^{\text{UF}} \downarrow & & \downarrow F^{\text{UF}} \\
L\text{-Top} & \xrightarrow{\iota_L^{\text{T}}} & \text{Top}
\end{array}$$

commutes (where  $F_L^{\text{UF}}$  and  $F^{\text{UF}}$  denote, respectively, the forgetful functors).

**Corollary 3.5.** *For each  $(X, \tau_X, \mathcal{U}_X) \in L\text{-UTop}$  and each  $p \in \text{Spec } L$  the mapping  $\iota_p : (\tau_X, \mathcal{U}_X) \rightarrow (\iota_L^{\text{T}}(\tau_X), \iota_L^{\text{UF}}(\mathcal{U}_X))$  is a uniform homomorphism.*

*Proof:* By Lemma 3.1(1) and (2), each  $\iota_p$  is a frame homomorphism. Clearly, it is moreover uniform, that is, for every  $U \in \mathcal{U}_X$ ,  $\iota_p[U] \in \iota_L^{\text{UF}}(\mathcal{U}_X)$ .  $\square$

## 4. $L$ -valued uniform frames

Let us consider the following system of uniform homomorphisms:

$$(\iota_p : (\tau_X, \mathcal{U}_X) \rightarrow (\iota_L^{\text{T}}(\tau_X), \iota_L^{\text{UF}}(\mathcal{U}_X)) \mid p \in \text{Spec } L).$$

**Remarks 4.1.** (1) As it was shown in [7], for each  $p \in \text{Spec } L$ ,

$$\iota_p = \bigvee \{ \iota_q \mid q \in \uparrow p \cap \text{Spec } L \}. \quad (4.1.1)$$

In particular, the assignment  $p \mapsto \iota_p$  is antitone.

(2) We have

$$(\iota_L^{\text{T}}(\tau_X), \iota_L^{\text{UF}}(\mathcal{U}_X)) = \left\langle \bigvee_{p \in \text{Spec } L} \iota_p(\tau_X, \mathcal{U}_X) \right\rangle, \quad (4.1.2)$$

where, by the previous expression we mean that the topology  $\iota_L^{\text{T}}(\tau_X)$  is generated by the subbase  $\{ \iota_p(a) \mid a \in \tau_X \text{ and } p \in \text{Spec } L \}$  and the uniformity  $\iota_L^{\text{UF}}(\mathcal{U}_X)$  is generated by the subbase  $\{ \iota_p[U] \mid U \in \mathcal{U}_X \text{ and } p \in \text{Spec } L \}$ .

(3) For each pair of distinct  $a, b \in \tau_X$  there exists  $x \in X$  such that  $a(x) \neq b(x)$ , hence there exists  $p \in \text{Spec } L$  such that either  $a(x) \leq p$  and  $b(x) \not\leq p$  or  $a(x) \not\leq p$  and  $b(x) \leq p$  and so  $[a \not\leq p] \neq [b \not\leq p]$ . It follows that

$$\text{if } a \neq b \in \tau_X \text{ then } \iota_p(a) \neq \iota_p(b) \text{ for some } p \in \text{Spec } L. \quad (4.1.3)$$

(4) As a consequence of the previous comments, the system of uniform homomorphisms

$$(\iota_p : (\tau_X, \mathcal{U}_X) \rightarrow (\iota_L^{\text{T}}(\tau_X), \iota_L^{\text{UF}}(\mathcal{U}_X)) \mid p \in \text{Spec } L)$$



satisfies conditions (4.1.1), (4.1.2) and (4.1.3) above.

**Proposition 4.2.** *Let  $(A, \mathcal{U})$  be a uniform frame and  $(X, \mathcal{T}_X, \mathcal{U}_X) \in 2\text{-UTop}$ . Let  $(\varphi_p : (A, \mathcal{U}) \rightarrow (\mathcal{T}_X, \mathcal{U}_X) \mid p \in \text{Spec } L)$  be a system of uniform homomorphisms satisfying:*

$$\bullet \text{ for each } p \in \text{Spec } L, \varphi_p = \bigvee \{\varphi_q \mid q \in \uparrow p \cap \text{Spec } L\}. \quad (4.2.1)$$

$$\bullet (\mathcal{T}_X, \mathcal{U}_X) = \langle \bigvee_{p \in \text{Spec } L} \varphi_p((A, \mathcal{U})) \rangle. \quad (4.2.2)$$

$$\bullet \text{ if } a \neq b \in A \text{ then } \varphi_p(a) \neq \varphi_p(b) \text{ for some } p \in \text{Spec } L. \quad (4.2.3)$$

Then there is a uniform frame  $(B, \mathcal{V})$  and a uniform isomorphism  $h : (A, \mathcal{U}) \rightarrow (B, \mathcal{V})$  satisfying the following:

- (1)  $B$  is an  $L$ -topology on  $X$  (hence  $(X, B, \mathcal{V}) \in L\text{-UTop}$ ).
- (2)  $\iota_L^{\text{UF}}(X, B, \mathcal{V}) = (X, \mathcal{T}_X, \mathcal{U}_X)$ .
- (3) For each  $p \in \text{Spec } L$ ,  $\iota_p \circ h = \varphi_p$ .

*Proof:* The proof follows the lines of Proposition 2.5 in [7].

Given  $a \in A$ , let  $h(a) \in L^X$  be the  $L$ -valued function induced by the family  $\{\varphi_p(a) \mid p \in \text{Spec } L\}$ , that is,

$$h(a)(x) = \bigwedge \{p \in \text{Spec } L \mid x \notin \varphi_p(a)\} \quad \text{for every } x \in X.$$

Take  $B = \{h(a) \mid a \in A\}$  and  $\mathcal{V} = \{h[U] : U \in \mathcal{U}\}$ . It is easy to check (see [7, Proposition 2.5]) that condition (4.2.1) implies that  $h(a)(x) \leq p \iff x \notin \varphi_p(a)$  and so  $\iota_p(h(a)) = [h(a) \not\leq p] = \varphi_p(a)$  for every  $a \in A$  and  $p \in \text{Spec } L$ , i.e.,  $\iota_p \circ h = \varphi_p$  for every  $p \in \text{Spec } L$ .

We now check that  $h$  is injective: Given  $a \neq b \in A$ , by (4.2.3) we have  $\varphi_p(a) \neq \varphi_p(b)$  for some  $p \in \text{Spec } L$ . We can assume, without loss of generality, that there exists  $x \in \varphi_p(a)$  such that  $x \notin \varphi_p(b)$ . Then  $h(b)(x) \leq p$  and  $h(a)(x) \not\leq p$  which implies  $h(b) \neq h(a)$ .

Moreover,  $h$  is a frame homomorphism. Indeed:

Let  $a, b \in A$ ,  $x \in X$  and  $p \in \text{Spec } L$ . We have

$$\begin{aligned} h(a \wedge b)(x) \leq p &\iff x \notin \varphi_p(a \wedge b) = \varphi_p(a) \cap \varphi_p(b) \\ &\iff x \notin \varphi_p(a) \text{ or } x \notin \varphi_p(b) \\ &\iff h(a)(x) \leq p \text{ or } h(b)(x) \leq p \\ &\iff h(a)(x) \wedge h(b)(x) \leq p \end{aligned}$$

and therefore, since  $L$  is a spatial frame, we conclude that  $h(a \wedge b) = h(a) \wedge h(b)$ . Further,  $h(1) = 1$ , since  $\varphi_p(1) = X$  for every  $p \in \text{Spec } L$ . Thus  $h$  preserves finite meets.

On the other hand, given  $\{a_i\}_{i \in I} \subseteq A$ ,  $x \in X$  and  $p \in \text{Spec } L$  we have

$$\begin{aligned} h(\bigvee_{i \in I} a_i)(x) \leq p &\iff x \notin \varphi_p(\bigvee_{i \in I} a_i) = \bigvee_{i \in I} \varphi_p(a_i) \\ &\iff x \notin \varphi_p(a_i) \text{ for all } i \in I \\ &\iff h(a_i)(x) \leq p \text{ for all } i \in I \\ &\iff \bigvee_{i \in I} h(a_i)(x) \leq p \end{aligned}$$

and thus, once again because  $L$  is a spatial frame, we may conclude that  $h(\bigvee_{i \in I} a_i) = \bigvee_{i \in I} h(a_i)$ .

Hence  $B = h(A)$  is a subframe of  $L^X$ , i.e. an  $L$ -topology on  $X$ , and therefore,  $h : A \rightarrow B$  is a frame isomorphism. The latter fact makes straightforward the proof that  $\mathcal{V} = \{h[U] : U \in \mathcal{U}\}$  is a uniformity on  $B$  and that  $h : (A, \mathcal{U}) \rightarrow (B, \mathcal{V})$  is a uniform isomorphism, since  $h$  and  $h^{-1}$  are trivially uniform.

Finally, it follows from (4.2.2) that

$$\begin{aligned} \iota_L^{\top}(B) &= \langle \{\iota_p(h(a)) \mid a \in A, p \in \text{Spec } L\} \rangle \\ &= \langle \{\varphi_p(a) \mid a \in A, p \in \text{Spec } L\} \rangle = \mathcal{T}_X \end{aligned}$$

and

$$\begin{aligned} \iota_L^{\text{UF}}(\mathcal{V}) &= \langle \{\iota_p[h(U)] \mid U \in \mathcal{U}, p \in \text{Spec } L\} \rangle \\ &= \langle \{\varphi_p[U] \mid U \in \mathcal{U}, p \in \text{Spec } L\} \rangle = \mathcal{U}_X, \end{aligned}$$

i.e.,  $\iota_L^{\text{TUF}}(X, B, \mathcal{V}) = (X, \mathcal{T}_X, \mathcal{U}_X)$ .  $\square$

Proposition 4.2 leads immediately to the following definition:

**Definition 4.3.** An  $L$ -valued uniform frame  $\mathfrak{A}$  is a system

$$\mathfrak{A} = (\varphi_p^{\mathfrak{A}} : (A^u, \mathcal{U}^u) \rightarrow (A^l, \mathcal{U}^l) \mid p \in \text{Spec } L)$$

of uniform homomorphisms ( $(A^u, \mathcal{U}^u)$  is the *upper uniform frame* and  $(A^l, \mathcal{U}^l)$  is the *lower uniform frame*) satisfying the following conditions:

(UF0) For every  $p \in \text{Spec } L$ ,  $\varphi_p^{\mathfrak{A}} = \bigvee \{\varphi_q^{\mathfrak{A}} \mid q \in \uparrow p \cap \text{Spec } L\}$ .

(UF1)  $(A^l, \mathcal{U}^l) = \langle \bigvee_{p \in \text{Spec } L} \varphi_p^{\mathfrak{A}}((A^u, \mathcal{U}^u)) \rangle$ .

(UF2) If  $a \neq b$  then  $\varphi_p^{\mathfrak{A}}(a) \neq \varphi_p^{\mathfrak{A}}(b)$  for some  $p \in \text{Spec } L$ .

An  $L$ -uniform homomorphism  $h : \mathfrak{A} \rightarrow \mathfrak{B}$  is an ordered pair of uniform homomorphisms

$$(h^u : (A^u, \mathcal{U}^u) \rightarrow (B^u, \mathcal{V}^u), h^l : (A^l, \mathcal{U}^l) \rightarrow (B^l, \mathcal{V}^l))$$

satisfying

$$\forall p \in \text{Spec } L, h^l \circ \varphi_p^{\mathfrak{A}} = \varphi_p^{\mathfrak{B}} \circ h^u.$$

The resulting category, with composition and identities component-wise in  $\mathbf{UFrm}$ , is denoted by  $L\text{-UFrm}$ .

**Remarks 4.4.** (1) Note that any  $L$ -valued uniform frame is, in particular, an  $L$ -valued frame.

(2) For  $L = 2$ , we have exactly one uniform homomorphism  $\varphi_0^{\mathfrak{A}}$  which automatically satisfies (UF0): conditions (UF1) and (UF2) imply that  $\varphi_0^{\mathfrak{A}}$  is an isomorphism  $(A^u, \mathcal{U}^u) \rightarrow (A^l, \mathcal{U}^l)$ . Thus a 2-uniform frame is a pair of isomorphic uniform frames. Further, each 2-uniform homomorphism is a pair of uniform homomorphisms  $(h^u, h^l)$  such that each factors through the other via isomorphisms. Each 2-uniform frame is a functor  $\mathbf{2} \rightarrow \mathbf{UFrm}$  where  $\mathbf{2}$  denotes the category

$$\begin{array}{ccc} \circlearrowleft & \xrightarrow{\quad} & \circlearrowright \\ \bullet & & \bullet \end{array}$$

It is then easy to conclude that the category  $\mathbf{2}\text{-UFrm}$  of 2-valued uniform frames is the category  $\mathbf{UFrm}^{\mathbf{2}}$  of functors  $\mathbf{2} \rightarrow \mathbf{UFrm}$  and natural transformations between such functors. Consequently,  $\mathbf{2}\text{-UFrm}$  is equivalent to the category  $\mathbf{UFrm}$ , since  $\mathbf{UFrm}^{\mathbf{2}}$  is a category equivalent to  $\mathbf{UFrm}$  via functors  $F : \mathbf{UFrm}^{\mathbf{2}} \rightarrow \mathbf{UFrm}$ , with  $F((A^u, \mathcal{U}^u), (A^l, \mathcal{U}^l)) = (A^u, \mathcal{U}^u)$  and  $F(h^u, h^l) = h^u$ , and  $G : \mathbf{UFrm} \rightarrow \mathbf{UFrm}^{\mathbf{2}}$ , with  $G(A, \mathcal{U}) = ((A, \mathcal{U}), (A, \mathcal{U}))$  and  $G(h) = (h, h)$ .

(3) For a general completely distributive  $L$ ,  $L\text{-UFrm}$  is a full subcategory of the category  $\mathbf{UFrm}^{\mathfrak{L}}$  where  $\mathfrak{L}$  denotes the category

$$\begin{array}{ccc} & p \in \text{Spec } L & \\ & \xrightarrow{\quad} & \\ \circlearrowleft & \xrightarrow{\quad} & \circlearrowright \\ \bullet & & \bullet \\ & \vdots & \\ & \xrightarrow{\quad} & \end{array}$$

(the morphisms between  $\mathfrak{u}$  and  $\mathfrak{l}$  are indexed by  $\text{Spec } L$ ).

**Theorem 4.5.** *The category  $L\text{-UFrm}$  is complete and cocomplete.*

*Proof:* Let  $\text{UF}_0$  and  $\text{UF}_1$  denote the categories defined by relaxing the axioms for objects in the definition of  $L\text{-UFrm}$  as follows: in  $\text{UF}_0$  the objects are systems of uniform homomorphisms for which only (UF0) is required and in  $\text{UF}_1$  the objects are systems of uniform homomorphisms for which only (UF0) and (UF1) are required.

**Claim 1.** *The category  $\text{UF}_0$  is complete and cocomplete.*

**Proof of Claim 1.** Given any family of objects  $(\mathfrak{A}_i)_{i \in I}$ , with  $\mathfrak{A}_i = (\varphi_p^{\mathfrak{A}_i} : (A_i^u, \mathcal{U}_i^u) \rightarrow (A_i^l, \mathcal{U}_i^l) \mid p \in \text{Spec } L)$ , in  $\text{UF}_0$ , consider the uniform frame coproduct [18]  $(A^u, \mathcal{U}^u)$  of the  $(A_i^u, \mathcal{U}_i^u)$  (with injections  $u_i^u$ ) and the uniform frame coproduct  $(A^l, \mathcal{U}^l)$  of the  $(A_i^l, \mathcal{U}_i^l)$  (with injections  $u_i^l$ ). Then, for each  $p \in \text{Spec } L$ , there exists a unique  $\varphi_p^{\mathfrak{A}} : (A^u, \mathcal{U}^u) \rightarrow (A^l, \mathcal{U}^l)$  such that  $\varphi_p^{\mathfrak{A}} \circ u_i^u = u_i^l \circ \varphi_p^{\mathfrak{A}_i}$  for every  $i \in I$ :

$$\begin{array}{ccc} (A_i^u, \mathcal{U}_i^u) & \xrightarrow{u_i^u} & (A^u, \mathcal{U}^u) \\ \downarrow \varphi_p^{\mathfrak{A}_i} & & \downarrow \varphi_p^{\mathfrak{A}} \\ (A_i^l, \mathcal{U}_i^l) & \xrightarrow{u_i^l} & (A^l, \mathcal{U}^l) \end{array}$$

Further,  $\mathfrak{A} = (\varphi_p^{\mathfrak{A}} : (A^u, \mathcal{U}^u) \rightarrow (A^l, \mathcal{U}^l))_{p \in \text{Spec } L}$  is in  $\text{UF}_0$ . Indeed, for each  $i$ ,

$$\begin{aligned} \varphi_p^{\mathfrak{A}} \circ u_i^u &= u_i^l \circ \varphi_p^{\mathfrak{A}_i} = u_i^l \circ \bigvee \{ \varphi_q^{\mathfrak{A}_i} \mid q \in \uparrow p \cap \text{Spec } L \} \\ &= \bigvee \{ u_i^l \circ \varphi_q^{\mathfrak{A}_i} \mid q \in \uparrow p \cap \text{Spec } L \} \\ &= \bigvee \{ \varphi_q^{\mathfrak{A}} \circ u_i^u \mid q \in \uparrow p \cap \text{Spec } L \} \\ &= (\bigvee \{ \varphi_q^{\mathfrak{A}} \mid q \in \uparrow p \cap \text{Spec } L \}) \circ u_i^u, \end{aligned}$$

from which it readily follows that  $\varphi_p^{\mathfrak{A}} = \bigvee \{ \varphi_q^{\mathfrak{A}} \mid q \in \uparrow p \cap \text{Spec } L \}$ .

Finally,  $(\mathfrak{A}, (u_i^u, u_i^l)_{i \in I}) = ((\varphi_p^{\mathfrak{A}} : (A^u, \mathcal{U}^u) \rightarrow (A^l, \mathcal{U}^l))_{p \in \text{Spec } L}, (u_i^u, u_i^l)_{i \in I})$  is the coproduct in  $\text{UF}_0$  of the system of all  $\mathfrak{A}_i$ . Indeed, given any  $\mathfrak{B} = (\varphi_p^{\mathfrak{B}} : (B^u, \mathcal{V}^u) \rightarrow (B^l, \mathcal{V}^l))_{p \in \text{Spec } L} \in \text{UF}_0$  and any collection of  $L$ -uniform homomorphisms  $\{h_i = (h_i^u, h_i^l) : \mathfrak{A}_i \rightarrow \mathfrak{B} \mid i \in I\}$ , since  $(A^u, \mathcal{U}^u)$  is the coproduct in  $\text{UFrm}$  of the  $(A_i^u, \mathcal{U}_i^u)$ , there exists a unique uniform homomorphism  $h^u : (A^u, \mathcal{U}^u) \rightarrow (B^u, \mathcal{V}^u)$  such that  $h^u \circ u_i^u = h_i^u$  for every  $i \in I$ . Similarly,

there exists a unique uniform homomorphism  $h^l : (A^l, \mathcal{U}^l) \rightarrow (B^l, \mathcal{V}^l)$  such that  $h^l \circ u_i^l = h_i^l$  for every  $i \in I$ .

$$\begin{array}{ccccc}
 & & h_i^u & & \\
 & \xrightarrow{\quad u_i^u \quad} & & \xrightarrow{\quad h^u \quad} & \\
 (A_i^u, \mathcal{U}_i^u) & \longrightarrow & (A^u, \mathcal{U}^u) & \dashrightarrow & (B^u, \mathcal{V}^u) \\
 \downarrow \varphi_p^{\mathfrak{A}_i} & & \downarrow \varphi_p^{\mathfrak{A}} & & \downarrow \varphi_p^{\mathfrak{B}} \\
 (A_i^l, \mathcal{U}_i^l) & \longrightarrow & (A^l, \mathcal{U}^l) & \dashrightarrow & (B^l, \mathcal{V}^l) \\
 & \xrightarrow{\quad u_i^l \quad} & & \xrightarrow{\quad h^l \quad} & \\
 & & h_i^l & & 
 \end{array}$$

Since, for each  $i \in I$  and  $p \in \text{Spec } L$ ,

$$\varphi_p^{\mathfrak{B}} \circ h^u \circ u_i^u = \varphi_p^{\mathfrak{B}} \circ h_i^u = h_i^l \circ \varphi_p^{\mathfrak{A}_i} = h^l \circ u_i^l \circ \varphi_p^{\mathfrak{A}_i} = h^l \circ \varphi_p^{\mathfrak{A}} \circ u_i^u,$$

then  $\varphi_p^{\mathfrak{B}} \circ h^u = h^l \circ \varphi_p^{\mathfrak{A}}$ , and the pair  $(h^u, h^l)$  is an  $L$ -uniform homomorphism  $\mathfrak{A} \rightarrow \mathfrak{B}$ . It is clearly the unique  $L$ -uniform homomorphism  $\mathfrak{A} \rightarrow \mathfrak{B}$  such that  $(h^u, h^l) \circ (u_i^u, u_i^l) = (h_i^u, h_i^l)$  for every  $i \in I$ .

In a similar way, one can construct the coequalizers, products and equalizers of  $\text{UF}_0$  from the corresponding constructions in  $\text{UFrm}$ .  $\square$

**Claim 2.**  $\text{UF}_1$  is mono-coreflective in  $\text{UF}_0$ . Consequently,  $\text{UF}_1$  inherits colimits from  $\text{UF}_0$  and the limits of  $\text{UF}_1$  are the coreflections of limits of  $\text{UF}_0$ , and hence  $\text{UF}_1$  is also complete and cocomplete.

**Proof of Claim 2.** Let  $\mathfrak{B} = (\varphi_p^{\mathfrak{B}} : (B^u, \mathcal{V}^u) \rightarrow (B^l, \mathcal{V}^l))_{p \in \text{Spec } L} \in \text{UF}_0$  be given and consider  $\mathfrak{A} = (\varphi_p^{\mathfrak{A}} : (A^u, \mathcal{U}^u) \rightarrow (A^l, \mathcal{U}^l))_{p \in \text{Spec } L}$  where  $(A^u, \mathcal{U}^u) = (B^u, \mathcal{V}^u)$ ,  $A^l$  is the subframe of  $B^l$  generated by  $\bigcup_{p \in \text{Spec } L} \varphi_p^{\mathfrak{B}}[B^u]$ ,  $\mathcal{U}^l$  is the uniformity contained in  $\mathcal{V}^l$  generated by  $\{\varphi_p^{\mathfrak{B}}[V] \mid p \in \text{Spec } L, V \in \mathcal{V}^u\}$  and  $\varphi_p^{\mathfrak{A}} = \varphi_p^{\mathfrak{B}}|_{\text{cod}=\varphi_p^{\mathfrak{B}}[B^u]}$ . It is immediate that  $\mathfrak{A}$  satisfies (UF0) and (UF1). Further, define  $h : \mathfrak{A} \rightarrow \mathfrak{B}$  by  $h^u = \text{id}_{B^u}$  and  $h^l : A^l \hookrightarrow B^l$ . It is straightforward to check that  $h$  is a monomorphism in  $\text{UF}_0$  satisfying the required universal property.  $\square$

**Claim 3.**  $\text{UF}_1$  is an  $(E, \mathcal{M})$ -category, for  $E$  the class  $\text{RegEpi}$  of regular epimorphisms and  $\mathcal{M}$  the class  $\text{Mono-source}$  of mono-sources.

**Proof of Claim 3.** By Proposition 15.13 of [1], it suffices to check that each source in  $\mathbf{UF}_1$  has a (RegEpi, Mono-source)-factorization.

So let  $(h_i : \mathfrak{A} \rightarrow \mathfrak{A}_i)_{i \in I}$  be a mono-source in  $\mathbf{UF}_1$  and consider the product  $((B^u, \mathcal{V}^u), p_i^u)$  (resp.  $((B^l, \mathcal{V}^l), p_i^l)$ ) of the domains  $(A_i^u, \mathcal{U}_i^u)$  (resp. codomains  $(A_i^l, \mathcal{U}_i^l)$ ) of the  $\varphi_p^{\mathfrak{A}_i}$ . Then there exists, for each  $p \in \text{Spec } L$ , a unique  $\varphi_p^{\mathfrak{B}} : (B^u, \mathcal{V}^u) \rightarrow (B^l, \mathcal{V}^l)$  such that  $p_i^l \circ \varphi_p^{\mathfrak{B}} = \varphi_p^{\mathfrak{A}_i} \circ p_i^u$  for every  $i \in I$ .

The function  $f^u : (A^u, \mathcal{U}^u) \rightarrow (B^u, \mathcal{V}^u)$  (resp.  $f^l : (A^l, \mathcal{U}^l) \rightarrow (B^l, \mathcal{V}^l)$ ) defined by  $p_i^u \circ f^u = h_i^u$  for each  $i \in I$  (resp.  $p_i^l \circ f^l = h_i^l$  for each  $i \in I$ ) has a factorization

$$(A^u, \mathcal{U}^u) \xrightarrow{e^u} (C^u, \mathcal{W}^u) \xrightarrow{m^u} (B^u, \mathcal{V}^u)$$

(resp.  $(A^l, \mathcal{U}^l) \xrightarrow{e^l} (C^l, \mathcal{W}^l) \xrightarrow{m^l} (B^l, \mathcal{V}^l)$ ) with  $e^u$  and  $e^l$  surjections and  $m^u$  and  $m^l$  injective functions.

Since, for each  $c \in C^u$ ,  $\varphi_p^{\mathfrak{B}}(m^u(c)) = \varphi_p^{\mathfrak{B}}(m^u(e^u(a))) = \varphi_p^{\mathfrak{B}}(f^u(a)) = f^l(\varphi_p^{\mathfrak{A}}(a)) = m^l(e^l(\varphi_p^{\mathfrak{A}}(a)))$  for some  $a \in A^u$ , we may define

$$\varphi_p^{\mathfrak{C}} : (C^u, \mathcal{W}^u) \rightarrow (C^l, \mathcal{W}^l)$$

by  $\varphi_p^{\mathfrak{C}}(c) = (m^l)^{-1}(\varphi_p^{\mathfrak{B}}(m^u(c))) = e^l(\varphi_p^{\mathfrak{A}}(a))$  (easily seen to be well defined).

Finally, if we define  $m_i^u : C^u \rightarrow A_i^u$  by  $m_i^u = p_i^u \circ m^u$  and  $m_i^l : C^l \rightarrow A_i^l$  by  $m_i^l = p_i^l \circ m^l(a)$  we have the following diagram:

$$\begin{array}{ccccc}
 & & h_i^u & & \\
 & & \curvearrowright & & \\
 (A^u, \mathcal{U}^u) & \xrightarrow{e^u} & (C^u, \mathcal{W}^u) & \xrightarrow{m_i^u} & (A_i^u, \mathcal{U}_i^u) \\
 & \searrow f^u & \downarrow m^u & \nearrow p_i^u & \\
 & & (B^u, \mathcal{V}^u) & & \\
 & & \downarrow \varphi_p^{\mathfrak{B}} & & \\
 & & (B^l, \mathcal{V}^l) & & \\
 & \nearrow f^l & \uparrow m^l & \searrow p_i^l & \\
 (A^l, \mathcal{U}^l) & \xrightarrow{e^l} & (C^l, \mathcal{W}^l) & \xrightarrow{m_i^l} & (A_i^l, \mathcal{U}_i^l) \\
 & \searrow e^l & \curvearrowleft & & \\
 & & h_i^l & & 
 \end{array}$$

$\varphi_p^{\mathfrak{A}}$  (left vertical arrow),  $\varphi_p^{\mathfrak{A}_i}$  (right vertical arrow),  $\varphi_p^{\mathfrak{C}}$  (bottom curved arrow)

It follows that  $(h_i^u, h_i^l) = (m_i^u \circ e^u, m_i^l \circ e^l)$  is a (RegEpi, Mono-source)-factorization of  $(h_i)_{i \in I} = (h_i^u, h_i^l)_{i \in I}$  in  $\mathbf{UF}_1$ .  $\square$

**Claim 4.**  $L\text{-UFrm}$  is a full subcategory of  $\text{UF}_1$  closed under the formation of mono-sources in  $\text{UF}_1$ .

**Proof of Claim 4.** First, notice that for any mono-source  $(h_i : \mathfrak{A} \rightarrow \mathfrak{A}_i)_{i \in I}$  in  $\text{UF}_1$ ,  $(h_i^u)_{i \in I}$  is a mono-source in  $\text{UFrm}$  and, therefore, if  $h_i^u(a) = h_i^u(b)$  for every  $i \in I$  then  $a = b$ . Thus, if each  $\mathfrak{A}_i$  belongs to  $L\text{-UFrm}$  and  $a, b \in A^u$  then, for each  $p \in \text{Spec } L$ ,

$$\begin{aligned} \varphi_p^{\mathfrak{A}}(a) = \varphi_p^{\mathfrak{A}}(b) &\implies h_i^l(\varphi_p^{\mathfrak{A}}(a)) = h_i^l(\varphi_p^{\mathfrak{A}}(b)), \text{ for all } i \in I \\ &\iff \varphi_p^{\mathfrak{A}_i}(h_i^u(a)) = \varphi_p^{\mathfrak{A}_i}(h_i^u(b)), \text{ for all } i \in I \\ &\implies h_i^u(a) = h_i^u(b), \text{ for all } i \in I \\ &\implies a = b, \end{aligned}$$

which shows that  $\mathfrak{A}$  is also in  $L\text{-UFrm}$ . Hence  $L\text{-UFrm}$  is closed under the formation of mono-sources in  $\text{UF}_1$ .  $\square$

In conclusion,  $L\text{-UFrm}$  is a full subcategory of an  $(E, \mathcal{M})$ -category  $\text{UF}_1$ , closed under the formation of  $\mathcal{M}$ -sources in  $\text{UF}_1$ . Hence, by Theorem 16.8 of [1],  $L\text{-UFrm}$  is  $E$ -reflective in  $\text{UF}_1$ . Consequently,  $L\text{-UFrm}$  inherits limits from  $\text{UF}_1$  and the colimits of  $L\text{-UFrm}$  are the reflections of colimits of  $\text{UF}_1$ , and hence  $L\text{-UFrm}$  is complete and cocomplete.  $\square$

## References

- [1] J. Adámek, H. Herrlich, G. E. Strecker, *Abstract and Concrete Categories: The Joy of Cats*, Wiley Interscience Pure and Applied Mathematics, Wiley, 1990.
- [2] C. L. Chang, Fuzzy topological spaces, *J. Math. Anal. Appl.* 24 (1968) 182–190.
- [3] J. Frith, The category of uniform frames, *Cahiers Topologie Géom. Différentielle Catég.* 31 (1990) 305–313.
- [4] G. Gierz, et al., *Continuous Lattices and Domains*, Cambridge University Press, Cambridge, 2003.
- [5] J.A. Goguen, The fuzzy Tychonoff theorem, *J. Math. Anal. Appl.* 43 (1973), 734–742.
- [6] J. Gutiérrez García, I. Mardones-Pérez, J. Picado, M. A. de Prada Vicente, Uniform-type structures on lattice-valued spaces and frames, *Fuzzy Sets and Systems* 159 (2008) 2469–2487.
- [7] J. Gutiérrez García, U. Höhle, M. A. de Prada Vicente, On lattice-valued frames: the completely distributive case, *Fuzzy Sets and Systems*, in press, doi: 10.1016/j.fss.2009.10.024.
- [8] U. Höhle, S. E. Rodabaugh, *Appendix to Chapter 6: weakening the requirement that  $L$  be a complete chain*, in: S. E. Rodabaugh, E. P. Klement (Eds.) *Topological and Algebraic Structures in Fuzzy Sets: A Handbook of Recent Developments in the Mathematics of Fuzzy Sets*, Trends in Logic, vol. 20, Kluwer Academic Publishers, Boston, Dordrecht, London, 2003, pp. 189–197.

- [9] T. Kubiak, *The topological modification of the L-fuzzy unit interval*, in: S. E. Rodabaugh, E. P. Klement, U. Höhle (Eds.), *Applications of Category Theory To Fuzzy Subsets, Theory and Decision Library—Series B: Mathematical and Statistical Methods*, vol. 14, Kluwer Academic Publishers, Boston, Dordrecht, London, 1992, pp. 275–305 (Chapter 11).
- [10] R. Lowen, Fuzzy topological spaces and fuzzy compactness, *J. Math. Anal. Appl.* 56 (1976) 621–633.
- [11] J. Picado, A. Pultr, A. Tozzi, *Locales*, in: M.C. Pedicchio, W. Tholen (Eds.), *Categorical Foundation – Special Topics in Order, Algebra and Sheaf Theory*, Encyclopedia of Mathematics and its Applications, Vol. 97, Cambridge Univ. Press, Cambridge, 2004, pp. 49–101 (Chapter 2).
- [12] A. Pultr, Pointless uniformities I. Complete regularity, *Comment. Math. Univ. Carolinae* 25 (1984) 91–104.
- [13] A. Pultr, *Frames*, in: *Handbook of Algebra*, Vol. 3, North-Holland, Amsterdam, 2003, pp. 791–857.
- [14] A. Pultr, S. E. Rodabaugh, *Lattice-valued frames, functor categories, and classes of sober spaces*, in: S. E. Rodabaugh, E. P. Klement (Eds.) *Topological and Algebraic Structures in Fuzzy Sets: A Handbook of Recent Developments in the Mathematics of Fuzzy Sets*, Trends in Logic, vol. 20, Kluwer Academic Publishers, Boston, Dordrecht, London, 2003, pp. 153–187 (Chapter 6).
- [15] A. Pultr, S. E. Rodabaugh, *Examples for different sobrieties in fixed-basis topology*, in: S. E. Rodabaugh, E. P. Klement (Eds.) *Topological and Algebraic Structures in Fuzzy Sets: A Handbook of Recent Developments in the Mathematics of Fuzzy Sets*, Trends in Logic, vol. 20, Kluwer Academic Publishers, Boston, Dordrecht, London, 2003, pp. 427–440 (Chapter 17).
- [16] A. Pultr, S. E. Rodabaugh, Category theoretic aspects of chain-valued frames: Part I: Categorical and presheaf theoretic foundations, *Fuzzy Sets and Systems* 159 (2008) 501–528.
- [17] A. Pultr, S. E. Rodabaugh, Category theoretic aspects of chain-valued frames: Part II: Applications to lattice-valued topology, *Fuzzy Sets and Systems* 159 (2008) 529–558.
- [18] J. Walters-Wayland, *Completeness and nearly fine uniform frames*, PhD Thesis, University of Cape Town, 1996.

J. GUTIÉRREZ GARCÍA

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DEL PAÍS VASCO-EUSKAL HERRIKO UNIBERTSITATEA, APARTADO 644, 48080, BILBAO, SPAIN  
*E-mail address:* javier.gutierrezgarcia@ehu.es

I. MARDONES-PÉREZ

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DEL PAÍS VASCO-EUSKAL HERRIKO UNIBERTSITATEA, APARTADO 644, 48080, BILBAO, SPAIN  
*E-mail address:* iraide.mardones@ehu.es

JORGE PICADO

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, 3001-454 COIMBRA, PORTUGAL  
*E-mail address:* picado@mat.uc.pt

M. A. DE PRADA VICENTE

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DEL PAÍS VASCO-EUSKAL HERRIKO UNIBERTSITATEA, APARTADO 644, 48080, BILBAO, SPAIN  
*E-mail address:* mariangeles.deprada@ehu.es