

PROPER MAPS FOR LAX ALGEBRAS AND THE KURATOWSKI-MRÓWKA THEOREM

MARIA MANUEL CLEMENTINO AND WALTER THOLEN

ABSTRACT. The characterization of stably closed maps of topological spaces as the closed maps with compact fibres and the role of the Kuratowski-Mrówka' Theorem in this characterization are being explored in the general context of lax (\mathbb{T}, V) -algebras, for a quantale V and a **Set**-monad \mathbb{T} with a lax extension to V -relations. The general results are being applied in standard (topological and metric) and non-standard (labeled graphs) contexts.

1. Introduction

Bourbaki [2] emphasized the importance of *proper* maps of topological spaces, defined as the *stably closed* continuous maps. Point-set topologists prefer to introduce them as the closed continuous maps with compact fibres and to usually call them *perfect* ([9]), often imposing Hausdorff separation conditions in addition. We will employ Bourbaki's terminology in this paper. The statement that closed maps with compact fibres are proper generalizes Kuratowski's Theorem which asserts that $X \rightarrow 1$ is proper when X is compact. Mrówka [16] showed that compactness of X is not only sufficient but also necessary for propriety of $X \rightarrow 1$, which then gives that proper maps have compact fibres.

Extrapolating from the Manes-Barr presentation (see [1]) of topological spaces as the lax relational algebras of the ultrafilter monad (induced by the underlying **Set**-functor of compact Hausdorff spaces [15]), in this paper we consider the question of to which extent the description of proper maps as closed maps with compact fibres may be transferable to the context of lax (\mathbb{T}, V) -algebras, as considered with slight variations in [3, 7, 17, 10] and other papers, where the quantale V replaces the two-element chain (so that V -relations replace ordinary relations) and the **Set**-monad \mathbb{T} replaces the ultrafilter monad. In order not to lose the V -categorical intuition [13], we prefer to call lax (\mathbb{T}, V) -algebras and their lax homomorphisms (\mathbb{T}, V) -categories and (\mathbb{T}, V) -functors, respectively. We define proper (\mathbb{T}, V) -functors as in [4, 12] equationally, as the strict homomorphisms amongst lax, and call an object X compact when $X \rightarrow 1$ is proper, with 1 denoting the terminal object. The

This work was partially supported by the Centro de Matemática da Universidade de Coimbra (CMUC), funded by the European Regional Development Fund through the program COMPETE and by the Portuguese Government through the FCT - Fundação para a Ciência e a Tecnologia under the project PEst-C/MAT/UI0324/2011, and by the Natural Sciences and Engineering Council of Canada.

Received by the editors 2012-05-01 and, in revised form, 2013-01-15.

Published on 2013-02-07 in the volume of articles from CT2011.

2010 Mathematics Subject Classification: 18D20, 54B30, 54D30, 54C10.

Key words and phrases: (\mathbb{T}, V) -category, compact space, proper map, Kuratowski-Mrówka Theorem.

© Maria Manuel Clementino and Walter Tholen, 2013. Permission to copy for private use granted.

terminal structure on a singleton set will generally be distinct from its discrete structure, which is being used when forming fibres. Keeping this distinction in mind, with the known and easily-established pullback stability of proper morphisms one obtains that their fibres are proper as maps, and then compact as objects whenever the terminal structure is discrete.

While there is a compelling definition of “proper map” in the general context, it is less obvious how to define “closed map”. We first pursue a technique that was already used in [12] in some key examples and relies on assigning to every (\mathbb{T}, V) -category structure on X a V -category structure on TX in a functorial manner, such that in the example $V = 2$ and \mathbb{T} the ultrafilter monad, closedness of a continuous map $f : X \rightarrow Y$ is equivalently described as propriety of the monotone map $Tf : TX \rightarrow TY$ in $\mathbf{2-Cat} \cong \mathbf{Ord}$, the category of (pre)ordered sets. (We often omit the prefix “pre” in this paper.) This leads us to the general characterization of proper (\mathbb{T}, V) -functors as those f with proper fibres for which Tf is proper (Theorem 3.2), as presented by the second author at CT2011. We then introduce a family of closure operators which work well when V is constructively completely distributive. Mrówka’s result in the general context relies, as in the topological role model, on the provision of suitable “test objects” (Theorem 5.2). With these at hand, proper (\mathbb{T}, V) -functors can be characterized entirely in terms of closure (Theorem 6.1). However, in the general context there are features not apparent at the level of the role model \mathbf{Top} , and we illustrate them by non-standard examples that leave the realm of categories considered in [12], like the categories of metric and of topological spaces and their natural hybrid, the category of approach spaces [14].

Following a preliminary version of this paper, Solovyov [18] gave a definition of closed morphism in (\mathbb{T}, V) - \mathbf{Cat} that avoids the use of closure operators and the hypothesis that V be constructively completely distributive: see Remark 4.3.

The authors are indebted to Dirk Hofmann who advised them about his proof of Lemma 7.1 in the cases that the quantale V is the two-element chain or the extended non-negative real half line. The proof given here is an easy adaptation of his argumentation to our more general context. They also thank George Janelidze for his careful reading of the paper and various helpful comments for its improvement.

2. The setting

Throughout the paper V is a cartesian closed, unital, associative and commutative quantale. Hence, V is a frame endowed with an associative and commutative binary operation \otimes which, like the binary meet \wedge , preserves arbitrary joins in each variable; in addition, we assume that the top element \top serves as the \otimes -neutral element.

We consider a monad $\mathbb{T} = (T, m, e)$ of the category \mathbf{Set} and, for simplicity, assume that T is *taut*, so that T preserves inverse images (i.e., pullbacks of monomorphisms along arbitrary maps). In particular, T preserves monomorphisms, and for $i : A \hookrightarrow X$ and $\mathfrak{x} \in TX$ we will often write $\mathfrak{x} \in TA$ when $\mathfrak{x} \in Ti(TA)$.

Furthermore, we assume that \mathbb{T} comes with a fixed *lax extension* \hat{T} to the category $V\text{-Rel}$ of V -relations, that is: to the category with objects sets and morphisms $r : X \dashrightarrow Y$ given by functions $r : X \times Y \rightarrow V$ whose composite with $s : Y \dashrightarrow Z$ is defined by

$$(s \cdot r)(x, z) = \bigvee_{y \in Y} r(x, y) \otimes s(y, z)$$

for all $x \in X, z \in Z$. Note that $V\text{-Rel}$ has an involution $r \mapsto r^\circ : Y \dashrightarrow X$ with $r^\circ(y, x) = r(x, y)$, and that every map $f : X \rightarrow Y$ may be considered a V -relation $f_\circ : X \dashrightarrow Y$ with $f_\circ(x, y) = \top$ when $f(x) = y$, and $f_\circ(x, y) = \perp$ (the bottom element) otherwise. Unless $|V| = 1$ there is no danger in identifying f_\circ with f ; its converse, $f^\circ : Y \dashrightarrow X$, serves as the right adjoint to f_\circ in the 2-category $V\text{-Rel}$, the 2-cells of which are given by pointwise order: $r \leq r'$ if and only if $r(x, y) \leq r'(x, y)$ for all $x \in X, y \in Y$.

We must clarify what we mean by *lax extension*: \hat{T} assigns to every V -relation r the V -relation $\hat{T}r : TX \dashrightarrow TY$ subject to the axioms (A)-(F) below.

(A) $Tf \leq \hat{T}f, (Tf)^\circ \leq \hat{T}(f^\circ),$

(B) $r \leq r' \Rightarrow \hat{T}r \leq \hat{T}r',$

(C) $\hat{T}s \cdot \hat{T}r \leq \hat{T}(s \cdot r),$

(D) $\hat{T}\hat{T}r \cdot m_X^\circ = m_Y^\circ \cdot \hat{T}r,$

(E) $r \cdot e_X^\circ \leq e_Y^\circ \cdot \hat{T}r,$

for all $r, r' : X \dashrightarrow Y, s : Y \dashrightarrow Z$ and $f : X \rightarrow Y$. (A)-(E) mean equivalently that $\hat{T} : V\text{-Rel} \rightarrow V\text{-Rel}$ is a lax functor, $m^\circ : \hat{T} \rightarrow \hat{T}\hat{T}$ a natural transformation (so that m satisfies in particular the Beck-Chevalley condition as used in [4, 6]), and $e^\circ : \hat{T} \rightarrow 1$ a lax natural transformation, extending T laxly (in the sense of (A)). They imply in particular the identities

$$\hat{T}(s \cdot f) = \hat{T}s \cdot Tf, \hat{T}(g^\circ \cdot r) = (Tg)^\circ \cdot \hat{T}r, \hat{T}1_X = \hat{T}(e_X^\circ) \cdot m_X^\circ$$

(with $g : Z \rightarrow Y$), see [17, 19]. We require in addition:

(F) $\hat{T}(h \cdot r) = Th \cdot \hat{T}r$

(with $h : Y \rightarrow Z$). We do *not* assume a priori that \hat{T} is *flat*, i.e., that $\hat{T}1_X = 1_{TX}$, which forces the inequalities (A) to become identities.

A (\mathbb{T}, V) -category (X, a) is a set X with a V -relation $a : TX \dashrightarrow X$ with $1_X \leq a \cdot e_X$ and $a \cdot \hat{T}a \leq a \cdot m_X$. A (\mathbb{T}, V) -functor $f : (X, a) \rightarrow (Y, b)$ is a map $f : X \rightarrow Y$ with $f \cdot a \leq b \cdot Tf$. This defines the (ordinary) category $(\mathbb{T}, V)\text{-Cat}$. For $\mathbb{T} = \mathbb{I}$ the identity monad (identically extended to $V\text{-Rel}$), $(\mathbb{T}, V)\text{-Cat}$ is the category $V\text{-Cat}$, i.e. the category of (small) categories enriched over the monoidal-closed category V .

The forgetful functor

$$(\mathbb{T}, V)\text{-Cat} \longrightarrow \mathbf{Set}, \quad (X, a) \longmapsto X, \quad f \longmapsto f$$

is topological, hence $(\mathbb{T}, V)\text{-Cat}$ is complete and cocomplete. In particular, $(1, \top)$, with $1 = \{*\}$ and $\top(\mathfrak{w}, *) = \top$ for every $\mathfrak{w} \in T1$, is the terminal object, and the structure d on the pullback of $f : (X, a) \rightarrow (Z, c)$ and $g : (Y, b) \rightarrow (Z, c)$

$$\begin{array}{ccc} X \times_Z Y & \xrightarrow{q} & Y \\ p \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

is given by

$$d(\mathfrak{w}, (x, y)) = a(Tp(\mathfrak{w}), x) \wedge b(Tq(\mathfrak{w}), y),$$

for any $\mathfrak{w} \in T(X \times_Z Y)$, $(x, y) \in X \times_Z Y$. The left adjoint to the forgetful functor assigns to each set X the *discrete structure*

$$1_X^\sharp = e_X^\circ \cdot \hat{T}1_X.$$

The monad \mathbb{T} may be extended to become a monad of $V\text{-Cat}$ which we again denote by $\mathbb{T} = (T, m, e)$: for a V -category (X, a_0) , let $T(X, a_0) = (TX, \hat{T}a_0)$. There is a comparison functor

$$K : (V\text{-Cat})^\mathbb{T} \longrightarrow (\mathbb{T}, V)\text{-Cat}$$

which commutes with the underlying-set functors; it sends $(X, a_0 : X \dashrightarrow X, \alpha : TX \rightarrow X)$ to $(X, a_0 \cdot \alpha : TX \dashrightarrow X)$ (see [20]). It is less trivial and requires the full extent of hypothesis (D) to show that K has a left adjoint, which sends a (\mathbb{T}, V) -category (X, a) to (TX, \hat{a}, m_X) and a (\mathbb{T}, V) -functor f to Tf , where

$$\hat{a} := \hat{T}a \cdot m_X^\circ$$

(see [11]). We will make use of the composite of this left adjoint with the forgetful functor $(V\text{-Cat})^\mathbb{T} \rightarrow V\text{-Cat}$:

$$(\mathbb{T}, V)\text{-Cat} \longrightarrow V\text{-Cat}, \quad (X, a) \longmapsto (TX, \hat{a}), \quad f \longmapsto Tf.$$

2.1. EXAMPLES.

1. For $V = 2 = \{\text{false} \leq \text{true}\}$, with $\otimes = \&$, an $(\mathbb{I}, 2)$ -category (X, a) is a set X equipped with a (pre)order, that is a relation \leq on X with

$$x \leq x, \quad (x \leq y \ \& \ y \leq z) \Rightarrow x \leq z,$$

for all $x, y, z \in X$ (no anti-symmetry assumed!), while $(\mathbb{I}, 2)$ -functors are exactly monotone maps. We write \mathbf{Ord} for $(\mathbb{I}, 2)\text{-Cat} = 2\text{-Cat}$.

If $V = [0, \infty]$ is the real half-line, ordered by the relation \geq , and $\otimes = +$ (with $v + \infty = \infty$ for every $v \in [0, \infty]$), then an $(\mathbb{I}, [0, \infty])$ -category (X, a) is a set X equipped with a (generalized) metric a , that is a map $a : X \times X \rightarrow [0, \infty]$ such that

$$0 \geq a(x, x), \quad a(x, y) + a(y, z) \geq a(x, z),$$

for all $x, y, z \in X$, and $(\mathbb{I}, [0, \infty])$ -functors are non-expansive maps ([13]). We write **Met** for $(\mathbb{I}, [0, \infty])$ -**Cat** = $[0, \infty]$ -**Cat**.

2. Let $V = 2$ and $\mathbb{P} = (P, m, e)$ be the power-set monad in **Set**, extended to **Rel** by

$$A(\hat{P}r)B \Leftrightarrow \forall x \in A \exists y \in B (x r y),$$

for $r : X \rightrightarrows Y$, $A \subseteq X$ and $B \subseteq Y$. (Note that \hat{P} is a non-flat extension of P .) Then, as shown in [17], $(\mathbb{P}, 2)$ -**Cat** is isomorphic to **Ord**. In particular, every ordered set (X, \leq) defines a $(\mathbb{P}, 2)$ -category (X, \preceq) via

$$A \preceq y \Leftrightarrow \forall x \in A (x \leq y),$$

and conversely.

For every $(X, a) \in (\mathbb{P}, 2)$ -**Cat**, $\hat{a} : PX \rightrightarrows PX$ is defined by

$$\begin{aligned} A \hat{a} B &\Leftrightarrow \exists \mathcal{A} \in PX \ (m_X(\mathcal{A}) = A \ \& \ \mathcal{A}(\hat{P}a) B) \\ &\Leftrightarrow \exists \mathcal{A} \in PX \ (\bigcup \mathcal{A} = A \ \& \ \forall A' \in \mathcal{A} \exists y \in B \ (A' \preceq y)) \\ &\Leftrightarrow \forall x \in A \exists y \in B \ (x \leq y). \end{aligned}$$

3. Let $V = 2$ and $\mathbb{F} = (F, m, e)$ be the filter monad on **Set**, extended to **Rel** by putting

$$\mathfrak{r}(\hat{F}r)\mathfrak{y} \Leftrightarrow \forall B \in \mathfrak{y} \exists A \in \mathfrak{r} \ \forall x \in A \exists y \in Y \ (x r y),$$

for a relation $r : X \rightrightarrows Y$, $\mathfrak{r} \in FX$, $\mathfrak{y} \in FY$. As shown in [17], $(\mathbb{F}, 2)$ -**Cat** is isomorphic to **Top**.

4. When restricted to ultrafilters, \hat{F} gives the lax extension \hat{U} of the ultrafilter **Set**-monad $\mathbb{U} = (U, m, e)$ to **Rel** which may be described by:

$$\mathfrak{r}(\hat{U}r)\mathfrak{y} \Leftrightarrow \forall A \in \mathfrak{r}, B \in \mathfrak{y} \exists x \in A, y \in B \ (x r y),$$

for a relation $r : X \rightrightarrows Y$ and $\mathfrak{r} \in UX$, $\mathfrak{y} \in UY$. As shown by Barr [1], the category $(\mathbb{U}, 2)$ -**Cat** is isomorphic to the category **Top** of topological spaces and continuous maps (see [3, 7] for details).

If (X, a) is an $(\mathbb{U}, 2)$ -category, then the ordered set (UX, \hat{a}) has the following structure:

$$\begin{aligned} \mathfrak{r} \hat{a} \mathfrak{y} &\Leftrightarrow \forall A \subseteq X, A \text{ closed } (A \in \mathfrak{y} \Rightarrow A \in \mathfrak{r}) \\ &\Leftrightarrow \forall A \subseteq X, A \text{ open } (A \in \mathfrak{r} \Rightarrow A \in \mathfrak{y}), \end{aligned}$$

for all $\mathfrak{r}, \mathfrak{v} \in UX$.

In fact, for any V, \mathbb{U} has a flat extension to $V\text{-Rel}$ given by:

$$(\hat{U}r)(\mathfrak{r}, \mathfrak{v}) := \bigwedge_{A \in \mathfrak{r}, B \in \mathfrak{v}} \bigvee_{x \in A, y \in B} r(x, y),$$

for a relation $r : X \rightrightarrows Y, \mathfrak{r} \in UX, \mathfrak{v} \in UY$.

When $V = [0, \infty]$ is the real half-line, it was shown in [3] that $(\mathbb{U}, [0, \infty])\text{-Cat}$ is isomorphic to the category **App** of approach spaces and non-expansive maps [14]. The structure \hat{a} , for a given approach space (X, a) , will be studied in Section 7.

5. Consider now the free-monoid monad $\mathbb{L} = (L, m, e)$ on **Set**, (flatly) extended to **Rel** by putting

$$(x_1, \dots, x_n) (\hat{L}r) (y_1, \dots, y_m) \Leftrightarrow n = m \ \& \ x_i r y_i, \text{ for all } i = 1, \dots, n,$$

for $r : X \rightrightarrows Y, (x_1, \dots, x_n) \in LX, (y_1, \dots, y_m) \in LY$. Then an $(\mathbb{L}, 2)$ -category (X, a) is a *multi-ordered set*, that is, the relation $a : LX \rightrightarrows X$ is such that

$$(x) a x, ((x_1^1, \dots, x_{n_1}^1), \dots, (x_1^l, \dots, x_{n_l}^l)) (\hat{L}a) (y_1, \dots, y_m) a z \Rightarrow (x_1^1, \dots, x_{n_l}^l) a z.$$

6. For a monoid (H, μ, η) , we consider the **Set**-monad $\mathbb{H} = (H \times -, m, e)$, with $m_X = \mu \times 1_X$ and $e_X = \langle \eta, 1_X \rangle$. \mathbb{H} has a flat extension to **Rel** given by

$$(\alpha, x) (\hat{H}r) (\beta, y) \Leftrightarrow \alpha = \beta \ \& \ x r y,$$

for any $r : X \rightrightarrows Y, (\alpha, x) \in H \times X$ and $(\beta, y) \in H \times Y$. Writing $x \xrightarrow{\alpha} y$ instead of $(\alpha, x) a y$ for a relation $a : H \times X \rightrightarrows X$, an $(\mathbb{H}, 2)$ -category (X, a) can be seen as an *H-labeled graph* such that

$$x \xrightarrow{\eta} x, \quad x \xrightarrow{\alpha} y \xrightarrow{\beta} z \Rightarrow x \xrightarrow{\beta \cdot \alpha} z,$$

for all $x, y, z \in X$ and $\alpha, \beta \in H$. An $(\mathbb{H}, 2)$ -functor $f : (X, a) \rightarrow (Y, b)$ is a map $f : X \rightarrow Y$ satisfying the condition:

$$x \xrightarrow{\alpha} y \Rightarrow f(x) \xrightarrow{\alpha} f(y).$$

For each *H*-labeled graph (X, a) , the (pre)order \hat{a} induced on $H \times X$ by a is given by:

$$(\alpha, x) \hat{a} (\beta, y) \Leftrightarrow \exists \gamma \in H (\alpha = \beta \cdot \gamma \ \& \ x \xrightarrow{\gamma} y).$$

3. Proper (\mathbb{T}, V) -functors

A (\mathbb{T}, V) -functor $f : (X, a) \rightarrow (Y, b)$ is *proper* if $f \cdot a = b \cdot Tf$. In order to be able to talk about fibres of f , we should first clarify that very term. For each $y \in Y$, the assignment $* \mapsto y$ defines a (\mathbb{T}, V) -functor $y : (1, 1^\sharp) \rightarrow (Y, b)$, where $1^\sharp = e_1^\circ \cdot \hat{T}1_1$ is the discrete structure on $1 = \{*\}$; explicitly, for $\mathfrak{w} \in T1$,

$$1^\sharp(\mathfrak{w}, *) = \hat{T}1_1(\mathfrak{w}, e_1(*)).$$

By *fibre of f on y* we mean the pullback $(f^{-1}y, \tilde{a}) \rightarrow (1, 1^\sharp)$ of f along the (\mathbb{T}, V) -functor $y : (1, 1^\sharp) \rightarrow (Y, b)$. We note that $(f^{-1}y, \tilde{a}) \rightarrow (X, a)$ is a monomorphism, but in general not regular, i.e., \tilde{a} does not need to be the restriction of $a : TX \times X \rightarrow V$ to $T(f^{-1}y) \times f^{-1}y$:

$$\begin{aligned} \tilde{a}(\mathfrak{r}, x) &= a(\mathfrak{r}, x) \wedge 1^\sharp(T!(\mathfrak{r}), *) && \text{(where } ! : f^{-1}y \rightarrow 1) \\ &= a(\mathfrak{r}, x) \wedge \hat{T}1_X(T!(\mathfrak{r}), e_1(*)), \end{aligned}$$

for every $\mathfrak{r} \in T(f^{-1}y)$ and $x \in f^{-1}y$.

Proper (\mathbb{T}, V) -functors have proper fibres, since:

3.1. PROPOSITION. [See [4]] *Proper maps are stable under pullback in (\mathbb{T}, V) -Cat.*

PROOF. Consider the pullback diagram of Section 2, with f proper. Then

$$\begin{aligned} b \cdot Tq &= (b \wedge b) \cdot Tq \\ &\leq ((g^\circ \cdot c \cdot Tg) \wedge b) \cdot Tq && (g \cdot b \leq c \cdot Tg \Rightarrow b \leq g^\circ \cdot g \cdot b \leq g^\circ \cdot c \cdot Tg) \\ &= (g^\circ \cdot c \cdot Tg \cdot Tq) \wedge (b \cdot Tq) \\ &= (g^\circ \cdot c \cdot Tf \cdot Tp) \wedge (b \cdot Tq) \\ &= (g^\circ \cdot f \cdot a \cdot Tp) \wedge (b \cdot Tq) && (f \text{ proper}) \\ &= (q \cdot p^\circ \cdot a \cdot Tp) \wedge (b \cdot Tq) && (\text{the diagram is a pullback}) \\ &= q \cdot ((p^\circ \cdot a \cdot Tp) \wedge (q^\circ \cdot b \cdot Tq)) && (V \text{ cartesian closed}) \\ &= q \cdot d. \end{aligned}$$

■

We can now prove a first characterization theorem.

3.2. THEOREM. *A (\mathbb{T}, V) -functor $f : (X, a) \rightarrow (Y, b)$ is proper if, and only if, all of its fibres are proper, and the V -functor $Tf : (TX, \hat{a}) \rightarrow (TY, \hat{b})$ is proper.*

PROOF. If f is proper, from $b \cdot Tf = f \cdot a$ one obtains

$$\begin{aligned} \hat{b} \cdot Tf &= \hat{T}b \cdot m_Y^\circ \cdot Tf &= \hat{T}b \cdot \hat{T}\hat{T}f \cdot m_X^\circ && \text{(D)} \\ &\leq \hat{T}(b \cdot \hat{T}f) \cdot m_X^\circ && \text{(C)} \\ &\leq \hat{T}(b \cdot Tf) \cdot m_X^\circ && (*) \\ &= \hat{T}(f \cdot a) \cdot m_X^\circ \\ &= Tf \cdot \hat{T}a \cdot m_X^\circ = Tf \cdot \hat{a}; && \text{(F)} \end{aligned}$$

here (*) comes about since

$$\begin{aligned} b \cdot \hat{T}f &= b \cdot \hat{T}1_Y \cdot Tf = b \cdot \hat{T}(e_Y^\circ) \cdot m_Y^\circ \cdot Tf \\ &\leq b \cdot \hat{T}b \cdot m_Y^\circ \cdot Tf && (1_Y \leq b \cdot e_Y \Rightarrow e_Y^\circ \leq b \cdot e_Y \cdot e_Y^\circ \leq b) \\ &\leq b \cdot m_Y \cdot m_Y^\circ \cdot Tf \leq b \cdot Tf. \end{aligned}$$

Conversely, assume all fibres of f to be proper in (\mathbb{T}, V) -**Cat** and Tf to be proper in V -**Cat**. Since

$$b = b \cdot e_{TY}^\circ \cdot m_Y^\circ \leq e_Y^\circ \cdot \hat{T}b \cdot m_Y^\circ = e_Y^\circ \cdot \hat{b},$$

for all $\mathfrak{x} \in TX$, $y \in Y$ one obtains:

$$\begin{aligned} b \cdot Tf(\mathfrak{x}, y) &= b(Tf(\mathfrak{x}), y) \\ &\leq \hat{b}(Tf(\mathfrak{x}), e_Y(y)) \\ &= \bigvee_{\mathfrak{z} \in (Tf)^{-1}(e_Y(y))} \hat{a}(\mathfrak{x}, \mathfrak{z}) && (Tf \text{ proper}) \\ &= \bigvee_{\mathfrak{z} \in (Tf)^{-1}(e_Y(y))} (\hat{T}a \cdot m_X^\circ)(\mathfrak{x}, \mathfrak{z}) \\ &= \bigvee_{\mathfrak{z} \in (Tf)^{-1}(e_Y(y))} \bigvee_{\mathfrak{x} \in m_X^{-1}\mathfrak{z}} \hat{T}a(\mathfrak{x}, \mathfrak{z}) \otimes \top \end{aligned}$$

Since tautness of T guarantees that the diagram

$$\begin{array}{ccc} T(f^{-1}y) & \xrightarrow{T!} & T1 \\ \downarrow & & \downarrow Ty \\ TX & \xrightarrow{Tf} & TY \end{array}$$

is a pullback, every $\mathfrak{z} \in (Tf)^{-1}(e_Y(y)) = (Tf)^{-1}(Ty(e_1(*)))$ satisfies $\mathfrak{z} \in T(f^{-1}y)$ and $T!(\mathfrak{z}) = e_1(*)$. Using propriety of $(f^{-1}y, \tilde{a}) \rightarrow (1, 1^\#)$ one gets:

$$\begin{aligned} \bigvee_{\mathfrak{z} \in (Tf)^{-1}(e_Y(y))} \bigvee_{\mathfrak{x} \in m_X^{-1}\mathfrak{z}} \hat{T}a(\mathfrak{x}, \mathfrak{z}) \otimes \top &\leq \bigvee_{\mathfrak{z} \in (T!)^{-1}(e_1(*))} \bigvee_{\mathfrak{x} \in m_X^{-1}\mathfrak{z}} \hat{T}a(\mathfrak{x}, \mathfrak{z}) \otimes \bigvee_{x \in f^{-1}y} \tilde{a}(\mathfrak{z}, x) \\ &\leq \bigvee_{\mathfrak{z} \in (T!)^{-1}(e_1(*))} \bigvee_{\mathfrak{x} \in m_X^{-1}\mathfrak{z}} \bigvee_{x \in f^{-1}y} \hat{T}a(\mathfrak{x}, \mathfrak{z}) \otimes a(\mathfrak{z}, x) \\ &\leq \bigvee_{\mathfrak{x} \in m_X^{-1}\mathfrak{z}} \bigvee_{x \in f^{-1}y} a(m_X(\mathfrak{x}), x) \\ &\leq \bigvee_{x \in f^{-1}y} a(\mathfrak{x}, x) \\ &= (f \cdot a)(\mathfrak{x}, y). \end{aligned}$$

Hence, f is proper. ■

Next we show that propriety of fibres trivializes whenever the lax natural transformation $e^\circ : \hat{T} \rightarrow 1$ is strict.

3.3. PROPOSITION. *If $e^\circ : \hat{T} \rightarrow 1$ is a natural transformation, then any (\mathbb{T}, V) -functor has proper fibres.*

PROOF. For a (\mathbb{T}, V) -functor $f : (X, a) \rightarrow (Y, b)$ and $y \in Y$, we must show that the diagram

$$\begin{array}{ccc} T(f^{-1}y) & \xrightarrow{T!} & T1 \\ \tilde{a} \downarrow & & \downarrow 1^\sharp \\ f^{-1}y & \xrightarrow{!} & 1 \end{array}$$

commutes, and for that it suffices to consider $\mathfrak{x} \in T(f^{-1}y)$ with

$$1^\sharp(T!(\mathfrak{x}), *) = \hat{T}1(T!(\mathfrak{x}), e_1(*)) > \perp$$

and show $\tilde{a}(\mathfrak{x}, *) = \top$. From the commutativity of the diagram

$$\begin{array}{ccccc} T(f^{-1}y) & \xrightarrow{T!} & T1 & \xrightarrow{\hat{T}1} & T1 \\ e^\circ \downarrow & & e_1^\circ \downarrow & & \downarrow e_1^\circ \\ f^{-1}y & \xrightarrow{!} & 1 & \xrightarrow{1} & 1 \end{array}$$

we first obtain

$$\perp < e_1^\circ \cdot \hat{T}1 \cdot T!(\mathfrak{x}, *) = e_1^\circ \cdot T!(\mathfrak{x}, *) = ! \cdot e^\circ(\mathfrak{x}, *) = \bigvee_{x \in f^{-1}y} e^\circ(\mathfrak{x}, x) = \top,$$

and then

$$! \cdot \tilde{a}(\mathfrak{x}, *) \geq ! \cdot e^\circ(\mathfrak{x}, x) = \top.$$

■

3.4. COROLLARY. *If $e^\circ : \hat{T} \rightarrow 1$ is a natural transformation, then a (\mathbb{T}, V) -functor $f : (X, a) \rightarrow (Y, b)$ is proper if, and only if, the V -functor Tf is proper.*

3.5. REMARK. This Corollary shows that, in Examples 2.1.4 and 2.1.5, propriety of (\mathbb{T}, V) -functors can be characterized at the V -categorical level. However our main example, the ultrafilter monad, shows that the hypothesis that e° be a natural transformation is essential for the validity of the Corollary.

The notion of proper morphism leads to a natural notion of compactness: a (\mathbb{T}, V) -category (X, a) is *compact* whenever $!_X : (X, a) \rightarrow (1, \top)$ is proper. When $T1 \cong 1$, so that the generator $(1, 1^\sharp)$ coincides with the terminal object $(1, \top)$, (X, a) is compact if, and only if, the only fibre of $!_X : (X, a) \rightarrow (1, \top)$ is proper. In general we can prove:

3.6. PROPOSITION. *If (X, a) is a compact (\mathbb{T}, V) -category, then the fibre of the (\mathbb{T}, V) -functor $!_X : (X, a) \rightarrow (1, \top)$ is proper. Furthermore, if \top is the discrete structure 1^\sharp on 1 (in particular, when $T1 \cong 1$), the converse is true.*

PROOF. Let (X, a) be compact and $t : (X, \tilde{a}) \rightarrow (1, 1^\sharp)$ be the fibre of $!_X$ along $* \in 1$. Then, for any $\mathfrak{r} \in TX$, since V is a frame,

$$t \cdot \tilde{a}(\mathfrak{r}, x) = \bigvee_{x \in X} \tilde{a}(\mathfrak{r}, x) = \bigvee_{x \in X} (a(\mathfrak{r}, x) \wedge 1^\sharp(T!(\mathfrak{r}), *)) = \top \wedge 1^\sharp(T!(\mathfrak{r}), *) = 1^\sharp(Tt(\mathfrak{r}), *),$$

so that t is proper. ■

3.7. COROLLARY. *If \top is the discrete structure on 1 , then the following conditions are equivalent, for a (\mathbb{T}, V) -functor f :*

- (i) f is proper;
- (ii) Tf is proper and f has compact fibres.

3.8. COROLLARY. *If \top is the discrete structure on 1 and e° a natural transformation, then every (\mathbb{T}, V) -category is compact.*

We point out that, when the lax extension \hat{T} is flat, $\top = 1^\sharp$ if and only if $T1 \cong 1$, since flatness of \hat{T} gives $1^\sharp(\mathfrak{r}, *) = e_1^\circ(\mathfrak{r}, *) = \top$ only if $\mathfrak{r} = e_1(*)$. In this case it is easily checked that \mathbb{T} must be the identity monad (see [6]).

We will be able to demonstrate easily that Corollary 3.7 generalizes the characterization of the proper maps in **Top** as the closed maps with compact fibres once we have interpreted the condition that “ Tf be proper” to mean equivalently that “ f be closed”. To this end, the next section introduces a suitable notion of closedness.

4. Closed (\mathbb{T}, V) -functors

Recall that an ordered set X is *constructively completely distributive (ccd)* if there are adjunctions

$$\Downarrow \dashv \bigvee \dashv \Downarrow : X \longrightarrow \text{Down}X$$

where $\text{Down}X$ is the lattice of down-closed sets in X , ordered by inclusion (cf. [21]). Writing $x \ll a$ instead of $x \in \Downarrow a$, one then has

$$x \ll a \Leftrightarrow \forall A \subseteq X (a \leq \bigvee A \Rightarrow \exists y \in A (x \leq y)),$$

and $a = \bigvee \{x \in X \mid x \ll a\}$.

Throughout the remainder of the paper, we assume V to be ccd. Fixing $v \in V$, for a (\mathbb{T}, V) -category (X, a) and $A \subseteq X$ we let

$$A^{(v)} := \{x \in X \mid \bigvee_{\mathfrak{r} \in TA} a(\mathfrak{r}, x) \geq v\}.$$

For a (\mathbb{T}, V) -functor $f : (X, a) \rightarrow (Y, b)$ one then has

$$\bigcap_{u \ll v} f(A^{(u)}) \subseteq f(A)^{(v)}.$$

Indeed, if $y \in f(A^{(u)})$ for every $u \ll v$ in V , so that we can write $y = f(x)$ for some $x \in A^{(u)}$, we obtain

$$u \leq \bigvee_{\mathfrak{r} \in TA} a(\mathfrak{r}, x) \leq \bigvee_{\mathfrak{r} \in TA} b(Tf(\mathfrak{r}), y)$$

and, with $Tf(TA) = T(f(A))$ (the Axiom of Choice granted), $v \leq \bigvee_{\mathfrak{h} \in T(f(A))} b(\mathfrak{h}, y)$.

We call $f : (X, a) \rightarrow (Y, b)$ *closed* if

$$\bigcap_{u \ll v} f(A^{(u)}) = f(A)^{(v)}$$

for all $v \in V$, $A \subseteq X$.

4.1. PROPOSITION. *Every proper (\mathbb{T}, V) -functor is closed, and the converse statement holds in $V\text{-Cat}$ (i.e., when $\mathbb{T} = \mathbb{I}$).*

PROOF. Let $f : (X, a) \rightarrow (Y, b)$ in $(\mathbb{T}, V)\text{-Cat}$ be proper, and $y \in f(A)^{(v)}$ for $v \in V$, so that

$$v \leq \bigvee_{\mathfrak{h} \in T(f(A))} b(\mathfrak{h}, y) = \bigvee_{\mathfrak{r} \in TA} b(Tf(\mathfrak{r}), y) \leq \bigvee_{\mathfrak{r} \in TA} \bigvee_{x \in f^{-1}y} a(\mathfrak{r}, x).$$

For every $u \ll v$ one then obtains $\mathfrak{r} \in TA$, $x \in f^{-1}y$ with $u \leq a(\mathfrak{r}, x)$, and $y \in \bigcap_{u \ll v} f(A^{(u)})$ follows.

Let now $\mathbb{T} = \mathbb{I}$ and f be closed. For all $x \in X$, $y \in Y$, with $v := b(f(x), y)$ and $A := \{x\}$, from

$$y \in f(A)^{(v)} \subseteq \bigcap_{u \ll v} f(A^{(u)})$$

one obtains for every $u \ll v$ some $z \in f^{-1}y$ with $a(x, z) \geq u$. Consequently,

$$v = b(f(x), y) \leq \bigvee_{z \in f^{-1}y} a(x, z),$$

as desired. ■

4.2. COROLLARY. *For every (\mathbb{T}, V) -functor $f : (X, a) \rightarrow (Y, b)$, the V -functor $Tf : (TX, \hat{a}) \rightarrow (TY, \hat{b})$ is proper if and only if it is closed.*

4.3. REMARK. Following the appearance of a preliminary version of this paper Solovyov [18] showed that $f : (X, a) \rightarrow (Y, b)$ is closed if, and only if,

$$\forall A \subseteq X \quad (b \cdot Tf \cdot Ti_A \cdot !_{TA}^\circ \leq f \cdot a \cdot Ti_A \cdot !_{TA}^\circ),$$

with $i_A : A \hookrightarrow X$, $!_{TA} : TA \rightarrow 1$ denoting the obvious maps. This characterization makes the implication (proper \Rightarrow closed) trivial and makes it possible to obtain the results of the following two sections without the blanket assumption that V be completely distributive.

5. The Kuratowski-Mrówka Theorem

In order to be able to characterize compactness of a (\mathbb{T}, V) -category (X, a) by the condition

(KM) *the projection $X \times Z \rightarrow Z$ along any (\mathbb{T}, V) -category (Z, c) is closed,*

one needs to provide suitable test objects (Z, c) that can be used in the sufficiency proof of the condition. For that purpose, using a particular instance of a construction given in [5], for every set X and $\mathfrak{x} \in TX$ we consider the set

$$Z := X \cup \{\omega\} \quad (\text{for some } \omega \notin X)$$

and the V -relation $c : TZ \dashrightarrow Z$ with

$$c(\mathfrak{z}, z) = \begin{cases} \top & \text{if } \mathfrak{z} = e_Z(z) \text{ or } (\mathfrak{z} = \mathfrak{x} \text{ and } z = \omega), \\ \perp & \text{else,} \end{cases}$$

for all $\mathfrak{z} \in TZ$, $z \in Z$, assuming $TX \subseteq TZ$ (and $TTX \subseteq TTZ$) without loss of generality. In order to determine when c will provide Z with the structure of a (\mathbb{T}, V) -category, we highlight two convenient properties of the V -relation c :

1. With i denoting the inclusion map $X \hookrightarrow Z$, c satisfies $i^\circ \cdot c = e_X^\circ \cdot (Ti)^\circ$. Consequently, when \hat{T} is flat, $(Ti)^\circ \cdot \hat{T}c = (Te_X)^\circ \cdot (TTi)^\circ$ and in particular

$$\hat{T}c(\mathfrak{z}, \mathfrak{z}) > \perp \Rightarrow \mathfrak{z} = Te_X(\mathfrak{z}) \tag{1}$$

for all $\mathfrak{z} \in TTZ$, $\mathfrak{z} \in TX$.

2. The V -relation c has finite fibres, that is:

$$c^\circ(z) = \{\mathfrak{z} \in TZ \mid c(\mathfrak{z}, z) > \perp\}$$

is finite for all $z \in Z$. Consequently, if the lax natural transformation $e^\circ : \hat{T} \rightarrow 1$ is finitely strict, so that

$$\begin{array}{ccc} TX & \xrightarrow{\hat{T}r} & TY \\ e_X^\circ \downarrow & & \downarrow e_Y^\circ \\ X & \xrightarrow{r} & Y \end{array}$$

commutes strictly whenever r has finite fibres, then $e_Z^\circ \cdot \hat{T}c = c \cdot e_{TZ}^\circ$, and in particular

$$\hat{T}c(\mathfrak{z}, e_Z(z)) > \perp \Rightarrow \exists \mathfrak{w} \in TZ \ (\mathfrak{z} = e_{TZ}(\mathfrak{w}) \ \& \ c(\mathfrak{w}, z) = \top) \tag{2}$$

for all $\mathfrak{z} \in TTZ$, $z \in Z$.

Note that for the ultrafilter monad e° is finitely strict although, as mentioned in Remark 3.5, it is not a strict natural transformation.

5.1. PROPOSITION. *If \hat{T} is flat and e° finitely strict, then (Z, c) is a (\mathbb{T}, V) -category.*

PROOF. It suffices to show

$$\hat{T}c(\mathfrak{z}, \mathfrak{z}) \otimes c(\mathfrak{z}, z) > \perp \Rightarrow c(m_Z(\mathfrak{z}), z) = \top$$

for all $\mathfrak{z} \in TTZ$, $\mathfrak{z} \in TZ$, $z \in Z$. The premiss implies $\hat{T}c(\mathfrak{z}, \mathfrak{z}) > \perp$ and $c(\mathfrak{z}, z) = \top$. If $\mathfrak{z} \in TX$, one obtains $\mathfrak{z} = Te_X(\mathfrak{z}) = Te_Z(\mathfrak{z})$ from (1) and therefore $c(m_Z(\mathfrak{z}), z) = c(\mathfrak{z}, z) = \top$. If $\mathfrak{z} \notin TX$, since $c(\mathfrak{z}, z) = \top$, we must have $z = \omega$ and $\mathfrak{z} = e_Z(\omega)$, and (2) gives $\mathfrak{w} \in TZ$ with $\mathfrak{z} = e_{TZ}(\mathfrak{w})$, and we may conclude again $c(m_Z(\mathfrak{z}), z) = c(\mathfrak{w}, z) = \top$. ■

5.2. THEOREM. *Let \hat{T} be flat and $e^\circ : \hat{T} \rightarrow 1$ be finitely strict. Then a (\mathbb{T}, V) -category (X, a) is compact if, and only if, (KM) holds.*

PROOF. As a pullback of $X \rightarrow 1$, the second projection $q : X \times Z \rightarrow Z$ is proper for every (\mathbb{T}, V) -category (Z, c) when (X, a) is compact, and therefore closed. Conversely, let (X, a) be such that (KM) holds. We must now show

$$\bigvee_{x \in X} a(\mathfrak{x}, x) = \top,$$

for every $\mathfrak{x} \in TX$. For $Z = X \cup \{\omega\}$ and c as defined above, one considers the set $\Delta_X = \{(x, x) \mid x \in X\} \subseteq X \times Z$. Since $q(\Delta_X) = X \subseteq Z$, from $c(\mathfrak{x}, \omega) = \top$ with $\mathfrak{x} \in TX$ one obtains $\omega \in q(\Delta_X)^{(\top)}$, hence

$$\omega \in \bigcap_{u \ll \top} q(\Delta_X^{(u)})$$

by hypothesis. Consequently, for all $u \ll \top$ one can find $x \in X$ with $(x, \omega) \in \Delta_X^{(u)}$, that is (using the product structure of $X \times Z$):

$$\bigvee_{\mathfrak{w} \in T\Delta_X} a(Tp(\mathfrak{w}), x) \wedge c(Tq(\mathfrak{w}), \omega) \geq u,$$

with $p : X \times Z \rightarrow X$ the first projection. For any $\mathfrak{w} \in T\Delta_X$ one has $Tq(\mathfrak{w}) \in TX$, so that when (without loss of generality) $u > \perp$, we must have $c(Tq(\mathfrak{w}), \omega) = \top$ with $Tq(\mathfrak{w}) = \mathfrak{x}$, and then also $Tp(\mathfrak{w}) = \mathfrak{x}$. Hence, for all $\perp < u \ll \top$ we have found an $x \in X$ with $a(\mathfrak{x}, x) \geq u$, which implies

$$\bigvee_{x \in X} a(\mathfrak{x}, x) = \top,$$

as desired. ■

6. Characterization of propriety via closure

We now have all the ingredients that allow for a characterization of propriety of a (\mathbb{T}, V) -functor $f : (X, a) \rightarrow (Y, b)$ in terms of closure, making essential use of the V -functor $Tf : (TX, \hat{a}) \rightarrow (TY, \hat{b})$ again. V continues to be constructively completely distributive.

6.1. THEOREM. *Let $T1 \cong 1$, \hat{T} be flat and e° be finitely strict. Then the following conditions are equivalent for a (\mathbb{T}, V) -functor f :*

- (i) f is proper;
- (ii) every pullback of f is closed, and Tf is closed;
- (iii) all fibres of f are compact, and Tf is closed.

PROOF. (i) \Rightarrow (ii): From Theorem 3.2 and Propositions 3.1 and 4.1. (ii) \Rightarrow (iii): From Theorem 5.2. (iii) \Rightarrow (i): From Corollary 3.7. ■

6.2. REMARK.

- (1) Without the hypothesis $T1 \cong 1$, stably-closed maps need not be proper (see 7.2), and proper maps may have non-compact fibres (see 7.6).
- (2) In Theorem 6.1 we do not know whether the condition that Tf be closed may be removed from (ii) or be replaced in (iii) by the condition that f be closed.

7. Examples

7.1. V -CATEGORIES (See [12].) By Corollary 3.8 every V -category is compact, and by Corollary 4.2 closed V -functors are exactly the proper ones. In case $V = 2$, for a monotone map $f : (X, \leq) \rightarrow (Y, \leq)$,

$$\begin{aligned} f \text{ proper} &\Leftrightarrow \forall x \in X \quad (\uparrow_Y f(x) \subseteq f(\uparrow_X x)) \\ &\Leftrightarrow \forall A \subseteq X \quad (\uparrow_Y f(A) \subseteq f(\uparrow_X A)), \end{aligned}$$

with $\uparrow_X A = \{x' \in X \mid \exists x \in A : x \leq x'\}$.

When $V = [0, \infty]$, for a non-expansive map $f : (X, a) \rightarrow (Y, b)$,

$$\begin{aligned} f \text{ proper} &\Leftrightarrow \forall x \in X, y \in Y \quad b(f(x), y) = \inf\{a(x, x') \mid x' \in X, f(x') = y\} \\ &\Leftrightarrow \forall A \subseteq X, y \in Y \quad b(f(A), y) = \inf\{a(A, x') \mid x' \in X, f(x') = y\}, \end{aligned}$$

with $a(A, x') = \inf_{x \in A} a(x, x')$.

7.2. ORDERED SETS AS $(\mathbb{P}, 2)$ -CATEGORIES (See [12].) Consider the lax extension \hat{P} of the power-set monad as described in Example 2.1.2. Then a monotone map $f : (X, \preceq) \rightarrow (Y, \preceq)$ is proper if, and only if, for all $A \subseteq X$,

$$\uparrow_Y f(A) \subseteq f(\uparrow_X A), \tag{3}$$

where $\uparrow_X A = \{x \in X \mid A \preceq x\}$. Taking $A = \emptyset$ in (3) one sees immediately that proper maps are surjective, while putting $A = \{x\}$ shows that they are $(\mathbb{I}, 2)$ -proper. Here closedness of f is equivalent to surjectivity since, for any $A \subseteq X$, $A^{(\top)} = X$. So, stably-closed $(\mathbb{P}, 2)$ -functors need not be proper. Note, however, that neither of the hypotheses of Theorem 6.1 is satisfied here.

7.3. TOPOLOGICAL SPACES AS $(\mathbb{F}, 2)$ -CATEGORIES If \hat{F} is the lax (non-flat) extension of F considered in Example 2.1.3, an $(\mathbb{F}, 2)$ -functor is proper if, and only if, it is closed (in the ordinary topological sense) and every fibre has a largest element with respect to the underlying (pre)order of X (that is, $x \leq x'$ when $e_X(x) \rightarrow x'$): see [12]. In particular, proper $(\mathbb{F}, 2)$ -functors must be surjective stably-closed maps.

7.4. TOPOLOGICAL AND APPROACH SPACES AS (\mathbb{U}, V) -CATEGORIES For an (\mathbb{U}, V) -category (X, a) and $\mathfrak{x}, \mathfrak{y} \in UX$ one has, by definition,

$$\hat{a}(\mathfrak{x}, \mathfrak{y}) = \bigvee_{\mathfrak{x} \in m_X^{-1}\mathfrak{x}} \hat{U}a(\mathfrak{X}, \mathfrak{y}) = \bigvee_{\mathfrak{x} \in m_X^{-1}\mathfrak{x}} \bigwedge_{A \in \mathfrak{x}, B \in \mathfrak{y}} \bigvee_{\mathfrak{z} \in A, y \in B} a(\mathfrak{z}, y).$$

Using the hypothesis that V is ccd, we first show that $\hat{a}(\mathfrak{x}, \mathfrak{y})$ can be written more conveniently, provided that V is linearly ordered.

7.5. LEMMA. *If the order of V is linear, then $\hat{a}(\mathfrak{x}, \mathfrak{y}) = \bigvee\{u \in V \mid \forall A \in \mathfrak{x} \ (A^{(u)} \in \mathfrak{y})\}$.*

PROOF. For “ \leq ”, consider any $\mathfrak{X} \in UUX$ with $m_X(\mathfrak{X}) = \mathfrak{x}$. It suffices to show that every $u \ll \bigwedge_{A \in \mathfrak{x}, B \in \mathfrak{y}} \bigvee_{\mathfrak{z} \in A, y \in B} a(\mathfrak{z}, y)$ has the property that $A^{(u)} \in \mathfrak{y}$ for all $A \in \mathfrak{x}$. But if for $A \in \mathfrak{x}$

we assume $A^{(u)} \notin \mathfrak{y}$, so that $B := X \setminus A^{(u)} \in \mathfrak{y}$, considering

$$\mathcal{A} := A^\sharp = \{\mathfrak{z} \in UX : A \in \mathfrak{z}\} \in \mathfrak{X} \text{ (since } A \in \mathfrak{x}\text{)}$$

we would conclude

$$u \ll \bigvee_{\mathfrak{z} \in \mathcal{A}, y \in B} a(\mathfrak{z}, y)$$

and therefore $A^{(u)} \cap B \neq \emptyset$, a contradiction.

For “ \geq ”, consider $v \ll \bigvee \{u \in V \mid \forall A \in \mathfrak{r} (A^{(u)} \in \mathfrak{h})\}$ in V . For all $A \in \mathfrak{r}$, $B \in \mathfrak{h}$, the ultrafilter \mathfrak{h} contains $A^{(v)} \cap B \neq \emptyset$, so that $v \leq \bigvee_{\mathfrak{z} \in A^\#} a(\mathfrak{z}, y)$ for some $y \in B$, and

$$v \leq \bigwedge_{B \in \mathfrak{h}} \bigvee_{\mathfrak{z} \in A^\#, y \in B} a(\mathfrak{z}, y)$$

follows for every $A \in \mathfrak{r}$. Now,

$$\mathfrak{F} = \{\mathcal{A} \subseteq UX \mid A^\# \subseteq \mathcal{A} \text{ for some } A \in \mathfrak{r}\}$$

is a filter on UX , and

$$\mathfrak{J} := \{\mathcal{B} \subseteq UX \mid v > \bigwedge_{B \in \mathfrak{h}} \bigvee_{\mathfrak{z} \in B, y \in B} a(\mathfrak{z}, y)\}$$

is an ideal on UX that is disjoint from \mathfrak{F} . Here, in order to establish closure of \mathfrak{J} under binary union we use the linearity of the order of V , as follows: If \mathcal{B} and \mathcal{C} belong to \mathfrak{J} , then

$$v > \bigvee_{\mathfrak{z} \in \mathcal{B}, y \in B} a(\mathfrak{z}, y) \text{ and } v > \bigvee_{\mathfrak{z} \in \mathcal{C}, y \in C} a(\mathfrak{z}, y) \text{ for some } B, C \in \mathfrak{h}, \text{ hence}$$

$$\begin{aligned} v &> \left(\bigvee_{\mathfrak{z} \in \mathcal{B}, y \in B} a(\mathfrak{z}, y) \right) \vee \left(\bigvee_{\mathfrak{z} \in \mathcal{C}, y \in C} a(\mathfrak{z}, y) \right) \\ &\geq \bigvee_{\mathfrak{z} \in \mathcal{B} \cup \mathcal{C}, y \in B \cap C} a(\mathfrak{z}, y), \end{aligned}$$

and then

$$v > \bigwedge_{D \in \mathfrak{h}} \bigvee_{\mathfrak{z} \in \mathcal{B} \cup \mathcal{C}, y \in D} a(\mathfrak{z}, y)$$

since $B \cap C$ belongs to \mathfrak{h} . Now, the filter \mathfrak{F} must be contained in an ultrafilter \mathfrak{X} which does not meet the ideal \mathfrak{J} . By definition of \mathfrak{F} one has $\mathfrak{r} = m_X(\mathfrak{X})$, and by definition of \mathfrak{J}

$$v \leq \bigwedge_{A \in \mathfrak{X}, B \in \mathfrak{h}} \bigvee_{\mathfrak{z} \in A, y \in B} a(\mathfrak{z}, y) \leq \hat{a}(\mathfrak{r}, \mathfrak{h})$$

follows. ■

7.6. PROPOSITION. *If V is linearly ordered, for an (\mathbb{U}, V) -functor $f : (X, a) \rightarrow (Y, b)$ one has:*

$$f \text{ closed} \Leftrightarrow Uf \text{ closed.}$$

PROOF. As a V -functor, Uf is closed if, and only if, it is proper. We must show that propriety of Uf is equivalent to closedness of f . First let f be closed. For $\mathfrak{x} \in UX$, $\eta \in UY$, we must show $\hat{b}(Uf(\mathfrak{x}), \eta) \leq \bigvee_{\mathfrak{z} \in (Uf)^{-1}\eta} \hat{a}(\mathfrak{x}, \mathfrak{z})$, and for that, by Lemma 7.1, it suffices to show that, whenever $u \ll v$ in V with $B^{(v)} \in \eta$ for all $B \in Uf(\mathfrak{x})$, one has some $\mathfrak{z} \in UX$ with $Uf(\mathfrak{z}) = \eta$ and $A^{(u)} \in \mathfrak{z}$ for all $A \in \mathfrak{x}$. But since f is closed, for every $A \in \mathfrak{x}$ one has $f(A)^{(v)} \subseteq f(A^{(u)}) \in \eta$. Therefore, any ultrafilter \mathfrak{z} on X containing the filterbase $\{A^{(u)} \mid A \in \mathfrak{x}\}$ disjoint from the ideal $\{C \subseteq X \mid f(C) \notin \eta\}$ will be as required.

Conversely, let Uf be proper and $y \in f(A)^{(v)}$ with $A \subseteq X$, $v \in V$. For every $u \ll v$ we must show $y \in f(A^{(u)})$. Since every ultrafilter η on Y containing $f(A)$ is the image of an ultrafilter \mathfrak{x} on X containing A , one has:

$$\begin{aligned} u \ll v &\leq \bigvee_{\eta \in Uf(A)} b(\eta, y) \\ &= \bigvee_{\eta \in Uf(A)} \hat{b}(\eta, e_Y(y)) \\ &= \bigvee_{\mathfrak{x} \in UA} \hat{b}(Uf(\mathfrak{x}), e_Y(y)) \\ &= \bigvee_{\mathfrak{x} \in UA} \bigvee_{\mathfrak{x}' \in (Uf)^{-1}(e_Y(y))} \hat{a}(\mathfrak{x}, \mathfrak{x}') \\ &= \bigvee_{\mathfrak{x} \in UA} \bigvee_{\mathfrak{x}' \in (Uf)^{-1}(e_Y(y))} \bigvee \{w \in V \mid \forall B \in \mathfrak{x} (B^{(w)} \in \mathfrak{x}')\}. \end{aligned}$$

Hence there exist $\mathfrak{x} \in UA$, $\mathfrak{x}' \in (Uf)^{-1}(e_Y(y))$ and $w \geq u$ such that $B^{(w)} \in \mathfrak{x}'$ whenever $B \in \mathfrak{x}$. In particular, $A^{(u)} \in \mathfrak{x}'$, and so $f(A^{(u)}) \in Uf(\mathfrak{x}') = e_Y(y)$, that is $y \in f(A^{(u)})$. ■

Since $U1 = 1$, \hat{U} is flat and e° is finitely strict (although not strict in general), Theorem 6.1 gives:

7.7. COROLLARY. *If V is linearly ordered, for an (\mathbb{U}, V) -functor $f : (X, a) \rightarrow (Y, b)$ the following conditions are equivalent:*

- (i) f is proper;
- (ii) f is stably closed;
- (iii) f is closed with compact fibres.

In case $V = 2$ this Corollary recovers the classical results for **Top**, while in case $V = [0, \infty]$ it recovers the results obtained in [8].

7.8. MULTI-ORDERED SETS AS $(\mathbb{L}, 2)$ -CATEGORIES The extension \hat{L} of the free-monoid monad given in Example 2.1.5 is flat, with $L1 \not\cong 1$, and e° a strict natural transformation. Hence, by Corollary 3.4, an $(\mathbb{L}, 2)$ -functor $f : (X, a) \rightarrow (Y, b)$ is proper whenever Lf is closed. Closedness of f does not imply propriety. In fact, an $(\mathbb{L}, 2)$ -functor $f : (X, a) \rightarrow (Y, b)$ is:

1. proper if, and only if, whenever $(f(x_1), \dots, f(x_n)) b y$, there exists $x \in f^{-1}y$ such that $(x_1, \dots, x_n) a x$;
2. closed if, and only if, whenever $(f(x_1), \dots, f(x_n)) b y$, there exists a list (x'_1, \dots, x'_m) , with $\{x'_1, \dots, x'_m\} \subseteq \{x_1, \dots, x_n\}$ and $x \in f^{-1}y$ such that $(x'_1, \dots, x'_m) a x$.

7.9. LABELED GRAPHS AS $(\mathbb{H}, 2)$ -CATEGORIES For the flat extension \hat{H} of $\mathbb{H} = (H \times -, m, e)$ (H a monoid) of Example 2.1.6, we have:

1. Since e° is a natural transformation, every $(\mathbb{H}, 2)$ -functor $f : (X, a) \rightarrow (Y, b)$ has proper fibres; hence,

$$f \text{ proper} \Leftrightarrow Hf \text{ proper} \Leftrightarrow Hf \text{ closed.}$$

2. Although propriety of fibres is trivial, compactness is not: for (X, a) an H -labeled graph,

$$(X, a) \text{ compact} \Leftrightarrow \forall \alpha \in H, x \in X \exists x' \in X (x \xrightarrow{\alpha} x').$$

3. Closed $(\mathbb{H}, 2)$ -functors need not be proper: for an $(\mathbb{H}, 2)$ -functor $f : (X, a) \rightarrow (Y, b)$,

$$f \text{ proper} \Leftrightarrow \forall \alpha \in H, x \in X, y \in Y (f(x) \xrightarrow{\alpha} y \Rightarrow \exists x' \in f^{-1}y (x \xrightarrow{\alpha} x')),$$

$$f \text{ closed} \Leftrightarrow \forall \alpha \in H, x \in X, y \in Y (f(x) \xrightarrow{\alpha} y \Rightarrow \exists x' \in f^{-1}y \exists \beta \in H (x \xrightarrow{\beta} x')).$$

References

- [1] M. Barr, Relational algebras, in: Reports of the Midwest Category Seminar, IV, *Lecture Notes in Mathematics* 137, Springer, Berlin, 1970, pp. 39-55.
- [2] N. Bourbaki, *Topologie Générale*, Herman, Paris, 1961.
- [3] M.M. Clementino, D. Hofmann, Topological features of lax algebras, *Appl. Categ. Structures* 11 (2003), 267–286.
- [4] M.M. Clementino, D. Hofmann, Effective descent morphisms in categories of lax algebras, *Appl. Categ. Structures* 12 (2004), 413–425.
- [5] M.M. Clementino, D. Hofmann, Descent morphisms and a van Kampen Theorem in categories of lax algebras, *Topology Appl.* **159** (2012), 2310–2319.

- [6] M.M. Clementino, D. Hofmann, G. Janelidze, The monads of classical algebra are seldom weakly cartesian, DMUC Preprint 12-46, University of Coimbra 2012.
- [7] M.M. Clementino, W. Tholen, Metric, Topology and Multicategory – A Common Approach, *J. Pure Appl. Algebra* 179 (2003) 13–47.
- [8] E. Colebunders, R. Lowen, P. Wuyts, A Kuratowski-Mrówka theorem in approach theory, *Topology Appl.* 153 (2005), 756–766.
- [9] R. Engelking, *General Topology*, revised and completed edition (Heldermann Verlag, Berlin 1989).
- [10] D. Hofmann, Topological theories and closed objects, *Adv. Math.* **215** (2007) 789-824.
- [11] D. Hofmann, Dualities for distributive spaces, arxiv: math.CT/1009.3892
- [12] D. Hofmann, W. Tholen, Lax algebra meets topology, *Topology Appl.* **159** (2012), 2434–2452.
- [13] F.W. Lawvere, Metric spaces, generalized logic, and closed categories, *Rend. Sem. Mat. Fis. Milano* **43** (1973) 135–166. *Reprints in Theory and Applications of Categories* **1** (2002) 1–37.
- [14] R. Lowen, *Approach Spaces: The missing link in the Topology-Uniformity-Metric Triad*, Oxford Mathematical Monographs (Oxford University Press, Oxford 1997).
- [15] E. Manes, A triple theoretic construction of compact algebras, in: *Lecture Notes in Math.* **80** (Springer, Berlin 1969) pp. 91-118.
- [16] S. Mrówka, Compactness and product spaces, *Colloq. Math.* 7 (1959), 19–22.
- [17] G. Seal, Canonical and op-canonical lax algebras, *Theory Appl. Categ.* 14 (2005), 221-243.
- [18] S. Solovyov, On a lax-algebraic characterization of closed maps. Preprint, Masaryk University, Brno (Czech Republic) 2012.
- [19] W. Tholen, Lax-algebraic Methods in General Topology. Lecture notes. <http://www.math.yorku.ca/~tholen> (2007)
- [20] W. Tholen, Ordered topological structures, *Topology Appl.* 156 (2009), 2148-2157.
- [21] R. Wood, Ordered sets via adjunction, *Categorical Foundations*, pp. 5-47, Encyclopedia Math. Appl., 97, Cambridge Univ. Press, Cambridge, 2004.

*CMUC, Department of Mathematics,
University of Coimbra,
3001-454 Coimbra, Portugal*

*Department of Mathematics and Statistics,
York University,
Toronto, ON M3J 1P3, Canada*

Email: `mmc@mat.uc.pt`

`tholen@mathstat.yorku.ca`

This article may be accessed at <http://www.tac.mta.ca/tac/> or by anonymous ftp at <ftp://ftp.tac.mta.ca/pub/tac/html/volumes/27/14/27-14.{dvi,ps,pdf}>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS The typesetting language of the journal is $\text{T}_{\text{E}}\text{X}$, and $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}2\text{e}$ strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

$\text{T}_{\text{E}}\text{X}$ TECHNICAL EDITOR Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT $\text{T}_{\text{E}}\text{X}$ EDITOR Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS

Clemens Berger, Université de Nice-Sophia Antipolis, cberger@math.unice.fr

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva: valeria.depaiva@gmail.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne : kathryn.hess@epfl.ch

Martin Hyland, University of Cambridge: M.Hyland@dpmms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Edinburgh, Tom.Leinster@ed.ac.uk

Ieke Moerdijk, University of Utrecht: moerdijk@math.uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Edinburgh: Alex.Simpson@ed.ac.uk

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: robert.walters@uninsubria.it

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca