

# SOME REMARKS ON PULLBACKS IN GUMM CATEGORIES

MARINO GRAN AND DIANA RODELO

*Dedicated to Manuela Sobral on the occasion of her seventieth birthday*

ABSTRACT. We extend some properties of pullbacks which are known to hold in a Mal'tsev context to the more general context of Gumm categories. The varieties of universal algebras which are Gumm categories are precisely the congruence modular ones. These properties lead to a simple alternative proof of the known property that central extensions and normal extensions coincide for any Galois structure associated with a Birkhoff subcategory of an exact Goursat category.

## 1. INTRODUCTION

A categorical approach to the property of congruence modularity, well known in universal algebra, was proposed in [5, 6] via a categorical formulation of the so-called *Shifting Lemma* (recalled in Section 3). The categories satisfying this categorical property are called *Gumm categories*, since it was the mathematician H.P. Gumm who proved that, for a variety of universal algebras, the validity of the Shifting Lemma is equivalent to congruence modularity [13]. As examples of Gumm categories, we also have regular Mal'tsev categories [9, 8] and regular Goursat categories [8], which are defined by the property that any pair of equivalence relations  $R$  and  $S$  in  $\mathbb{C}$  (on a same object) 3-permute:  $RSR = SRS$ .

In this context [3] D. Bourn established an interesting permutability result (see Theorem 3.1), that we use in the present paper to prove the following property of regular Gumm categories (Proposition 4.1). Given a commutative

---

2000 *Mathematics Subject Classification.* 18C05, 08B10, 08C05, 18B10, 18E10.

*Key words and phrases.* regular category, Mal'tsev category, Goursat category, Gumm category, congruence modularity, pullback properties.

The second author was supported by the Centro de Matemática da Universidade de Coimbra (CMUC), funded by the European Regional Development Fund through the program COMPETE and by the Portuguese Government through the FCT - Fundação para a Ciência e a Tecnologia under the projects PEst-C/MAT/UI0324/2013 and PTDC/MAT/120222/2010 and grant number SFRH/BPD/69661/2010.



and  $m$  a monomorphism. It is well known that such factorisations are necessarily stable under pullbacks in a regular category. The subobject determined by the monomorphism  $m$  in the factorisation of  $f = m \cdot p$  is usually called the *image* of  $f$ .

A relation  $R$  from  $A$  to  $B$  is a subobject  $\langle r_1, r_2 \rangle: R \rightarrow A \times B$ . The opposite relation, denoted by  $R^\circ$ , is the relation from  $B$  to  $A$  given by the subobject  $\langle r_2, r_1 \rangle: R \rightarrow B \times A$ . A morphism  $f: A \rightarrow B$  can be identified with the relation  $\langle 1_A, f \rangle: A \rightarrow A \times B$ ; its opposite relation  $\langle f, 1_A \rangle: A \rightarrow B \times A$  is usually referred to as  $f^\circ$ . Given another relation  $\langle s_1, s_2 \rangle: S \rightarrow B \times C$  from  $B$  to  $C$ , the composite relation of  $R$  and  $S$  is a relation  $SR$  from  $A$  to  $C$ , obtained as the image of the induced arrow  $\langle r_1\pi_1, s_2\pi_2 \rangle: R \times_B S \rightarrow A \times C$ , where  $(R \times_B S, \pi_1, \pi_2)$  is the pullback of  $r_2$  and  $s_1$ . With the above notation, we can write any relation  $\langle r_1, r_2 \rangle: R \rightarrow A \times B$  as  $R = r_2 r_1^\circ$ . The following properties are well known, and also easy to check (see [8], for instance):

**Lemma 2.1.** *Let  $f: A \rightarrow B$  be any morphism in a regular category  $\mathbb{C}$ . Then:*

- (1)  $ff^\circ f = f$  and  $f^\circ f f^\circ = f^\circ$ ;
- (2)  $ff^\circ = 1_B$  if and only if  $f$  is a regular epimorphism.

A relation  $R$  from an object  $A$  to  $A$  is called a *relation on  $A$* . Such a relation is *reflexive* if  $1_A \leq R$ , *symmetric* if  $R^\circ \leq R$ , and *transitive* when  $RR \leq R$ . A relation  $R$  on  $A$  is called an *equivalence relation* when it is reflexive, symmetric and transitive. Any kernel pair  $\langle f_1, f_2 \rangle: R_f \rightarrow A \times A$  of a morphism  $f: A \rightarrow B$  is an equivalence relation, called an *effective* equivalence relation. By using the composition of relations, it can be written either as  $R_f = f^\circ f$ , or as  $R_f = f_2 f_1^\circ$ . Of course, if  $f = m \cdot p$  is the (regular epimorphism, monomorphism) factorisation of an arbitrary morphism  $f$ , then  $R_f = R_p$ , so that an effective equivalence relation is always the kernel pair of a regular epimorphism.

### 3. GUMM CATEGORIES

A lattice  $(L, \vee, \wedge)$  is called *modular* when, for  $x, y, z \in L$ , one has


$$x \leq z \Rightarrow x \vee (y \wedge z) = (x \vee y) \wedge z.$$

A variety  $\mathcal{V}$  of universal algebras is called *congruence modular* when every lattice of congruences (= effective equivalence relations) on any algebra in  $\mathcal{V}$  is modular. It is well known from [13] that a variety  $\mathcal{V}$  is congruence modular if and only if the following property, called the Shifting Lemma, holds in  $\mathcal{V}$ :

#### Shifting Lemma

Given congruences  $R, S$  and  $T$  on the same algebra  $X$  such that  $R \wedge S \leq T$ ,

whenever  $x, y, t, z$  are elements in  $X$  with  $(x, y) \in R \wedge T$ ,  $(x, t) \in S$ ,  $(y, z) \in S$  and  $(t, z) \in R$ , it then follows that  $(t, z) \in T$ :



This notion has been extended to a categorical context in [6]. Indeed, the property expressed by the Shifting Lemma can be equivalently reformulated in any finitely complete category  $\mathbb{C}$  by asking that a specific class of internal functors are discrete fibrations, as we are now going to recall. For any object  $X$  in  $\mathbb{C}$  and any equivalence relations  $R, S, T$  on  $X$  with

$$R \wedge S \leq T \leq R$$

there is a canonical inclusion  $(i, j): T \square S \rightarrow R \square S$  of equivalence relations, depicted as

$$(3.1) \quad \begin{array}{ccc} T \square S & \xrightarrow{j} & R \square S \\ \pi_1 \downarrow & & \downarrow \pi_1 \\ & & T \\ \pi_2 \downarrow & & \downarrow \pi_2 \\ & \xrightarrow{i} & R \end{array}$$

where  $T \square S$  (respectively,  $R \square S$ ) is the largest double equivalence relation on  $T$  and  $S$  (respectively, on  $R$  and  $S$ ) and  $\pi_1$  and  $\pi_2$  are the projections on  $T$  (respectively, on  $R$ ).

**Definition 3.1.** [6] *A finitely complete category  $\mathbb{C}$  is called a Gumm category when any inclusion  $(i, j): T \square S \rightarrow R \square S$  as in (3.1) is a discrete fibration. This means that any of the commutative squares in (3.1) is a pullback.*

Let us recall that a *Mal'tsev* category  $\mathbb{C}$  is a finitely complete category such that every reflexive relation in  $\mathbb{C}$  is an equivalence relation. A regular category  $\mathbb{C}$  is a Mal'tsev category when the composition of (effective) equivalence relations on any object in  $\mathbb{C}$  is 2-permutable:  $RS = SR$ , where  $R$  and  $S$  are (effective) equivalence relations on a same object (see [9, 8]). The strictly weaker 3-permutability property for (effective) equivalence relations,  $RSR = SRS$ , defines the notion of regular *Goursat categories* [8]. Goursat categories, thus in particular Mal'tsev categories, have the property that every lattice of equivalence relations (on the same object) is modular (Proposition 3.2 in [8]). This

fact implies that any regular Mal'tsev category and, more generally, any regular Goursat category is a Gumm category. Thanks to the characterization theorem in [13] one knows that a variety of universal algebras is congruence modular if and only if it is a Gumm category.

The following property of regular Gumm categories, due to D. Bourn, will play a fundamental role in the next section.

**Theorem 3.1.** (Theorem 7.12 of [3]) *Let  $\mathbb{C}$  be a regular Gumm category. Consider equivalence relations  $R, S$  and  $T$  on a same object such that  $RS = SR$  and  $R \wedge S \leq T \leq R$ . Then  $TS = ST$ .*

#### 4. PULLBACK PROPERTIES IN REGULAR GUMM CATEGORIES

In this section we extend some useful properties of pullbacks from the context of Mal'tsev categories to that of Gumm categories.

The first observation concerns a generalisation of Proposition 3.4 in [7]:

**Proposition 4.1.** *Let  $\mathbb{C}$  be a regular Gumm category. Consider a commutative diagram*

$$(4.2) \quad \begin{array}{ccccc} Z \times_V U & \xrightarrow{x} & X & \xrightarrow{u} & U \\ \varphi \updownarrow \sigma & & f \updownarrow s & & \downarrow w \\ Z & \xrightarrow{y} & Y & \xrightarrow{v} & V, \end{array}$$

such that the whole rectangle is a pullback and the left square  $\boxed{1}$  is composed by vertical split epimorphisms and horizontal regular epimorphisms. Then both squares  $\boxed{1}$  and  $\boxed{2}$  are pullbacks.

**PROOF.** The comparison morphism  $\langle \varphi, x \rangle: Z \times_V U \rightarrow Z \times_Y X$  is clearly a monomorphism. We are now going to show that it is also a regular epimorphism, i.e. that  $\varphi x^\circ = y^\circ f$ . This will imply that the square  $\boxed{1}$  is a pullback.

Consider the effective equivalence relations  $R = R_{ux}, S = R_\varphi$  and  $T = R_x$  (to use the same notations as in Theorem 3.1). The fact that the whole rectangle  $\boxed{1} \boxed{2}$  is a pullback, implies that  $R_{ux} \wedge R_\varphi = 1$  and  $R_{ux} R_\varphi = R_\varphi R_{ux}$ . Since  $R_{ux} \wedge R_\varphi = 1 \leq R_x \leq R_{ux}$ , we conclude that  $R_x R_\varphi = R_\varphi R_x$  by Theorem 3.1. Equivalently, one has the equality  $x^\circ x \varphi^\circ \varphi = \varphi^\circ \varphi x^\circ x$ .

Since the vertical morphisms in  $\boxed{1}$  are split epimorphisms, then the comparison morphism  $R_x \rightarrow R_y$  is also a split epimorphism, thus the image of  $R_x$  along  $\varphi$  is  $R_y$ :  $\varphi(R_x) = R_y$  or, equivalently,  $\varphi x^\circ x \varphi^\circ = y^\circ y$ . One then has the

following equalities:

$$\begin{aligned}
\varphi x^\circ &= \varphi \varphi^\circ \varphi x^\circ x x^\circ \\
&= \varphi x^\circ x \varphi^\circ \varphi x^\circ \\
&= y^\circ y \varphi x^\circ \\
&= y^\circ f x x^\circ \\
&= y^\circ f.
\end{aligned}$$

Consequently, the arrow  $\langle \varphi, x \rangle: Z \times_V U \rightarrow Z \times_Y X$  is an isomorphism, and the square  $\square 1$  is a pullback. The square  $\square 2$  is then also a pullback, since the change-of-base functor along a regular epimorphism in a regular category reflects isomorphisms (see Proposition 2.7 of [15], for instance).  $\square$

From the proposition above we shall deduce a general result in the context of almost exact Gumm categories, which extends Lemma 1.1 in [10]. Recall that a regular category  $\mathbb{C}$  is called an *almost exact category* [16] when any regular epimorphism in  $\mathbb{C}$  is an effective descent morphism. This property can be expressed as follows [17]: for any regular epimorphism  $f: X \twoheadrightarrow Y$  and any vertical discrete fibration  $(g, h): R \rightarrow R_f$  in  $\mathbb{C}$

$$\begin{array}{ccc}
R & \begin{array}{c} \xrightarrow{r_1} \\ \xleftarrow{r_2} \end{array} & X' \\
\downarrow h & \lrcorner & \downarrow g \\
R_f & \begin{array}{c} \xrightarrow{f_1} \\ \xleftarrow{f_2} \end{array} & X \xrightarrow{f} Y
\end{array}$$

with  $R$  an equivalence relation on  $X'$ , then  $R$  is an effective equivalence relation.

**Proposition 4.2.** *Let  $\mathbb{C}$  be an almost exact Gumm category. Consider a commutative diagram*

$$(4.3) \quad \begin{array}{ccccc}
Z \times_V U & \xrightarrow{x} & X & \xrightarrow{u} & U \\
\downarrow \varphi & & \downarrow f & & \downarrow w \\
Z & \xrightarrow{y} & Y & \xrightarrow{v} & V,
\end{array}$$

such that the whole rectangle is a pullback and the left square  $\square 1$  is a pushout of regular epimorphisms. Then both squares  $\square 1$  and  $\square 2$  are pullbacks.

PROOF. Consider the following commutative diagram where  $\square 3$  gives the image factorisation of  $\langle x\varphi_1, x\varphi_2 \rangle: R_\varphi \rightarrow X \times X$ , and the morphism  $\rho$  is induced by

the equality  $wur_1 = wur_2$ :

$$\begin{array}{ccccc}
 R_\varphi & \xrightarrow{p} & R & \xrightarrow{\rho} & R_w \\
 \varphi_1 \downarrow & & \downarrow r_1 & & \downarrow w_1 \\
 \uparrow \varphi_2 & & \uparrow r_2 & & \uparrow w_2 \\
 Z \times_V U & \xrightarrow{x} & X & \xrightarrow{u} & U \\
 \varphi \downarrow & & \downarrow f & & \downarrow w \\
 Z & \xrightarrow{y} & Y & \xrightarrow{v} & V.
 \end{array}$$

Since  $\boxed{1} \boxed{2}$  is a pullback, then (any of the commutative squares)  $\boxed{3} \boxed{4}$  is also a pullback. We can apply Proposition 4.1 to  $\boxed{3} \boxed{4}$  to conclude that (any of the commutative squares)  $\boxed{3}$  and  $\boxed{4}$  are pullbacks.

Note that  $R = x(R_\varphi)$  is an equivalence relation: it is necessarily reflexive and symmetric being the image of the equivalence relation  $R_\varphi$  along a regular epimorphism  $x$ . It is also transitive: indeed, as in the proof of Proposition 4.1, the assumptions still guarantee that  $R_\varphi R_x = R_x R_\varphi$ , and this implies that

$$\begin{aligned}
 RR &= x\varphi^\circ\varphi x^\circ x\varphi^\circ\varphi x^\circ \\
 &= xx^\circ x\varphi^\circ\varphi\varphi^\circ\varphi x^\circ \\
 &= x\varphi^\circ\varphi x^\circ \\
 &= R.
 \end{aligned}$$

Since regular epimorphisms are effective for descent in  $\mathbb{C}$ , the equivalence relation  $R$  is effective, and it is then the kernel pair of its coequaliser. Moreover, the fact that the square  $\boxed{1}$  is a pushout easily implies that this coequaliser is  $f: X \twoheadrightarrow Y$ , and  $R = R_f = f^\circ f$ . To complete the proof, one applies the Barr-Kock Theorem [1] twice to conclude that  $\boxed{1}$  and  $\boxed{2}$  are pullbacks.  $\square$

**Remark 4.1.** Any efficiently regular category in the sense of [4] is almost-exact, so that the result above is true in particular in efficiently regular Gumm categories. For instance, this is the case for any category of topological Mal'tsev algebras [19], then in particular for the category of topological groups. Also any almost abelian category in the sense of [21] is almost exact (see [12]). Another example of almost exact category is provided by the category of regular epimorphisms in an exact Goursat category [16].

## 5. AN APPLICATION TO GALOIS THEORY

In this section we give an application of the results from Section 4 in Categorical Galois Theory [14, 15].

A commutative square of regular epimorphisms

$$\begin{array}{ccc} P & \xrightarrow{x} & X \\ a \downarrow & & \downarrow u \\ A & \twoheadrightarrow & U \end{array}$$

is called *right saturated* [11] when the comparison morphism  $\bar{x}: R_a \rightarrow R_u$  is also a regular epimorphism.

**Proposition 5.1.** *Let  $\mathcal{C}$  be a regular Gumm category. Consider a commutative cube*

$$(5.4) \quad \begin{array}{ccccc} P & \xrightarrow{x} & X & & \\ \varphi \uparrow & \searrow a & \uparrow f & \searrow u & \\ \sigma \downarrow & & A & \twoheadrightarrow & U \\ & & g \uparrow & & \uparrow i \\ Z & \dashrightarrow y & Y & & \\ & \searrow b & \downarrow v & & \\ & & B & \twoheadrightarrow & V \end{array}$$

of vertical split epimorphisms and regular epimorphisms. If the left and back faces are pullbacks and the top and bottom faces are right saturated, then the front and right faces are also pullbacks.

PROOF. We take the kernel pairs of  $a, b, u$  and  $v$  and the induced morphisms between them:

$$\begin{array}{ccccc} R_a & \xrightarrow{\bar{x}} & R_u & & \\ \bar{\varphi} \uparrow & \searrow a_2 & \uparrow \bar{f} & \searrow u_2 & \\ \bar{\sigma} \downarrow & & P & \xrightarrow{x} & X \\ & & \varphi \uparrow & & \uparrow f \\ R_b & \dashrightarrow \bar{y} & R_v & & Y \\ & \searrow b_2 & \downarrow v_2 & & \\ & & Z & \xrightarrow{y} & Y \end{array}$$

Note that  $\bar{x}$  and  $\bar{y}$  are regular epimorphisms since the top and bottom faces of the cube (5.4) are right saturated. The left and front faces above are pullbacks,



so that the rectangle formed by the back and right faces is a pullback. We can apply Proposition 4.1 to conclude that both the back and right faces above are pullbacks. By the Barr-Kock Theorem the right face of diagram (5.4) is then a pullback, hence so is the front face of (5.4).  $\square$

As a consequence of Proposition 5.1 we give a new proof of Theorem 4.8 of [15] stating that every central and split extension is a trivial extension for the Galois structure associated with any Birkhoff subcategory of an exact Goursat category. Let us briefly recall the main definitions, and we refer to [15] for more details.

When  $\mathbb{C}$  is an exact category and  $\mathbb{X}$  a full replete subcategory of  $\mathbb{C}$

$$(5.5) \quad \mathbb{C} \begin{array}{c} \xrightarrow{I} \\ \perp \\ \xleftarrow{\quad} \end{array} \mathbb{X},$$

one calls  $\mathbb{X}$  a *Birkhoff* subcategory of  $\mathbb{C}$  when  $\mathbb{X}$  is stable in  $\mathbb{C}$  under subobjects and regular quotients. Equivalently, all  $X$ -components  $\eta_X$  of the unit  $\eta: 1_{\mathbb{C}} \Rightarrow I$  of the adjunction are regular epimorphisms (the right adjoint is assumed to be a full inclusion and will not be mentioned explicitly), and the naturality square

$$(5.6) \quad \begin{array}{ccc} X & \xrightarrow{\eta_X} & IX \\ f \downarrow & & \downarrow If \\ Y & \xrightarrow{\eta_Y} & IY \end{array}$$

is a pushout for any regular epimorphism  $f: X \twoheadrightarrow Y$ .

A regular epimorphism  $f: X \twoheadrightarrow Y$  is called a *trivial extension* when the naturality square (5.6) is a pullback. It is called a *central extension* when it is “locally” trivial: there exists a regular epimorphism  $y: Z \twoheadrightarrow Y$  such that the pullback of  $f$  along  $y$  is a trivial extension.

**Theorem 5.2.** (Theorem 4.8 of [15]) *Let  $\mathbb{C}$  be an exact Goursat category, and  $\mathbb{X}$  a Birkhoff subcategory of  $\mathbb{C}$ . Then every central and split extension is necessarily a trivial extension.*

PROOF. Let  $f: X \twoheadrightarrow Y$  be both a central extension and a split epimorphism. By definition, there exists a regular epimorphism  $y: Z \twoheadrightarrow Y$  such that the pullback of  $f$  along  $y$  is a trivial extension. So, in the following commutative

cube

$$\begin{array}{ccccc}
 P & \xrightarrow{x} & X & & \\
 \eta_P \searrow & & \uparrow f & \eta_X \searrow & \\
 & & IP & \xrightarrow{s} & IX \\
 \sigma \downarrow & & \downarrow f & & \downarrow s \\
 Z & \xrightarrow{y} & Y & & \\
 \eta_Z \searrow & & \downarrow f & \eta_Y \searrow & \\
 & & IZ & \xrightarrow{Iy} & IY \\
 I\varphi \uparrow & & \uparrow I\sigma & & \uparrow If \\
 & & IP & \xrightarrow{I\sigma} & IX \\
 & & \downarrow I\sigma & & \downarrow Is \\
 & & IZ & \xrightarrow{I\varphi} & IY
 \end{array}$$

the back face is a pullback by construction, and the left face is a pullback by the assumption that  $\varphi$  is a trivial extension. Note that the top and bottom faces are pushouts of regular epimorphisms and are then right saturated (by Proposition 7.1 of [8]). By Proposition 5.1 we conclude that the front and right faces are pullbacks. It follows that  $f$  is a trivial extension.  $\square$

A regular epimorphism  $f: X \twoheadrightarrow Y$  is called a *normal extension* if the pullback of  $f$  along itself is a trivial extension. By definition any normal extension is central, but the converse is false in general, as various counter-examples given in [15] show. In the exact Goursat context the notions of central and normal extensions coincide:

**Corollary 5.3.** [15] *Consider a Birkhoff adjunction (5.5) where  $\mathbb{C}$  is an exact Goursat category. Then every central extension is normal.*

PROOF. This follows immediately from Theorem 5.2 and the fact that central extensions are pullback stable (this follows from the fact that adjunction is “admissible” in the sense of Categorical Galois Theory, see the Remark 5.1 here below).  $\square$

**Remark 5.1.** A Birkhoff subcategory  $\mathbb{X}$  of an exact category  $\mathbb{C}$  is called *admissible* when  $I$  preserves pullbacks of the form

$$(5.7) \quad \begin{array}{ccc}
 A & \xrightarrow{n} & U \\
 \varphi \downarrow & \lrcorner & \downarrow w \\
 B & \xrightarrow{m} & V,
 \end{array}$$

where  $w: U \twoheadrightarrow V$  is a regular epimorphism of  $\mathbb{X}$  (see Proposition 3.3 of [15]). In [20] V. Rossi proved that any Birkhoff subcategory of an almost exact Gumm category is admissible, extending a result due to G. Janelidze and G.M. Kelly

[15]. The proof of the more general Proposition 4.2 above is actually similar to the one given in [20]. In order to deduce the admissibility result from Proposition 4.2 it suffices to decompose the square (5.7) above as

$$\begin{array}{ccccc}
 A & \xrightarrow{\eta_A} & IA & \xrightarrow{In} & U \\
 \downarrow \varphi & & \downarrow I\varphi & & \downarrow w \\
 B & \xrightarrow{\eta_B} & IB & \xrightarrow{Im} & V
 \end{array}$$

to conclude that both squares are pullbacks.

#### REFERENCES

- [1] M. Barr, P.A. Grillet, D.H. van Osdol, *Exact Categories and Categories of Sheaves*, Springer Lecture Notes in Mathematics, vol. 236 (1971)
- [2] F. Borceux, D. Bourn, Mal'cev, protomodular, homological and semi-abelian categories, *Math. and its Appl.* Vol. 566, Kluwer (2004)
- [3] D. Bourn, *Fibration of points and congruence modularity*, *Algebra Universalis* 52 (2004) 403-429
- [4] D. Bourn, *Baer sums in homological categories*, *J. Algebra* 308 (2007) 414-443
- [5] D. Bourn, M. Gran, *Categorical Aspects of Modularity*. Galois Theory, Hopf Algebras and Semiabelian Categories, *Fields Instit. Commun.*, 43, Amer. Math. Soc., Providence RI (2004) 77-100
- [6] D. Bourn, M. Gran, *Normal sections and direct product decompositions* *Comm. Algebra*. 32 (2004) no. 10, 3825-3842
- [7] D. Bourn, D. Rodelo, *Comprehensive factorization and I-central extensions*, *J. Pure Appl. Algebra* 216 (2012) no. 3, 598-617
- [8] A. Carboni, G.M. Kelly, M.C. Pedicchio, *Some remarks on Maltsev and Goursat categories*, *Appl. Categ. Structures* 1 (1993), no.4, 385-421
- [9] A. Carboni, J. Lambek, M.C. Pedicchio, *Diagram chasing in Mal'cev categories*, *J. Pure Appl. Algebra* 69 (1991), no. 3, 271-284
- [10] M. Gran, *Central extensions and internal groupoids in Maltsev categories*, *J. Pure Appl. Algebra* 155 (2001), no. 2-3, 139-166
- [11] M. Gran, Z. Janelidze, D. Rodelo, A. Ursini, *Symmetry of regular diamonds, the Goursat property, and subtractivity*, *Th. Appl. Categories* 27 (2012), no. 6, 80-96
- [12] M. Gran, O. Ngaha Ngaha, *Effective descent morphisms in star-regular categories*, *Homology, Homotopy Appl.* 15 (2013), no. 2, 127-144
- [13] H.P. Gumm, *Geometrical methods in congruence modular algebras*, *Mem. Amer. Math. Soc.* 45 (1983), no. 286.
- [14] G. Janelidze, *Pure Galois theory in categories*, *J. Algebra* 132 (1990), no. 2, 270-286
- [15] G. Janelidze, G.M. Kelly, *Galois theory and a general notion of central extension*, *J. Pure Appl. Algebra* 97 (1994), no. 2, 135-161
- [16] G. Janelidze, M. Sobral, *Descent for regular epimorphisms in Barr exact Goursat categories*, *Appl. Categ. Structures* 19 (2011), no. 1, 271-276
- [17] G. Janelidze, M. Sobral, W. Tholen, *Beyond Barr exactness: Effective Descent Morphisms*, *Categorical Foundations*, 359-405, *Encyclopaedia Math. Appl.*, 97, Cambridge Univ. Press, Cambridge (2004)

- [18] G. Janelidze, W. Tholen, *Facets of descent I*, Appl. Cat. Structures 2 (1994), no. 3, 245-281
- [19] P.T. Johnstone and M.C. Pedicchio, *Remarks on continuous Mal'cev algebras*, Rend. Istit. Mat. Univ. Trieste, 25 (1993), no. 1-2, 277-297
- [20] V. Rossi, *Admissible Galois Structures and Coverings in Regular Mal'cev Categories*, Appl. Categ. Structures 14 (2006), no. 4, 291-311
- [21] W. Rump, *Almost abelian categories*, Cahiers Topologie Géom. Diiférentielle Catég. 42, (2001), no. 3, 163-225
- [22] J.D.H. Smith, *Mal'cev Varieties*, Lect. Notes Math. 554, Springer Verlag (1976)

INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE, UNIVERSITÉ CATHOLIQUE DE LOUVAIN, CHEMIN DU CYCLOTRON 2, 1348 LOUVAIN-LA-NEUVE, BELGIUM

*E-mail address:* [marino.gran@uclouvain.be](mailto:marino.gran@uclouvain.be)

CMUC, UNIVERSIDADE DE COIMBRA, 3001-501 COIMBRA, PORTUGAL  
DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS E TECNOLOGIA, UNIVERSIDADE DO ALGARVE, CAMPUS DE GAMBELAS, 8005-139 FARO, PORTUGAL

*E-mail address:* [dodelo@ualg.pt](mailto:dodelo@ualg.pt)