



Coherent pairs of linear functionals on the unit circle[☆]

A. Branquinho^a, A. Foulquié Moreno^b, F. Marcellán^{c,*}, M.N. Rebocho^{d,e}

^aCMUC, Department of Mathematics, University of Coimbra, Largo D. Dinis, 3001-454 Coimbra, Portugal

^bDepartamento de Matemática, Universidade de Aveiro, Campus de Santiago, 3810 Aveiro, Portugal

^cDepartamento de Matemáticas, Escuela Politécnica Superior, Universidad Carlos III, Avenida de la Universidad, 30, 28911 Leganés-Madrid, Spain

^dDepartamento de Matemática, Universidade da Beira Interior, 6200-001 Covilhã, Portugal

^eCMUC, University of Coimbra, 3001-454 Coimbra, Portugal

Received 9 October 2007; received in revised form 28 January 2008; accepted 11 March 2008

Communicated by Leonid Golinskii

Available online 16 March 2008

Abstract

In this paper we extend the concept of coherent pairs of measures from the real line to Jordan arcs and curves. We present a characterization of pairs of coherent measures on the unit circle: it is established that if (μ_0, μ_1) is a coherent pair of measures on the unit circle, then μ_0 is a semi-classical measure. Moreover, we obtain that the linear functional associated with μ_1 is a specific rational transformation of the linear functional corresponding to μ_0 . Some examples are given.

© 2008 Elsevier Inc. All rights reserved.

MSC: primary 42C05

Keywords: Orthogonal polynomials; Differential equations; Three-term recurrence relations; Measures on the unit circle; Hermitian functionals

[☆] The work of the first and fourth author was supported by (CMUC) Department of Mathematics, University of Coimbra. The second author would like to thank *Unidade de investigação (Matemática e Aplicações) da Universidade de Aveiro*. The work of the third author was supported by Dirección General de Investigación (Ministerio de Educación y Ciencia) of Spain under Grant *MTM 2006-13000-C03-02*. The work of M.N. Rebocho was supported by FCT, Fundação para a Ciência e Tecnologia, with Grant ref. SFRH/BD/25426/2005.

* Corresponding author. Fax: +34 916249451.

E-mail addresses: ajplb@mat.uc.pt (A. Branquinho), foulquie@ua.pt (A.F. Moreno), pacomarc@ing.uc3m.es (F. Marcellán), mneves@mat.ubi.pt (M.N. Rebocho).

1. Introduction

Let μ be a nontrivial positive Borel measure supported on a subset E of the real line. There exists a unique sequence $\{P_n\}$ of monic polynomials, with $\deg P_n = n$, such that

$$\int_E P_n(x)P_m(x) d\mu(x) = d_n^2\delta_{n,m}, \quad d_n \neq 0.$$

In this case $\{P_n\}$ is said to be the *sequence of monic orthogonal polynomials associated with μ* . It is well known that $\{P_n\}$ satisfies a three-term recurrence relation

$$xP_n(x) = P_{n+1}(x) + b_nP_n(x) + c_nP_{n-1}(x), \quad n \geq 0, \tag{1}$$

where $P_{-1}(x) = 0$ and

$$c_{n+1} = \frac{\int_E P_{n+1}^2(x) d\mu(x)}{\int_E P_n^2(x) d\mu(x)}, \quad b_n = \frac{\int_E xP_n^2(x) d\mu(x)}{\int_E P_n^2(x) d\mu(x)}, \quad n \geq 0.$$

On the other hand, if (1) holds with $c_n > 0$, there exists the sequence of monic polynomials defined by (1) orthogonal with respect to the measure μ .

Let (μ_0, μ_1) be a pair of nontrivial positive Borel measures supported on subsets E_0 and E_1 of the real line. We introduce an inner product in the linear space \mathbb{P} of polynomials with real coefficients

$$(p, q) = \int_{E_0} p(x)q(x) d\mu_0(x) + \lambda \int_{E_1} p'(x)q'(x) d\mu_1(x), \tag{2}$$

where $p, q \in \mathbb{P}$ and $\lambda \geq 0$.

This kind of inner products define a sequence $\{Q_n(\cdot, \lambda)\}$ of monic polynomials that is orthogonal with respect to (2). It can be constructed using the standard Gram–Schmidt process. But these polynomials do not satisfy a three-term recurrence relation as (1). If $\{P_n\}$ and $\{R_n\}$ denote, respectively, the sequences of monic polynomials orthogonal with respect to μ_0, μ_1 , then Iserles et al. introduced the concept of coherent pairs of measures in [6].

A pair of nontrivial Borel measures (μ_0, μ_1) supported on subsets of the real line is said to be *coherent* if the corresponding sequences of monic orthogonal polynomials satisfy

$$R_n(x) = \frac{P'_{n+1}}{n+1}(x) + \alpha_n \frac{P'_n}{n}(x), \quad \alpha_n \neq 0, \quad n = 1, 2, \dots \tag{3}$$

From here, a relation between $\{P_n\}$ and $\{Q_n(\cdot, \lambda)\}$ follows:

$$P_n(x) + \frac{n}{n-1}\alpha_{n-1}P_{n-1}(x) = Q_n(x, \lambda) + \beta_{n-1}(\lambda)Q_{n-1}(x, \lambda),$$

where $\beta_{n-1}(\lambda) = \frac{\gamma_{n-2}(\lambda)}{\gamma_{n-1}(\lambda)}$, γ_n is a polynomial of degree n in the variable λ , and $\{\gamma_n\}$ satisfies a three-term recurrence relation.

In [6] the authors ask about the description of all coherent pairs of measures. The answer was given by Meijer [8], where he proves that at least one of the measures must be a classical one (Laguerre or Jacobi). In particular, when the support is a compact subset of the real axis,

the following cases appear:

(a) $d\mu_0 = (1 - x)^\alpha(1 + x)^\beta dx, \quad \alpha, \beta > -1,$

$$d\mu_1 = \frac{(1 - x)^{\alpha+1}(1 + x)^{\beta+1}}{|x - \xi|} dx + M\delta(x - \xi), \quad |\xi| \geq 1, \quad M \geq 0,$$

(b) $d\mu_0 = (1 - x)^\alpha(1 + x)^\beta|x - \xi| dx, \quad d\mu_1 = (1 - x)^{\alpha+1}(1 + x)^{\beta+1} dx, \quad \alpha, \beta > -1,$

(c) $d\mu_0 = (1 - x)^\alpha dx + M\delta(x + 1), \quad d\mu_1 = (1 - x)^{\alpha+1} dx, \quad \alpha > -1,$

(d) $d\mu_0 = (1 + x)^\beta dx + M\delta(x - 1), \quad d\mu_1 = (1 + x)^{\beta+1} dx, \quad \beta > -1.$

The aim of this contribution is the analysis of the concept of coherent pairs of measures supported on compact subsets of the complex plane. In particular, we will focus our attention when the support is the unit circle.

The structure of the manuscript is as follows. In Section 2 we define coherent pairs of measures supported on Jordan arcs or curves using the connection between the corresponding sequences of orthogonal polynomials as in (3). As a consequence, the relation between these sequences and the sequence of monic polynomials orthogonal with respect to the Sobolev inner product associated with the pair of measures (μ_0, μ_1) is deduced. In Section 3 we present the basic results concerning Hermitian orthogonality on the unit circle which will be used in the forthcoming sections. We give a sufficient condition for a sequence of orthogonal polynomials on the unit circle (OPUC) satisfying a first order structure relation to be semi-classical (see Theorem 3). This result is an extension to the result deduced by Branquinho and Rebocho [3]. In Section 4 we present a characterization of pairs of coherent measures on the unit circle; we prove that if (μ_0, μ_1) is a coherent pair of measures on the unit circle (μ_0, μ_1) then μ_0 is a semi-classical measure and the linear functional associated with μ_1 is a specific rational transformation of the linear functional corresponding to μ_0 (see, for example, [2]). Finally, in Section 5, we study the companion coherent measure associated with the Bernstein–Szegő measure supported on the unit circle.

2. Coherent pairs of measures supported on Jordan arcs and curves

Let μ_0, μ_1 be positive Borel measures on E_0, E_1 , respectively, which are Jordan curves or arcs. For $\lambda \in \mathbb{R}^+$, consider the inner product

$$\langle f, g \rangle_S = \langle f, g \rangle_0 + \lambda \langle f', g' \rangle_1 \quad \text{where } \langle f, g \rangle_k = \int_{E_k} f(\xi) \overline{g(\xi)} d\mu_k(\xi), \quad k = 0, 1.$$

Let us denote by $\{Q_n(\cdot; \lambda)\}, \{P_n\}, \{R_n\}$, the sequences of monic polynomials orthogonal with respect to $\langle \cdot, \cdot \rangle_S, \langle \cdot, \cdot \rangle_0, \langle \cdot, \cdot \rangle_1$, respectively.

We also denote

$$S_{m,n} := \langle z^m, z^n \rangle_S = c_{m,n}^0 + \lambda m n c_{m-1,n-1}^1, \quad m, n \in \mathbb{N},$$

where $\{c_{m,n}^k\}_{n \in \mathbb{N}}$ are the moments with respect to the measures μ_k for $k = 0, 1$, respectively.

Taking into account this expression, we obtain the following representation in a determinantal form for the polynomials Q_n :

$$Q_n(z; \lambda) = \frac{\begin{vmatrix} c_{0,0}^0 & c_{1,0}^0 & \cdots & c_{n,0}^0 \\ c_{0,1}^0 & c_{1,1}^0 + \lambda c_{0,0}^1 & \cdots & c_{n,1}^0 + \lambda n c_{n-1,0}^1 \\ \vdots & \vdots & \ddots & \vdots \\ c_{0,n-1}^0 & c_{1,n-1}^0 + \lambda(n-1)c_{0,n-2}^1 & \cdots & c_{n,n-1}^0 + \lambda n(n-1)c_{n-1,n-2}^1 \\ 1 & z & \cdots & z^n \end{vmatrix}}{\begin{vmatrix} c_{0,0}^0 & c_{1,0}^0 & \cdots & c_{n-1,0}^0 \\ c_{0,1}^0 & c_{1,1}^0 + \lambda c_{0,0}^1 & \cdots & c_{n-1,1}^0 + \lambda(n-1)c_{n-2,0}^1 \\ \vdots & \vdots & \ddots & \vdots \\ c_{0,n-1}^0 & c_{1,n-1}^0 + \lambda(n-1)c_{0,n-2}^1 & \cdots & c_{n-1,n-1}^0 + \lambda(n-1)^2 c_{n-2,n-2}^1 \end{vmatrix}}.$$

Since the coefficients of the above polynomial are rational functions in λ , when λ tends to infinity we get the sequence of limit polynomials, $\{S_n\}$. It is straightforward to prove that the polynomial S_n satisfies

$$\langle S_n, 1 \rangle_0 = 0, \quad n \geq 1, \quad \langle S'_n, z^k \rangle_1 = 0, \quad 0 \leq k \leq n - 2, \quad n \geq 2,$$

and so $S'_n(z) = nR_{n-1}(z)$, $n = 1, 2, \dots$. See [4] for an analysis of such limit polynomials when a pair of measures supported on the real line is considered.

Therefore, using the same arguments as in [6], we get the Fourier expansions of S_n with respect to the sequences $\{P_n\}$ and $\{Q_n\}$, i.e.

$$S_n(z) = \sum_{k=1}^n a_{n-1,k} P_k(z), \quad S_n(z) = Q_n(z; \lambda) + \sum_{j=0}^{n-1} \beta_{n,j}(\lambda) Q_j(z; \lambda). \tag{4}$$

From this we do not get more information, but if in (4) we assume that $a_{n-1,k} = 0$ for $k < n - s$ (with s a fixed nonnegative integer number), it follows that $\beta_{n,j}(\lambda) = 0$ for $j < n - s$. Thus, for $n \geq s$,

$$\sum_{k=n-s}^n a_{n-1,k} P_k(z) = \sum_{j=n-s}^n \beta_{n,j}(\lambda) Q_j(z; \lambda). \tag{5}$$

Conversely, notice that if (5) holds, and $a_{n-1,n-s} \neq 0$, $\beta_{n,n-s}(\lambda) \neq 0$, then

$$\int_{E_1} \sum_{j=n-s}^n a_{n-1,j} P'_j(z) p'(z) d\mu_1 = 0, \quad p \in \mathbb{P}_{n-s-1}.$$

From this the following relation holds:

$$\sum_{j=n-s}^n a_{n-1,j} P'_j(z) = \sum_{j=n-s-1}^{n-1} b_{n,j} R_j(z).$$

Therefore the following problem arises: *To describe the measures μ_0, μ_1 such that the corresponding sequences of monic orthogonal polynomials $\{P_n\}$ and $\{R_n\}$ are related by*

$$R_{n-1}(z) = \frac{P'_n(z)}{n} + \alpha_{n-1} \frac{P'_{n-1}(z)}{n-1}, \quad \alpha_{n-1} \neq 0, \quad n = 2, 3, \dots \tag{6}$$

From now on, for a sake of simplicity, we write β_n instead of $\beta_{n,n}$, as well as a_n instead of $a_{n,n}$.

For a coherent pair of measures we get some extra information about the sequence $(\beta_n(\lambda))$. Indeed,

$$P_n(z) + a_{n-1}P_{n-1}(z) = Q_n(z; \lambda) + \beta_{n-1}(\lambda)Q_{n-1}(z; \lambda), \tag{7}$$

where for $n = 2, 3, \dots$

$$a_{n-1} = \frac{n}{n-1} \alpha_{n-1},$$

$$\beta_{n-1}(\lambda) = a_{n-1} \frac{\langle P_{n-1}, Q_{n-1}(\cdot; \lambda) \rangle_0}{\langle Q_{n-1}(\cdot; \lambda), Q_{n-1}(\cdot; \lambda) \rangle_S} = a_{n-1} \frac{\|P_{n-1}\|_0^2}{\|Q_{n-1}(\cdot; \lambda)\|_S^2}. \tag{8}$$

Therefore, taking into account (6) and (7), after some calculations we get

$$\begin{aligned} \|Q_{n-1}(\cdot; \lambda)\|_S^2 &= \langle Q_{n-1}(\cdot; \lambda), P_{n-1} \rangle_S \\ &= \|P_{n-1}\|_0^2 + \lambda(n-1)^2 \|R_{n-2}\|_1^2 + \bar{a}_{n-2} [a_{n-2} - \beta_{n-2}(\lambda)] \|P_{n-2}\|_0^2. \end{aligned}$$

Now, substituting in (8), and using the preceding notation for $n = 3, 4, \dots$, we deduce that

$$\beta_{n-1}(\lambda) = \frac{A_n}{B_n - \beta_{n-2}(\lambda)}, \tag{9}$$

where

$$A_n = \frac{a_{n-1} \|P_{n-1}\|_0^2}{\bar{a}_{n-2} \|P_{n-2}\|_0^2}, \quad B_n = a_{n-2} + \frac{\|P_{n-1}\|_0^2 + \lambda(n-1)^2 \|R_{n-2}\|_1^2}{\bar{a}_{n-2} \|P_{n-2}\|_0^2},$$

with $\beta_1(\lambda) = \frac{\|P_1\|_0^2 a_1}{\lambda \|R_0\|_1^2 + \|P_1\|_0^2}$.

Notice that B_n is a polynomial of degree one in λ . In this way, once we obtain the coherent pairs we can deduce a representation for $\beta_{n-1}(\lambda)$, which are rational functions of λ and, eventually, from (7) we get an explicit expression for $Q_n(\cdot; \lambda)$ in terms of $\{P_n\}$.

Theorem 1. *The sequence $(\beta_n(\lambda))$ is given by*

$$\beta_{n-1}(\lambda) = \frac{\gamma_{n-2}(\lambda)}{\gamma_{n-1}(\lambda)}, \quad n = 2, 3, \dots, \tag{10}$$

where $\{\gamma_n\}$ is a sequence of orthogonal polynomials associated with a positive Borel measure supported on \mathbb{R} .

Proof. Taking into account β_1 is a rational function in λ such that the degree of the numerator is zero and the degree of the denominator is one, by induction we get (10) where γ_n is a polynomial of degree n . Moreover, from (9), we get

$$\gamma_n(\lambda) = \frac{B_{n+1}}{A_{n+1}} \gamma_{n-1}(\lambda) - \frac{1}{A_{n+1}} \gamma_{n-2}(\lambda). \tag{11}$$

Taking into account that B_n is a polynomial of degree one in λ , we get that $\{\gamma_n\}$ is a sequence of polynomials orthogonal with respect to a linear functional. This is a straightforward consequence of the Favard Theorem, since they satisfy a three-term recurrence relation (see [5]).

Indeed, if $\gamma_n(\lambda) = s_n \lambda^n + \text{lower degree terms}$, then (11) becomes

$$s_n \tilde{\gamma}_n(\lambda) = \frac{B_{n+1}}{A_{n+1}} s_{n-1} \tilde{\gamma}_{n-1}(\lambda) - \frac{s_{n-2}}{A_{n+1}} \tilde{\gamma}_{n-2}(\lambda),$$

or, equivalently, for $n = 2, 3, \dots$

$$\tilde{\gamma}_n(\lambda) = (\lambda + c_{n-1}) \tilde{\gamma}_{n-1}(\lambda) - d_{n-1} \tilde{\gamma}_{n-2}(\lambda),$$

where

$$c_{n-1} = \frac{|a_{n-1}|^2 \|P_{n-1}\|_0^2 + \|P_n\|_0^2}{n^2 \|R_{n-1}\|_1^2}, \quad d_{n-1} = \frac{\|P_{n-1}\|_0^4 |a_{n-1}|^2}{n^2 (n-1)^2 \|R_{n-1}\|_1^2 \|R_{n-2}\|_1^2} > 0,$$

and initial conditions $\tilde{\gamma}_0(\lambda) = 1, \tilde{\gamma}_1(\lambda) = \lambda + \|P_1\|_0^2 / \|R_0\|_1^2$. Notice that, according to the Favard Theorem, $\{\tilde{\gamma}_n\}$ is a sequence of monic polynomials orthogonal with respect to a finite positive Borel measure supported on \mathbb{R} . \square

3. Quasi-orthogonality on the unit circle

Let $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$, and $\Lambda = \text{span}\{z^k : k \in \mathbb{Z}\}$, be the linear space of Laurent polynomials with complex coefficients. Given a linear functional $u : \Lambda \rightarrow \mathbb{C}$, and the sequence of moments $(c_n)_{n \in \mathbb{Z}}$ of $u, c_n = \langle u, \xi^n \rangle, n \in \mathbb{Z}, c_0 = 1$, define the minors of the Toeplitz matrix $\Delta = (c_{k-j})$, by

$$\Delta_k = \begin{vmatrix} c_0 & \cdots & c_k \\ \vdots & \ddots & \vdots \\ c_{-k} & \cdots & c_0 \end{vmatrix}, \quad \Delta_0 = c_0, \quad \Delta_{-1} = 1, \quad k \in \mathbb{N}.$$

u is said to be *Hermitian* if $c_{-n} = \bar{c}_n, \forall n \in \mathbb{N}$, and *quasi-definite* (respectively, *positive definite*) if $\Delta_n \neq 0$ (respectively, $\Delta_n > 0$), $\forall n \in \mathbb{N}$. We will denote by \mathcal{H} the set of Hermitian linear functionals defined on Λ .

In the positive-definite case, u has an integral representation given in terms of a nontrivial probability measure μ with infinite support on the unit circle \mathbb{T} ,

$$\langle u, e^{in\theta} \rangle = \frac{1}{2\pi} \int_0^{2\pi} e^{in\theta} d\mu(\theta), \quad n \in \mathbb{Z}.$$

The corresponding sequence of orthogonal polynomials, called OPUC, is then defined by

$$\frac{1}{2\pi} \int_0^{2\pi} P_n(e^{i\theta}) \bar{P}_m(e^{-i\theta}) d\mu(\theta) = e_n \delta_{n,m}, \quad e_n > 0, \quad n, m = 0, 1, \dots$$

If $P_n(z) = z^n + \text{lower degree terms}$, $\{P_n\}$ will be called a *sequence of monic orthogonal polynomials*, and we will denote it by MOPS. It is well known that MOPS on the unit circle satisfy the following recurrence relations, known as *Szegő recurrence relations*, for $n \geq 1$:

$$P_n(z) = z P_{n-1}(z) + a_n P_{n-1}^*(z), \quad P_n^*(z) = P_{n-1}^*(z) + \bar{a}_n z P_{n-1}(z),$$

with $a_n = P_n(0)$, $P_0(z) = 1$, and $P_n^*(z) = z^n \bar{P}_n(1/z)$, $n = \deg(P_n)$. In the literature of OPUC, the polynomials $\{P_n^*\}$ are called either reversed or reciprocal polynomials (see [9]).

$\{P_n^*\}$ satisfies, for $n \in \mathbb{N}$,

$$\langle u, P_n^*(z)z^{-k} \rangle = 0, \quad k = 1, \dots, n, \quad \langle u, P_n^*(z) \rangle = e_n. \tag{12}$$

The following relation holds (see [7]):

$$(P_n')^*(z) = nP_n^*(z) - z(P_n^*)'(z), \quad n \geq 1. \tag{13}$$

For $u \in \mathcal{H}$ and $A \in \mathbb{P}$, we define

$$\begin{aligned} \langle Au, f \rangle &= \langle u, A(z)f(z) \rangle, \quad f \in \Lambda, \\ \langle (A + \bar{A})u, f \rangle &= \langle u, (A(z) + \bar{A}(1/z))f(z) \rangle, \quad f \in \Lambda. \end{aligned}$$

Notice that $(A + \bar{A})u$ is a Hermitian linear functional. We will use the notation

$$u^A = (A(z) + \bar{A}(1/z))u.$$

Definition 1 (cf. Alfaro and Moral [1]). Let $v \in \mathcal{H}$, $p \in \mathbb{N}$, and let $\{P_n\}$ be a sequence of monic polynomials. $\{P_n\}$ is said to be \mathbb{T} -quasi-orthogonal of order p with respect to v if

- (i) $\langle v, P_n(z)z^{-k} \rangle = 0$, for every k with $p \leq k \leq n - p - 1$ and for every $n \geq 2p + 1$.
- (ii) There exists $n_0 \geq 2p$ such that $\langle v, P_{n_0}(z)z^{-n_0+p} \rangle \neq 0$.

Theorem 2 (cf. Alfaro and Moral [1]). Let $u \in \mathcal{H}$ be quasi-definite and let $\{P_n\}$ be the MOPS with respect to u . Then $\{P_n\}$ is \mathbb{T} -quasi-orthogonal of order p with respect to $v \in \mathcal{H} - \{0\}$ if and only if there exists only one polynomial B ($B \neq 0$) with $\deg(B) = p$, such that $v = u^B$.

Taking into account Theorem 4.1 of [1] we give the following definition.

Definition 2. Let $u \in \mathcal{H}$ be quasi-definite and let $\{P_n\}$ be the MOPS associated with u . u is said to be *semi-classical* if there exists $\hat{u} \in \mathcal{H} - \{0\}$ such that the sequence $\{\tilde{P}_n\}$ given by $\tilde{P}_n(z) = \frac{1}{n}zP_n'(z)$, $n \geq 1$, $\tilde{P}_0(z) = 1$, is \mathbb{T} -quasi-orthogonal with respect to \hat{u} . In such a situation $\{P_n\}$ is said to be a *semi-classical sequence* of orthogonal polynomials.

In the sequel we define $f_n(z) = P_n(z)/P_n^*(z)$, $\forall n \in \mathbb{N}$, and we study the conditions in order to $\{f_n\}$ satisfies a Riccati differential equation. This result will be useful to the following theorem. Using the Szegő recurrence relations we get

$$zf_n(z) = \frac{f_{n+1}(z) - a_{n+1}}{1 - \bar{a}_{n+1}f_{n+1}(z)}, \quad n = 1, \dots \tag{14}$$

Lemma 1. Let $\{P_n\}$ be a sequence of monic orthogonal polynomials on the unit circle and $\{P_n^*\}$ the sequence of reversed polynomials. If $\{f_n\}$ satisfies a Riccati differential equation with bounded degree polynomial coefficients, i.e.,

$$A_n(z)f_n'(z) = B_n(z)f_n^2(z) + C_n(z)f_n(z) + E_n(z), \quad \forall n \in \mathbb{N} \tag{15}$$

then, for every $n \in \mathbb{N}$, the following relations hold:

$$A_{n+1} = A_n, \tag{16}$$

$$zB_{n+1} = \lambda_n^{-1} \left\{ B_n - \bar{a}_{n+1}(zC_n + A_n) + \bar{a}_{n+1}^2 z^2 E_n \right\}, \tag{17}$$

$$zC_{n+1} = \lambda_n^{-1} \left\{ (-2a_{n+1}B_n + (zC_n + A_n)(1 + |a_{n+1}|^2) - 2\bar{a}_{n+1}z^2 E_n) \right\}, \tag{18}$$

$$zE_{n+1} = \lambda_n^{-1} \left\{ a_{n+1}^2 B_n - a_{n+1}(zC_n + A_n) + z^2 E_n \right\}, \tag{19}$$

with $\lambda_n = (1 - |a_{n+1}|^2)$.

Proof. If f_n satisfies (15), then

$$zA_n(zf_n)' = B_n(zf_n)^2 + (zC_n + A_n)zf_n + z^2 E_n.$$

Using (14) in previous equation we get

$$zA_n \left(\frac{f_{n+1} - a_{n+1}}{1 - \bar{a}_{n+1}f_{n+1}} \right)' = B_n \left(\frac{f_{n+1} - a_{n+1}}{1 - \bar{a}_{n+1}f_{n+1}} \right)^2 + (zC_n + A_n) \left(\frac{f_{n+1} - a_{n+1}}{1 - \bar{a}_{n+1}f_{n+1}} \right) + z^2 E_n.$$

Since

$$\left(\frac{f_{n+1} - a_{n+1}}{1 - \bar{a}_{n+1}f_{n+1}} \right)' = \frac{\lambda_n f'_{n+1}}{(1 - \bar{a}_{n+1}f_{n+1})^2} \quad \text{with } \lambda_n = 1 - |a_{n+1}|^2,$$

from the previous equations we get

$$\begin{aligned} zA_n \frac{\lambda_n f'_{n+1}}{(1 - \bar{a}_{n+1}f_{n+1})^2} &= B_n \left(\frac{f_{n+1}^2 + a_{n+1}^2 - 2a_{n+1}f_{n+1}}{(1 - \bar{a}_{n+1}f_{n+1})^2} \right) \\ &\quad + (zC_n + A_n) \left(\frac{f_{n+1} - a_{n+1}}{1 - \bar{a}_{n+1}f_{n+1}} \right) + z^2 E_n, \end{aligned}$$

as well as

$$\begin{aligned} \lambda_n zA_n f'_{n+1} &= \left\{ B_n - \bar{a}_{n+1}(zC_n + A_n) + \bar{a}_{n+1}^2 z^2 E_n \right\} f_{n+1}^2 \\ &\quad + \left\{ (-2a_{n+1}B_n + (zC_n + A_n)(1 + |a_{n+1}|^2) - 2\bar{a}_{n+1}z^2 E_n) \right\} f_{n+1} \\ &\quad + a_{n+1}^2 B_n - a_{n+1}(zC_n + A_n) + z^2 E_n. \end{aligned}$$

If we divide by $\lambda_n = (1 - |a_{n+1}|^2)$ then

$$\begin{aligned} zA_n f'_{n+1} &= \lambda_n^{-1} \left\{ B_n - \bar{a}_{n+1}(zC_n + A_n) + \bar{a}_{n+1}^2 z^2 E_n \right\} f_{n+1}^2 \\ &\quad + \lambda_n^{-1} \left\{ (-2a_{n+1}B_n + (zC_n + A_n)(1 + |a_{n+1}|^2) - 2\bar{a}_{n+1}z^2 E_n) \right\} f_{n+1} \\ &\quad + \lambda_n^{-1} \left\{ a_{n+1}^2 B_n - a_{n+1}(zC_n + A_n) + z^2 E_n \right\}. \end{aligned}$$

Now, comparing previous equation with (15) to $n + 1$ and multiplied by z , i.e., with

$$zA_{n+1}f'_{n+1} = zB_{n+1}f^2_{n+1} + zC_{n+1}f_{n+1} + zE_{n+1},$$

we get (16)–(19). \square

Theorem 3. Let $\{P_n\}$ be an MOPS the unit circle and $\{P_n^*\}$ be the sequence of reversed polynomials. If $\{P_n\}$ satisfies a structure relation with bounded degree polynomials, $n \geq 1$,

$$z\Pi_n(z)P'_n(z) = G_n(z)P_n(z) + H_n(z)P_n^*(z), \tag{20}$$

$$z\Pi_n(z)(P_n^*)'(z) = S_n(z)P_n(z) + T_n(z)P_n^*(z) \tag{21}$$

then Π_n does not depend on n .

Let $p = \max\{\deg(G_n), \deg(H_n)+1, \deg(S_n), \deg(\Pi_1-T_n)\}$, $\forall n \in \mathbb{N}$. If there exists $n_0 \geq 2p$ such that $\deg(\Pi_1 - T_{n_0}) = p$, then $\{P_n\}$ is semi-classical.

Proof. If we multiply (20) by P_n^* , (21) by P_n , and divide the resulting equations by $(P_n^*)^2$, we get, after subtracting the corresponding equations,

$$\begin{aligned} z\Pi_n \left(\frac{P'_n P_n^* - P_n (P_n^*)'}{(P_n^*)^2} \right) &= \frac{(G_n - T_n)P_n P_n^* + H_n (P_n^*)^2 - S_n (P_n)^2}{(P_n^*)^2} \\ \Leftrightarrow z\Pi_n \left(\frac{P_n}{P_n^*} \right)' &= -S_n \left(\frac{P_n}{P_n^*} \right)^2 + (G_n - T_n) \frac{P_n}{P_n^*} + H_n. \end{aligned}$$

Thus,

$$z\Pi_n f'_n = -S_n f_n^2 + (G_n - T_n) f_n + H_n.$$

From previous lemma, $\Pi_n = \Pi_{n-1}$, $\forall n \in \mathbb{N}$. Thus, $\Pi_n = \Pi_1$, $\forall n \in \mathbb{N}$.

Let us write (20) and (21) in the form

$$A \frac{zP'_n}{n} = \tilde{G}_n P_n + \tilde{H}_n P_n^*, \tag{22}$$

$$A \frac{z(P_n^*)'}{n} = \tilde{S}_n P_n + \tilde{T}_n P_n^*, \quad n \geq 1, \tag{23}$$

with $A = \Pi_1$, $\tilde{G}_n = G_n/n$, $\tilde{H}_n = H_n/n$, $\tilde{S}_n = S_n/n$, $\tilde{T}_n = T_n/n$. Furthermore, if we use (13) in (23) then

$$A \left(\frac{zP'_n}{n} \right)^* = -\tilde{S}_n P_n + (A - \tilde{T}_n) P_n^*. \tag{24}$$

On the other hand, from the Hermitian character of u , we have

$$\left\langle u^A, \frac{zP'_n}{n} z^{-k} \right\rangle = \left\langle u, A \frac{zP'_n}{n} z^{-k} \right\rangle + \overline{\left\langle u, A \left(\frac{zP'_n}{n} \right)^* z^{k-n} \right\rangle}.$$

Using (22) and (24) in previous equation we get

$$\begin{aligned} \left\langle u^A, \frac{zP'_n}{n} z^{-k} \right\rangle &= \langle u, \tilde{G}_n P_n z^{-k} \rangle + \langle u, \tilde{H}_n P_n^* z^{-k} \rangle \\ &\quad - \overline{\langle u, \tilde{S}_n P_n z^{k-n} \rangle} + \overline{\langle u, (A - \tilde{T}_n) P_n^* z^{k-n} \rangle}. \end{aligned} \tag{25}$$

Since

$$\begin{aligned} \langle u, \tilde{G}_n P_n z^{-k} \rangle &= 0, \quad k = \deg(\tilde{G}_n), \dots, n - 1, \\ \langle u, \tilde{H}_n P_n^* z^{-k} \rangle &= 0, \quad k = \deg(\tilde{H}_n) + 1, \dots, n, \\ \langle u, \tilde{S}_n P_n z^{k-n} \rangle &= 0, \quad k = 1, \dots, n - \deg(\tilde{S}_n), \\ \langle u, (A - \tilde{T}_n) P_n^* z^{k-n} \rangle &= 0, \quad k = 0, \dots, n - \deg(A - \tilde{T}_n) - 1 \end{aligned}$$

then, with $p = \max\{\deg(\tilde{G}_n), \deg(\tilde{H}_n) + 1, \deg(\tilde{S}_n), \deg(A - \tilde{T}_n)\}$, $\forall n \in \mathbb{N}$, it follows that

$$\left\langle u^A, \frac{z P'_n}{n} z^{-k} \right\rangle = 0 \quad \text{for every } p \leq k \leq n - p - 1 \text{ and for every } n \geq 2p + 1.$$

Next we show that condition (ii) of Definition 2,

$$\exists n_0 \geq 2p : \left\langle u^A, \frac{z P'_{n_0}}{n_0} z^{-n_0+p} \right\rangle \neq 0,$$

holds for $n_0 \geq 2p$ if and only if $\deg(A - \tilde{T}_{n_0}) = p$.

From (25)

$$\begin{aligned} \left\langle u^A, \frac{z P'_{n_0}}{n_0} z^{-n_0+p} \right\rangle &= \langle u, \tilde{G}_{n_0} P_{n_0} z^{-n_0+p} \rangle + \langle u, \tilde{H}_{n_0} P_{n_0}^* z^{-n_0+p} \rangle \\ &\quad - \overline{\langle u, \tilde{S}_{n_0} P_{n_0} z^{-p} \rangle} + \overline{\langle u, (A - \tilde{T}_{n_0}) P_{n_0}^* z^{-p} \rangle}. \end{aligned} \tag{26}$$

Since $\deg(\tilde{G}_n) \leq p$, $\deg(\tilde{H}_n) \leq p - 1$, $\deg(\tilde{S}_n) \leq p$, $\forall n \in \mathbb{N}$, and $n_0 - p \geq p$, then

$$\langle u, \tilde{G}_{n_0} P_{n_0} z^{-n_0+p} \rangle = \langle u, \tilde{H}_{n_0} P_{n_0}^* z^{-n_0+p} \rangle = \overline{\langle u, \tilde{S}_{n_0} P_{n_0} z^{-p} \rangle} = 0.$$

Therefore, (26) is equivalent to

$$\left\langle u^A, \frac{z P'_{n_0}}{n_0} z^{-n_0+p} \right\rangle = \overline{\langle u, (A - \tilde{T}_{n_0}) P_{n_0}^* z^{-p} \rangle}.$$

Taking into account the orthogonality relations (12) and $\deg(A - T_n) \leq p$, we get

$$\langle u, (A - \tilde{T}_{n_0}) P_{n_0}^* z^{-p} \rangle \neq 0 \Leftrightarrow \deg(A - \tilde{T}_{n_0}) = p.$$

Thus,

$$\left\langle u^A, \frac{z P'_{n_0}}{n_0} z^{-n_0+p} \right\rangle \neq 0 \Leftrightarrow \deg(A - \tilde{T}_{n_0}) = p.$$

Therefore, if there exists $n_0 \geq 2p$ such that $\deg(A - \tilde{T}_{n_0}) = p$, then the sequence $\{\frac{1}{n} z P'_n\}$ is \mathbb{T} -quasi-orthogonal of order p with respect to the Hermitian functional u^A and we conclude that $\{P_n\}$ is semi-classical. \square

4. Characterization theorem

In the sequel we will use the vectors defined by

$$\psi_n(z) = [P_n(z) \ P_n^*(z)]^T, \quad \vartheta_n(z) = [R_n(z) \ R_n^*(z)]^T, \quad n \in \mathbb{N}.$$

We will use the Szegő recurrence relations in the matrix form for $\{\psi_n\}$,

$$\psi_n(z) = \mathcal{A}_n(z)\psi_{n-1}(z), \quad \mathcal{A}_n(z) = \begin{bmatrix} z & a_n \\ \bar{a}_n z & 1 \end{bmatrix}, \quad n \in \mathbb{N}, \quad a_n = P_n(0), \tag{27}$$

and for $\{\vartheta_n\}$,

$$\vartheta_n(z) = \mathcal{B}_n(z)\vartheta_{n-1}(z), \quad \mathcal{B}_n(z) = \begin{bmatrix} z & b_n \\ \bar{b}_n z & 1 \end{bmatrix}, \quad n \in \mathbb{N}, \quad b_n = R_n(0). \tag{28}$$

We will write $X^{(i,j)}$ to denote the entry (i, j) of a matrix X , $i, j = 1, 2$.

Theorem 4. *Let (u, v) be a coherent pair of Hermitian linear functionals on the unit circle and $\{P_n\}, \{R_n\}$ the corresponding MOPS. Then, there exist $A \in \mathbb{P}$ and matrices $\mathcal{K}_n, \mathcal{M}_n$ of order two whose elements are bounded degree polynomials such that, for $n \geq 1$,*

$$zA(z)\psi'_n(z) = \mathcal{K}_n(z)\psi_n(z), \tag{29}$$

$$zA(z)\vartheta_n(z) = \mathcal{M}_n(z)\psi_n(z). \tag{30}$$

Moreover,

- (a) $\{P_n\}$ is semi-classical;
- (b) $\{R_n\}$ is quasi-orthogonal of order p ($p \leq 6$) with respect to the functional u^{zA} . Thus, there exists a unique polynomial B of degree p such that $u^{zA} = v^B$.

Proof. From

$$R_n = \frac{P'_{n+1}}{n+1} + \alpha_n \frac{P'_n}{n} \tag{31}$$

we get

$$R_n^* = \frac{(P'_{n+1})^*}{n+1} + \bar{\alpha}_n z \frac{(P'_n)^*}{n}.$$

Using (13), last equation is equivalent to

$$R_n^* = P_{n+1}^* + \bar{\alpha}_n z P_n^* - z \frac{(P_{n+1}^*)'}{n+1} - \bar{\alpha}_n z^2 \frac{(P_n^*)'}{n}. \tag{32}$$

If we write (31) and (32) in a matrix form and use (27), we obtain

$$\vartheta_n = \mathcal{S}_n \psi_n + \mathcal{T}_n \psi'_n, \quad n \geq 1, \tag{33}$$

with

$$\mathcal{S}_n = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \mathcal{A}_{n+1} + \begin{bmatrix} 0 & 0 \\ 0 & \bar{\alpha}_n z \end{bmatrix} + \begin{bmatrix} 1/(n+1) & 0 \\ 0 & -z/(n+1) \end{bmatrix} \begin{bmatrix} 1 & 0 \\ \bar{a}_{n+1} & 0 \end{bmatrix},$$

$$\mathcal{T}_n = \begin{bmatrix} 1/(n+1) & 0 \\ 0 & -z/(n+1) \end{bmatrix} \mathcal{A}_{n+1} + \begin{bmatrix} \alpha_n/n & 0 \\ 0 & -\bar{\alpha}_n z^2/n \end{bmatrix}.$$

Using (33) for $n + 1$ and the recurrence relations (27) and (28), we get

$$\mathcal{H}_n \psi'_n = \widetilde{\mathcal{M}}_n \psi_n, \tag{34}$$

where the matrices \mathcal{H}_n and $\widetilde{\mathcal{M}}_n$ are given by

$$\mathcal{H}_n = \mathcal{B}_{n+1} \mathcal{T}_n - \mathcal{T}_{n+1} \mathcal{A}_{n+1}, \quad \widetilde{\mathcal{M}}_n = \mathcal{S}_{n+1} \mathcal{A}_{n+1} + \mathcal{T}_{n+1} \begin{bmatrix} 1 & 0 \\ \bar{a}_{n+1} & 0 \end{bmatrix} - \mathcal{B}_{n+1} \mathcal{S}_n.$$

Now, if we multiply (34) by the adjoint matrix of \mathcal{H}_n , $\text{adj } \mathcal{H}_n$, we get

$$h_n \psi'_n = \mathcal{K}_n \psi_n,$$

where $h_n = \det(\mathcal{H}_n)$ is a nonzero polynomial and $\mathcal{K}_n = \text{adj}(\mathcal{H}_n) \widetilde{\mathcal{M}}_n$. Moreover, $h_n(0) = 0$, $\forall n \in \mathbb{N}$, and $\deg(h_n) \leq 5$, $\forall n \geq 1$. From Theorem 3 it follows that h_n is independent of n . Thus, we obtain (29) with $zA = h_1$ and \mathcal{K}_n defined as above.

To obtain (30) we multiply (33) by zA and use (29). Thus, we obtain (30) with $\mathcal{M}_n = zA \mathcal{S}_n + \mathcal{T}_n \mathcal{K}_n$.

To prove assertion (a) we remind that Eqs. (29) can be written as equations of the same type as (20) and (21) of Theorem 3. Moreover, if

$$p = \max\{\deg(\mathcal{K}_n^{(1,1)}), \deg(\mathcal{K}_n^{(1,2)} + 1), \deg(\mathcal{K}_n^{(2,1)}), \deg(A - \mathcal{K}_n^{(2,2)})\}, \quad \forall n \in \mathbb{N},$$

then one can see that $p \leq 4$ and $\deg(A - \mathcal{K}_n^{(2,2)}) = p$, $n \geq 1$. Thus, from Theorem 3 we conclude that $\{P_n\}$ is semi-classical.

To prove assertion (b) we use an analogue argument as in the proof of Theorem 3. We write (30) in the form

$$zAR_n = G_n P_n + H_n P_n^*, \tag{35}$$

$$zAR_n^* = S_n P_n + T_n P_n^*, \quad n \geq 1, \tag{36}$$

with $G_n, H_n, S_n, T_n \in \mathbb{P}$. From the definition of u^{zA} and the Hermitian character of u , we have

$$\langle u^{zA}, R_n z^{-k} \rangle = \langle u, zAR_n z^{-k} \rangle + \overline{\langle u, zAR_n^* z^{k-n} \rangle}. \tag{37}$$

On the other hand, using (35) and (36) in (37) we get, for $n, k \geq 0$,

$$\langle u^{zA}, R_n z^{-k} \rangle = \langle u, G_n P_n z^{-k} \rangle + \langle u, H_n P_n^* z^{-k} \rangle + \overline{\langle u, S_n P_n z^{k-n} \rangle} + \overline{\langle u, T_n P_n^* z^{k-n} \rangle}. \tag{38}$$

Using a similar reasoning as in the proof of Theorem 3, we obtain for

$$p = \max\{\deg(G_n), \deg(H_n) + 1, \deg(S_n), \deg(T_n)\}, \quad \forall n \in \mathbb{N},$$

that

$$\langle u^A, R_n z^{-k} \rangle = 0 \quad \text{for every } p \leq k \leq n - p - 1 \text{ as well as for every } n \geq 2p + 1.$$

Thus the condition (i) of Definition 2 is satisfied.

Then, we can also establish that condition (ii) of Definition 2,

$$\exists n_0 \geq 2p : \langle u^A, R_{n_0} z^{-n_0+p} \rangle \neq 0$$

holds for $n_0 \geq 2p$ if and only if $\deg(T_{n_0}) = p$. Moreover, we get that $p \leq 6$ and $\deg(T_n) = p$, $\forall n \geq 1$.

Thus $\{R_n\}$ is quasi-orthogonal of order p with respect to the functional u^{z^A} . In this case, from Theorem 2, we conclude that there exists a polynomial B with $\deg(B) = p$ such that $u^{z^A} = v^B$. \square

5. Examples of coherent pairs on the unit circle

In this section we present the examples of coherent pairs corresponding to the Bernstein–Szegő class.

Theorem 5. *Let (μ_0, μ_1) be a coherent pair of measures supported on the unit circle. If μ_0 is the Lebesgue measure, then μ_1 belongs to the Bernstein–Szegő class, and the corresponding MOPS, $\{R_n\}$, is given by, $R_n(z) = z^{n-1}(z + c)$, $n \geq 1$, with c a constant, $|c| < 1$.*

Furthermore, $d\mu_1 = d\theta / (2\pi|z + c|^2)$.

Proof. If in (6) we assume the sequence $\{P_n\}$ is a classical Hahn MOPS in the sense that $\{P'_{n+1}/(n+1)\}$ is a sequence of monic polynomials orthogonal with respect to a measure supported on the unit circle, we know that $P_n(z) = z^n$ (see [7]). Therefore,

$$R_{n-1}(z) = z^{n-1} + \alpha_{n-1}z^{n-2}.$$

If we want that $\{R_n\}$ is a monic orthogonal polynomial sequence on the unit circle, then it will satisfy a forward recurrence relation

$$zR_{n-1}(z) + R_n(0)R_{n-1}^*(z) = R_n(z), \tag{39}$$

and so $\alpha_n = \alpha_{n-1} = \dots = \alpha_2 = c$. As a consequence,

$$R_n(z) = z^{n-1}(z + c).$$

Thus the MOPS $\{R_n\}$ belongs to the Bernstein–Szegő class and μ_1 is defined as stated (see [2], for example). \square

Theorem 6. *The only Bernstein–Szegő measure, μ_0 , that admits a companion measure μ_1 supported on the unit circle such that it yields a coherent pair, is the Lebesgue measure.*

Proof. Let (μ_0, μ_1) be a coherent pair of measures supported on the unit circle and $\{P_n\}, \{R_n\}$ the corresponding MOPS. We will prove that if P_n belongs to the Bernstein–Szegő class, then $P_n(z) = z^n$.

Let us suppose that the monic orthogonal polynomial sequence $\{P_n\}$ is defined by $P_n(z) = z^{n-k} P_k(z)$ for $n \geq k$ (for a fixed nonnegative integer number k), where P_k is a monic polynomial of degree k with zeros of absolute value less than 1 and such that $P_k(0) \neq 0$. Thus

$$P'_n(z) = (n - k)z^{n-k-1} P_k(z) + z^{n-k} P'_k(z).$$

From (6) it follows that

$$R_n(z) = z^{n-k-1} P_k(z) \left[\frac{n - k + 1}{n + 1} z + \alpha_n \frac{n - k}{n} \right] + z^{n-k} P'_k(z) \left[\frac{z}{n + 1} + \frac{\alpha_n}{n} \right].$$

Since $R_n(0) = 0$ for $n \geq k + 2$ and taking into account (39), we have

$$R_n(z) = z R_{n-1}(z), \quad n \geq k + 2.$$

Thus, for $n \geq k + 2$

$$P_k(z) \left[\left(\frac{n - k + 1}{n + 1} - \frac{n - k}{n} \right) z + \frac{n - k}{n} \alpha_n - \frac{n - k - 1}{n - 1} \alpha_{n-1} \right] + z P'_k(z) \left[\left(\frac{1}{n + 1} - \frac{1}{n} \right) z + \frac{\alpha_n}{n} - \frac{\alpha_{n-1}}{n - 1} \right] = 0. \tag{40}$$

Hence, taking into account that $P_k(0) \neq 0$, we get from (40) with $z = 0$,

$$\alpha_n = \frac{n}{(k + 1)(n - k)} \alpha_{k+1}, \quad n \geq k + 2.$$

Substituting this expression in (40),

$$k P_k(z) - P'_k(z) \left[z + \frac{n(n + 1)}{(k + 1)(n - k)(n - k - 1)} \alpha_{k+1} \right] = 0, \quad n \geq k + 2$$

then $\alpha_{k+1} = 0$, as well as $P_k(z) = z^k$. But this contradicts the fact $P_k(0) \neq 0$, up to $k = 0$. In such a case we are in the previous situation. So we obtain that $P_n(z) = z^n, n \in \mathbb{N}$. \square

Lemma 2. Let (u_n) be a sequence of complex numbers. If a sequence of monic polynomials $\{P_n\}$ orthogonal with respect to a linear functional v on the unit circle satisfies

$$\frac{z^n}{n} + u_{n-1} = \frac{P_n(z)}{n} + \alpha_{n-1} \frac{P_{n-1}(z)}{n - 1}, \quad n = 2, 3, \dots, \tag{41}$$

where we assume that $\alpha_{n-1} \neq 0, n = 2, 3, \dots$ then $u_n = 0, n = 1, 2, \dots$

Furthermore, the moments c_n , associated with v , are zero for $n = 2, 3, \dots$ and $c_1 \neq 0$.

Proof. Take $n = 2, 3, \dots$, multiply (41) by $1, 1/z, \dots, 1/z^{n-1}$, respectively, and use the linear functional v to get

$$\frac{c_n}{n} + u_{n-1} c_0 = 0, \tag{42}$$

$$\frac{c_{n-j}}{n} + u_{n-1} \bar{c}_j = 0, \quad j = 1, 2, \dots, n - 2, \tag{43}$$

$$\frac{c_1}{n} + u_{n-1} \bar{c}_{n-1} = \frac{\alpha_{n-1}}{n - 1} \langle v, P_{n-1}(z) \bar{P}_{n-1}(1/z) \rangle. \tag{44}$$

From (42) and (44) with $n = 2$ we get that $c_1 \neq 0$. From (42) to (44) with $n = 3, 4$ we get that $c_2 = c_3 = 0$ and, as consequence, $u_1 = u_2 = u_3 = 0$.

Now, we use induction arguments to conclude the proof, i.e. assuming $u_{k-1} = 0$ as well as $c_k = 0$ for $k = 2, 3, \dots, n - 1$, then from (43) we get that $u_{n-1} = 0$ and thus, from (42), $c_n = 0$. \square

Theorem 7. *Let (μ_0, μ_1) be a coherent pair of measures supported on the unit circle. If μ_1 is the Lebesgue measure then μ_0 must be an absolutely continuous measure*

$$d\mu_0 = |z - \alpha|^2 \frac{d\theta}{2\pi}, \quad z = e^{i\theta}.$$

Proof. If we assume μ_1 is the Lebesgue measure supported on the unit circle, i.e., $R_n(z) = z^n$, then (6) becomes

$$z^{n-1} = \frac{P'_n(z)}{n} + \alpha_{n-1} \frac{P'_{n-1}(z)}{n-1}, \quad n = 2, 3, \dots$$

Integrating the above expression, there exists a sequence of complex numbers (u_n) such that

$$\frac{z^n}{n} + u_{n-1} = \frac{P_n(z)}{n} + \alpha_{n-1} \frac{P_{n-1}(z)}{n-1}, \quad n = 2, 3, \dots$$

According to the previous lemma, the moments, c_n , associated with the linear functional, v , such that $\{P_n\}$ is the corresponding MOPS, satisfy $c_n = 0, n = 2, 3, \dots$, and c_0, c_1 are two complex arbitrary constants.

Furthermore, since v is a positive-definite linear functional associated with a positive Borel measure μ_0 supported on the unit circle, then we get an integral representation of such a functional taking into account its moments c_0 and c_1 . Indeed,

$$c_0 = \frac{A}{2\pi} \int_0^{2\pi} |z - \alpha|^2 d\theta, \quad c_1 = \frac{A}{2\pi} \int_0^{2\pi} z|z - \alpha|^2 d\theta,$$

with $z = e^{i\theta}$. Thus, $c_0 = (1 + |\alpha|^2)A, c_1 = -\alpha A$. In other words, $\frac{\alpha}{1 + |\alpha|^2} = -\frac{c_1}{c_0}$. \square

Acknowledgment

We thank the referees by their suggestions and remarks in order to improve the presentation of the manuscript.

References

[1] M. Alfaro, L. Moral, Quasi-orthogonality on the unit circle and semi-classical forms, Portugal. Math. 51 (1) (1994) 47–62.
 [2] A. Branquinho, L. Golinskii, F. Marcellán, Rational modifications of Lebesgue measure on the unit circle and an inverse problem, Complex Variables Theory Appl. 38 (2) (1999) 137–154.
 [3] A. Branquinho, M.N. Rebocho, Characterizations of Laguerre–Hahn affine orthogonal polynomials on the unit circle, J. Comput. Anal. Appl., to appear.
 [4] M. de Bruin, H.G. Meijer, Zeros of orthogonal polynomials in a non-discrete Sobolev space, Ann. Numer. Math. 27 (1995) 233–246.
 [5] T.S. Chihara, An Introduction to Orthogonal Polynomials, Gordon and Breach, New York, 1978.

- [6] A. Iserles, P.E. Koch, S.P. Norsett, J.M. Sanz-Serna, On polynomials orthogonal with respect to certain Sobolev inner products, *J. Approx. Theory* 65 (1991) 151–175.
- [7] F. Marcellán, P. Maroni, Orthogonal polynomials on the unit circle and their derivatives, *Constr. Approx.* 7 (1991) 341–348.
- [8] H.G. Meijer, Determination of all coherent pairs, *J. Approx. Theory* 89 (1997) 321–343.
- [9] B. Simon, *Orthogonal Polynomials on the Unit Circle*, American Mathematical Society Colloquium Publication, Part 1, vol. 54, American Mathematical Society, Providence, RI, 2005.